

## Review Article

# Effect of different spacing and nutrient management on growth and yield of maize: A Review

### Abstract

Optimal plant density influenced by crop geometric/planting techniques factors are crucial in achieving high yield to maize. In addition, nutrient management further enhances productivity. Increasing plant density can lead to higher grain yield per unit area, but it also reduces the yield per individual plant as the competition between plant to plant will increase. From different research it indicates that plant density lower than the optimum number result in higher productivity rate per plant but lower yield per area unit. After reviewing various research articles, it was concluded that the maximum growth and yield attributes of maize obtained with spacing 60: 20-30 cm. Highest nitrogen application up to 160 kg N per ha<sup>-1</sup> by split application at sowing, 25 DAS and tasselling is most reliable to increase the maize productivity. The combined application of 70 kg ha<sup>-1</sup> phosphorus and 60 kg ha<sup>-1</sup> potassium is the most suitable to obtain the maximum growth and yield attributes such as plant height, number of leaves per plant, cob length, cobs per plant, dry matter accumulation and crop growth rate. Effective nutrient management, aligned with proper planting density, played a pivotal role in maximizing maize productivity, meeting the demands of a growing population, and contributing to sustainable agriculture.

**Keywords:** *Nutrient management, production, spacing, sustainable agriculture.*

### Introduction

Maize (*Zea mays*) is a major cereal crop belonging to family Poaceae. Around 9000 years ago it was first cultivated in Mexico and was expanded throughout the Americas. The primary countries globally recognized for their maize production include the United States, China, regions of South and Central Africa, Argentina, Brazil, and Mexico. Within India, the states of Uttar Pradesh, Bihar, Rajasthan, Madhya Pradesh, Punjab, Karnataka, Himachal Pradesh, and Andhra Pradesh are significant for their cultivation of maize. The area of cultivation of this crop is increasing year after year. In the year 2020-21 maize was cultivated in 9.86 M ha and recorded production of 31.51 Mt of maize grains and a yield of 3195 kg/ha (Directorate of Economics and Statistics, 2021).

Maize used as dual-purpose crop i.e. food and feed purpose. Maize plant is C4 compact with defined nodes and internodes and having a fibrous root system with well-established prop roots or stilted root system which provides extra support to the plants and absorbing nutrients. The leaves are opposite spirally arranged and comprises of a sheath, collar (ligules, auricles) and a blade with white coloured mid-rib which runs through the length of the leaf, providing support to the leaf. Maize has monoecious flowers and the male flowers bloom in the tassel and female flowers bloom on the ear. Six varieties of maize are cultivated widely i.e., dent corn, flint corn, flour corn, pod corn, popcorn, and sweet corn. Maize is one of the cereals with the largest grain yield productivity but is also one to be affected by plant density fluctuations. Modern hybrids can typically sustain larger population densities because they can more easily manage adverse environmental circumstances induced by increased intra-specific competition, resulting in fewer barren plants and less stem lodging. Maize is widely grown in subtropical, tropical and temperate regions. It is known as the "Queen of Cereals" in the world. It is consumed as raw, roasted, cooked or table purposes. Maize is used in many ways like other grain crops in food dishes including "Chapati". The stalk and leaves given as a raw (green fodder) to animals or made into silage or hay (dry fodder) for long-term usage. The stalk and leaves are also used in the production of biodiesel, pigments, polymers and fibre. Maize grain has about 10% protein, 4% oil, 70% carbohydrates, 2.3% crude fibre, 10.4% albuminoids and 1.4% ash. Maize grain has significant quantities of vitamin A, nicotinic acid, riboflavin, vitamin E and low in calcium but high in phosphorus.

Maize yield is closely related to plant population. Productivity with relation to population density is a widely researched topic in the field of agronomy. The primary objective behind increasing the plant density is to improve the output, be it in the form of grain or biomass, thereby making the crop system more effective and competitive per unit area Chauhan and Opeña (2013) [7]. However, higher plant population increase yield because grain yield is directly proportional to the number of plants standing in field at the time of harvesting of the crop. Maize growth and yield is also impacted by different crop geometries, intercropping and nutrient application, Thavaprakash *et. al.*, (2005). [51] Distance between rows and the distance between plants in a row influence the plant population in a unit area. Sowing of maize crop at optimum distance eases field operations like fertilizer application and weeding, reduces plant competition for resources like sunlight, water, nutrients and reduces the risk of infestations and diseases. Weed management is also an important factor for cultivation of maize to reduce the yield losses and enhance the crop growth and development (Deewan *et. al.*, 2017). [10]

Nutrient management is a key factor in cultivation of crops to achieve higher yields and also maintain the soil health. To meet out the requirement of human as well as animals (fodder) required a lot of agronomic practices to enhance the production of maize. Maize has distinct requirements of nutrients at various stages of growth. Nitrogen and phosphorus are required in early growth stages for root and shoot development. Potassium is provided to ensure plant health and resilience against certain abiotic and biotic stresses. Nutrients are provided to the plants in various forms and sources such as organic manures and chemicals like compost, vermicompost, FYM, poultry manure, *etc* which also enhances the soil structure as well as the soil microbial activity and chemical compounds like

rock phosphate, wood ash or conventional forms of fertilizers like urea, MOP, DAP, SSP, TSP, etc respectively. Understanding the requirement of the crop and the available nutrient in the soil is necessary before application to prevent excess and provide a balanced and precise level of fertilizers. Significance of providing optimum distance and nutrient on various parameters i.e., growth, yield and quality of maize will be discussed as follows:

### **Effect of spacing on growth and yield of Maize.**

Maintaining distance of row to row and plant to plant is an important factor in the cultivation of crops as it significantly influences the growth and development of crops by ensuring efficient utilization of resources. Proper spacing provides better vegetative growth as well as rooting, it also reduces the risks of disease incidence by reducing excessive humidity and weeds which may serve as a host. Maize being a C<sub>4</sub> plant has proper growth and photosynthesis rate when adequate and recommended rate of spacing is practiced, it also reduces competition of essential nutrients and water resources. Maintaining a proper space also ensures ease of weed control such as mechanical and manual weeding. Among plants if there is improper spacing, it limits the rate of photosynthesis, depletion of soil nutrients, water stress, incidence of pests and diseases and difficulties in weed control. Ideal spacing of maize can be different according to the topography and the climate of the area. Recommended spacing of maize in Punjab region is 60 x 20 cm. It is also important to consider the recommended spacing according to the objectives of cultivation, be it for fodder production or for human consumption.

Plant population and spacing affects the biomass production and grain yield, Harhoff and Swanepoel, (2022) [19] employed two different row spacing levels @ 52 cm and 76 cm where productivity was higher in the treatment with larger width. This maybe due to the competition of resources amongst the plants during vegetative stages, wider row spacings can also reduce crop failure caused by climate related factors. Similarly, Manan *et. al.*, (2016) [29] reported that the spacing from narrow to wider distance increase the grain yield of the maize crop. The effects of combination of sowing methods and intra-row spacing was observed in plant growth attributes and the effects of sowing methods were highly significant at grain yield of the maize crop. The grain yield of 3.53 t ha<sup>-1</sup> was the highest yield obtained from ridge sowing method integrated with 20 cm intra-row spacing was suitable for the satisfactory growth of maize crop as conducted by Hamid *et. al.*, (2022) [20] as compared to drilling method integrated with 20 cm intra row spacing, terrace method combined with 30 cm intra-row spacing. According to Meena *et. al.*, (2022) [31], fodder maize grown at a wider row spacing of 40 cm have better growth in terms of foliage, growth and yield of maize. Alam (2020) [3] reported that the highest plant height, stem diameter and the length of leaf was highest with 60cm x 30cm in contrast to the 50cm x 20cm, 55cm x 25cm, 65cm x 35cm and 70cm x 40cm of maize variety Bari hybrid Butta-09. In the same year, Rina *et. al.*, (2020) [41] found that providing a spacing of 60 x 20 cm gives better values in terms of growth and yield attributes like harvest index, test weight, etc. Similarly, Mahato *et. al.*, (2020) [28] observed that maintaining spacing of 60 x 20 cm and maintaining the plant population contributes to grain per unit area i.e., maximum grain yield (7.5 t/ha). It was also observed that wider spacing causes drastic reduction in crop yield. As compared to Rina *et. al.*, (2023) [41] using

a spacing half the size they have experimented, Paul (2019)[37] concluded that using spacing distance of 35cm x 10cm gives the maximum production of green fodder. Sharifi and Namvar (2016) [45] found that the significantly taller plants, total dry matter was found in intra row spacing of 45cm. In the same year Devi and Ghosh (2017) [17] recorded that the best performance with spacing of 60 x 15cm on plant growth attributes, grain yield and stover yield.

Singh *et. al.*, (2010) [47] demonstrated that intercropping maize with lentils in a ratio of 1:2 produced highest cob length cob weight per plant. The choice of crop companions, ratio and the spacing significantly influence the overall yield. Similarly, El-Ghobashy *et al* (2018) [13] observed higher land use efficiency when maize and cowpea were planted in a ratio of 2:4 respectively. Babu and Senthivel [5] conducted an experiment in Kharif season of 2014 and 15 to assess the significance of plant spacing and weed management of maize where it is indicated that maintaining a spacing of 60 x 25 cm and pre-emergence spraying of atrazine and hand weeding at 30 DAS had a favourable impact on increasing grain yield. In accordance with these findings, it may be considered emphasizing on plant spacing and weed management as strategies of achieving optimum yield and profit. Maintaining optimum spacing and cropping geometry can be considered a sustainable method as it reduces the use of inputs which may be harmful to the crop ecosystem, human health and the environment if wrongly applied. Practicing a proper spacing and crop geometry improve the maize production system, thus giving a major emphasis in sustainable agriculture.

Rina *et. al.*, (2020) [41] reported that the combination treatment of spacing 60 x 20 cm with 80 kg/ha P on PSB treated seeds obtained the best results with higher results namely in grain yield (5.22 ton/ha), harvest index (43.13%) and test weight (28.33 g). According to these findings it may be considered that offering strategies of spacing and nutrient management techniques like using of biofertilizers may improve the maize production system giving a major emphasis in sustainable agriculture.

Kebede (2019) [24] reported that the highest stover yield of 31.29 t ha<sup>-1</sup> and grain yield of 11.67 t ha<sup>-1</sup> was obtained in the combination of 75 cm x 25 cm in 20cm. Therefore, he concluded that optimum inter-row and intra-row spacing combination of 75 cm x 25 cm gives the highest grain yield for both varieties (BH-661 and BH-QPY-545) under sufficient amount and proper distribution of rainfall. According to his findings, knowing the appropriate agronomic practices is also a key factor to maximize maize production. Gaire *et. al.*, (2020) [16] reported that maintaining a crop geometry effects on the grain yield of the crop. The highest grain yield (4.11 Mt/ha) can be achieved by maintaining a spacing of 60 x 15 cm. The LAI is also found to be higher in the 60 x 15 cm spacing.

In the same year (2020) Koirala *et. al.*, [26] revealed highest yield was recorded at 60 x 25 cm spacing practiced. After all the findings we can say that maintaining a row to row spacing of 60 cm is advisable in the cultivation of maize. During the kharif season of 2018, Ezung and Jamir (2019)[14] conducted an experiment by practicing two different spacing patterns *i.e.*, x 25 cm has significantly influenced the growth and yield of the crop. Nagarajan *et. al.*, (2019) [33] revealed that the highest grain yield was achieved in 60 x 25 cm spacing.

**Table 1:** Effect of different cropping geometry on growth and yield attributes of maize.

<b>Cropping geometry</b>	<b>Results obtained</b>	<b>References</b>
Ridge sowing method combined with 20 cm intra-row spacing.	The grain yield 3.53 t ha <sup>-1</sup> were obtained.	Hamid <i>et. al.</i> , (2022)
Row spacing width of 40 cm	Obtained better foliage, height, occurrence of internodes and stem girth.	Meena <i>et. al.</i> , (2022)
Row spacing of (60 x 30) cm	Performed best at this spacing interval in terms of growth and yield.	Alam (2020)
Spacing of (60 x 15) cm,	Obtained highest yield (4.11 Mt/ha), leaf area index.	Gaire <i>et. al.</i> , (2020)
Spacing of (60 x 20) cm	Obtained higher harvest index, test weight, significantly highest grain yield (7.5 t/ha).	Rina <i>et. al.</i> , (2020), Mahato <i>et. al.</i> , (2020)
Spacing of (60 x 25) cm	Obtained significant highest yield attributes and grain yield of 3.5 t/ha.	Koirala <i>et. al.</i> , (2020)
Spacing of (75 × 25) cm	Obtained highest stover yield of 31.29 t ha <sup>-1</sup> and grain yield of 11.67 t ha <sup>-1</sup> .	Kebede (2019)
Spacing of (60 x 25) cm	Obtained the maximum grain yield (7431 kg/ha).	Nagarajan <i>et. al.</i> , (2019)
Intra row spacing of 45cm.	Significant result of plant height, total dry matter was recorded.	Sharifi and Namvar (2016)
Spacing of (60 × 15) cm	Obtained better plant growth parameters like plant height, plant dry weight, grain yield and stover yield.	Devi and Ghosh (2016)
Spacing of (60 x 25) cm	Increased the yield of grain.	Babu and Senthivel (2014)
Intercropping maize with lentils in a ratio of 1:2	Produced highest cob length cob weight per plant. The	Singh <i>et. al.</i> , (2010)

	choice of crop companions, ratio and the spacing significantly influence the overall yield.	
--	---	--

### Effect of nutrient management on growth of maize.

Maize is a heavy feeder and required a lot of nutrients for its physio-morphological growth. Plant requires various nutrients to be provided. According to Ariraman *et. al.*,(2020) [4] application of nitrogen at a range of 150 to 200 kg/ha was the suitable dose for maize cultivation and resulting a better growth, yield and quality mainly protein content and fibre content. Hirpara *et. al.*, (2017) [35] revealed that 120 kg/ha of N and 60 kg/ha P resulted in highest plant height and in maize cultivation in South Saurashtra region. Sabu *et. al.*, (2021) [42] recorded that combined spraying of 75 kg/ha P + 40 kg/ha K as a basal treatment have produced most significant better results on various plant growth parameters like plant height (95.4 cm), number of leaves (10.83) and dry weight (77.33 g) over the other nutrient treatment in baby corn. Interaction effect of both the nutrients also significantly enhance the growth and yield attributing characters over the individual one fertilizer. Sakandiya and Sanodiya (2021) [43] resulted that treatment combination of 70 kg/ha P + 60 kg/ha K gave the maximum values of growth parameters, such as plant height, number of leaves, dry matter and crop growth rate over the nutrient treatment in maize cultivation. Khan *et. al.*,(2017) [25] reported that growth attributes such as plant height, number of leaves per plant, dry weight and crop growth rate were recorded higher with 60 kg/ha P through DAP in sweet corn variety Sweety as compared with other levels of phosphorus during the kharif season.

Ezung and Jamir (2020) [14] reported that the application of 100% recommended dose had a significant positive impact on the plant growth attributes of the maize like plant height at monthly intervals and leaf area index. Kumar and Mehera (2022) [27] concluded that the combined treatment of Azotobacter + KSB + Potassium @ 50 kg/ha gives highest values of growth and yield attributes namely, height of plant (222.6 cm) and crop growth rate (27.4 g/m<sup>2</sup>/day). Similarly, by exploiting the combined effects of biofertilizers and organic fertilizer with conventional fertilizers Singh *et. al.*,(2017)[47] recorded that the maximum growth was observed in application of 100% RDF than the treatment combination of azotobacter, vermicompost and NPK fertilizers. It was also statistically on par with treatment of 50% RDF + vermicompost + FYM + Azotobacter and treatment of 75% RDF + vermicompost. In a similar manner Dilshad *et al* (2010) [11] found that integrated use of NPK fertilizers and organic inputs namely FYM and biofertilizers can yield higher grain harvest as compared to their sole treatments, it was also concluded that integrated application of nutrients has more nutrient use efficiency and reduced ecological hazards as compared to using only mineral fertilizers in lesser fertile soil and rainfed conditions. Similarly, Priya *et. al.*, (2014) [39] concluded that the integrated use of recommended dose of NPK with poultry manure increase the grain yield of hybrid maize compared to the control treatment *i.e* without any nutrient application. Mahato *et. al.*, (2020)[28] revealed that the

significant highest values of growth parameters mainly plant height, leaf area index and dry matter accumulation was found in application of vermicompost + 75% RDF + ZnSO<sub>4</sub>. Singh *et. al.*, (2021) [48] states that application of 150 kg/ha N and 30 kg/ha Zn resulted in higher values in growth and yield of maize. Kannan *et al.* (2013) [22] concluded that integrated use of inorganic fertilizers along with vermicompost at resulted in the best performance for maize growth parameters like leaf area, plant height, and yield parameters like number of grains per cob, 100-seed weight, and grain yield. The integrated nutrient management practice including vermicompost and the recommended dose of NPK also showed the highest organic carbon content, bulk density, and porosity in the soil compared to other treatments. Thus, for enhancing maize productivity and improving soil fertility, integrated nutrient management can result in better yield and soil health.

### **Effect of nutrient management on yield of maize.**

Baby corns are small and immature corn cobs that are harvested before the kernels become mature and hardened. During the zaid season of 2021, Sabu *et. al.*, [42] performed an experiment to estimate the influence of phosphorous and potassium on the yield attributes of baby corn. It was concluded that application of 75 kg/ha P + 40 kg/ha K have performed better as compared to other treatments with yield and yield contributing characters like cob length (12.97 cm), cob weight (9.7 g), total yield (5214 kg/ha) and stover yield (20.18 t/ha). According to the findings of Gaire *et. al.*, (2020) [16], maize tends to show a significant increase in yield with applying of nitrogen levels @ 90 kg/ha and 120 ka/ha. Several yield attributes like length of cob, cob diameter, kernels per row and thousand grain weight was highest in application of nitrogen fertilizer at 90 kg/ha. Hirpara *et. al.*, (2017) [35] concluded that application of 120 kg/ha of N resulted in highest plant growth, yield attributing characters and straw yield of maize.

Ponmozhi *et. al.*, (2019) [38] resulted that the application of recommended dose of fertilizers, vermicompost, FYM, and zinc sulphate gives significantly higher impacts on dry matter accumulation, seed yield, straw yield and harvest index with the treatment combination of 100% RDF + 25% vermicompost + 25% FYM + 25 kg ZnSO<sub>4</sub>. Similarly, Singh *et. al.*, (2017) [47] found that the combined application of 75% RDF + 5 t/ha vermicompost + 5 t/ha FYM + Azotobacter contributes in maximum values of yield and yield attributes namely cobs per plant (2.11), cob length (19.2 cm), no. of grain/cob (399.7), no. of grain/row (28.5), shelling % (76.67%) seed index (30.45 g), grain yield (63.7 q/ha) and harvest index (29.58%). In an analogous manner to Ponmozhi *et. al.*, (2019) [38] by using combination of chemical fertilizers with vermicompost and micronutrients, Mahato *et. al.*, (2020) [16] estimated the impact of combined application of nutrient management on the plant growth of hybrid maize cultivation where highest values of yield parameters mainly grain yield of 9.04 t/ha and straw yield of 13.50 t/ha was found in treatment combination of vermicompost + 75% RDF + ZnSO<sub>4</sub>.

Nsanzabaganwa *et. al.*, (2014) [34] stated that winter grown maize is a heavy feeder and lacks proper nutrient management practices. They studied the impacts of applying N and P fertilizers as sole treatments and as interactions in maize cultivation where highest yield and yield contributing characters mainly grain yield, stover yield, harvest index, total biomass index and 1000 grain weight

recorded were at application of 240 kg/ha N and 26.4 kg/ha P. According to Khan *et al.*, (2017) [25] yield attributes such as cob weight, grains per row, rows per cob, grains per cob and 100 grain weight were recorded maximum with application of 60 kg/ha Phosphorus applied as DAP in Sweety variety of sweet corn. Patil and Basavaraja (2017) [36] concluded that highest grain and stover yields can be achieved by application of 125% K through potassium schoenite. Sankadiya and Sanodiya (2021) [43] conducted a study on a sandy loam soil which have low N content and medium content of P and K. Various yield attributes like cob per plant, cob length, no. of row per cob, no. of grains per row, grains per cob, grain yield, stover yield, seedweight and harvest index was found in application of 70 kg/ha (Phosphorus) + 60 Kg/ha (Potassium).

Hammad *et al.*, (2022) [21] experimented on the response of different levels of nitrogen on maize hybrids Syngenta 7720 and Muqabla S25W87. It was observed that highest N use efficiency was achieved in low N application, *i.e.*, 75 kg/ha and contrastingly highest grain yield, dry weight and grain protein content was observed in application of 300 kg/ha N. Singh *et al.*, (2017) [47] evaluate the impact of planting date and foliar application of  $KNO_3$  on grain yield and yield contributing characters of spring maize. It was found that foliar application of 1%  $KNO_3$  at development stage, specifically at tassel initiation enhanced the grain yield. Dadhich *et al.*, (2019) [8] revealed that application of 60 kg/ha K and 45 kg/ha Mg has the highest grain and stover yield of 2690 kg/ha and 6592 kg/ha for K treatment and 2831 kg/ha and 7000 kg/ha for magnesium treatment.

Kumar and Mehera (2022) [27] concluded that combined treatment of Azotobacter + KSB + Potassium @ 50 kg/ha produced highest values of growth and yield attributes namely, plant height (222.6 cm), dry weight (229.5 g), crop growth rate (27.4 g/m<sup>2</sup>/day), cob yield (1.6/plant), grain yield (7.97 t/ha), straw yield (11.5 t/ha), harvest index (40.91%) and 1000 seed weight (243.3 g). Similarly, Raghavendra *et al.*, (2018) [40] concluded in his study that application of 50% recommended dose of K + KSB gives highest yield contributing characters in maize, *i.e.*, length of cob, cob girth, cob weight and biological yield.

Nutrients may be applied through incorporation of organic compounds like compost, vermicompost, humic acid *etc.* By using vermicompost as a source of nitrogen, Ezung and Jamir (2019) [14] conducted an experiment on four different levels of vermicompost where it is found that application of 100% recommended dose gives a significant positive impact on the yield and yield attributes of the crop, like grains per cob, 1000 seed weight, grain yield and straw yield of maize.

**Table 2:** Effect of nutrient management on cultivation of maize as cereal and fodder.

Nutrient Management Practice	Significance	References
Application of 300 N kg/ha	Obtained highest grain yield, dry weight and grain protein content.	Hammad <i>et al.</i> , (2022)
The combined treatment of Azotobacter + KSB + Potassium @ 50 kg/ha	Obtained the significant values of growth and yield attributes namely, plant height and crop growth rate, cob yield, grain yield, stover yield, harvest index and test weight.	Kumar and Mehera (2022)

Application of Vermicompost + 75% RDF + ZnSO <sub>4</sub>	Remarkable results in plant height, LAI and dry matter content.	Mahato <i>et al.</i> , (2020)
Combined application of 75 kg/ha P + 40 kg/ha K as a basal treatment	Obtained significant better results on plant growth parameters.	Sabuet <i>et al.</i> , (2021)
Application of 70 kg/ha P + 60 kg/ha K	Obtained highest values of growth, such as plant height, number of leaves per plant, dry matter accumulation and crop growth rate.	Sakandiya and Sanodiya (2021)
Application of 150 kg/ha N and 30 kg/ha Zn	Obtained higher values in growth and yield of maize.	Singh <i>et al.</i> , (2021)
100% RDF + 25% vermicompost + 25% FYM + 25kg ZnSO <sub>4</sub>	Significant results on dry matter accumulation, grain yield, stover yield and harvest index	Ponmozhi <i>et al.</i> , (2019)
Application of 60 kg/ha P through DAP	Obtained better growth attributes like height, foliage, dry weight and CGR	Khan <i>et al.</i> , (2017)
Application of 120 kg/ha N	Obtained significant values regarding cob characteristics, dry matter, grain yield and stover yield.	Hirpara <i>et al.</i> , (2017)
75% RDF + 5 t/ha vermicompost + 5 t/ha FYM + Azotobacter	Increases yield attributes namely cob/plant, cob length, no. of grain/cob, no. of grain/row, shelling %, seed index, grain yield and harvest index.	Singh <i>et al.</i> , (2017)

## Conclusion

In conclusion, the vital importance of management of plant distance and nutrient application in maize cultivation cannot be overstated. Through a review of numerous studies and experiments, it is evident that the choice of spacing configurations and optimal application of nutrients and have a crucial impact on the overall success in crop production. Maintaining proper spacing ensures that maize plants efficiently utilize essential resources such as sunlight, water, and nutrients which in turn, leads to better vegetative growth, root development, and a reduced risk of diseases and weed infestations. Overall, the findings emphasize the need to practice proper crop geometry and precise nutrient management practices to optimize maize cultivation, enhance crop performance, and ensure better quality harvests. These insights are valuable for farmers and agricultural practitioners seeking to enhance maize production and contribute to food security.

## References

1. Acharya, N. R., Sah, S. K., Gautam, A. K., &Regmi, A. P. (2020). Site specific nutrient management and its effect on growth and yield of winter maize. *Journal of Bioscience and Agriculture Research*, 25(02), 2128-2136.
2. Adhikari, K., Bhandari, S., Aryal, K., Mahato, M., & Shrestha, J. (2021). Effect of different levels of nitrogen on growth and yield of hybrid maize (*Zea mays* L.) varieties. *Journal of Agriculture and Natural Resources*, 4(2), 48-62.
3. Alam M J, Uddin M A, Nahar, M K, Ali M Y and Ahmed K S (2020). Enhancement of maize productivity through using improved techniques of spacing. *Journal of Experimental Bioscience* 11(2), 27-34
4. Ariraman. R, Prabhakaran, J, Selvakumar, S, Sowmya, SandMansingh, M. D. I. (2020). Effect of nitrogen levels on growth parameters, yield parameters, yield, quality and economics of maize: A review. *Journal of Pharmacognosy and Phytochemistry*9(6), 1558-1563
5. Babu T Rand Senthivel T (2019). Yield and economics of hybrid maize as influenced by plant spacing and weed management practices. *Journal of Pharmacognosy and Phytochemistry*8(2S): 876-879
6. Barbieri P ARozas H N R S, Andrade F H and Echeverria H N E (2000). Row spacing effects at different levels of nitrogen availability in maize. *Agronomy Journal*, 92(2), 283- 288
7. Chauhan, B. S. &Opeña J. L. (2013). Effect of plant geometry on growth and yield of corn in the rice-corn cropping system. *American Journal of Plant Sciences*, 4(10), 1928.
8. Dadhich S K, Sharma S K, Yadav G K and Charry G R (2019). Effect of Potassium and Magnesium on Productivity of Maize (*Zea mays* L.) under Dryland Conditions. *International Journal of Current Microbiology and Applied Science* 8(4), 3072-3077
9. Devi H S and Ghosh G (2017). Effect of planting geometry, nitrogen levels and zinc application on growth and yield of hybrid maize (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry* 6(4), 1067-1069
10. Deewan, P., Mundra, S. L., Singh, D., Meena, M., Verma, R., & Sharma, N. K. (2017). Effect of weed and nutrient management on growth, productivity and protein content of quality protein maize (*Zea mays* L.). *Journal of Pharmacognosy and Phytochemistry*, 6(1), 271-274.
11. Dilshad, M. D. Lone, M. I. Jilani, G. Malik, M. A. Yousaf, M. Khalid, R., &Shamim, F. (2010). Integrated plant nutrient management (IPNM) on maize under rainfed condition. *Pakistan Journal of Nutrition*,9(9), 896-901.
12. Du, X., Wang, Z., Lei, W., & Kong, L. (2021). Increased planting density combined with reduced nitrogen rate to achieve high yield in maize. *Scientific Reports*, 11(1), 358.
13. El-Ghobashy, Y. E. Shams, A. S. &Lamlom, M. M. (2018). Maximizing land use efficiency by intercropping cowpea with some maize cultivars under different maize planting geometries. *Agricultural sciences*, 9(12), 1601-1620.

14. Ezung N K and Jami J. (2019). Study on growth, yield and economics in maize as influenced by different levels of nitrogen applied through vermicompost and spacing in hill areas of North-East Indian. *Journal of Soils and Crops*29(2), 236-242.
15. Fan, Y., Wang, Z. Liao, D. Raza, M. A. Wang, B. Zhang, J. & Yang, F. (2020). Uptake and utilization of nitrogen, phosphorus and potassium as related to yield advantage in maize-soybean intercropping under different row configurations. *Scientific Reports*, 10(1), 9504.
16. Gaire R, Pant C, Sapkota N, Dhamaniya R and Bhusal T (2020). Effect of Spacing and Nitrogen Level on Growth and Yield of Maize (*Zea mays* L.) in mid hill of Nepal. *Malaysia Journal of Halal Research*3(2): 50-55
17. Ghosh, D. Brahmachari, K. Brestic, M. Ondrisik, P. Hossain, A. Skalicky, M. & Bell, R. W. (2020). Integrated weed and nutrient management improve yield, nutrient uptake and economics of maize in the rice-maize cropping system of Eastern India. *Agronomy*, 10(12), 1906.
18. Golla, B. Mintesnot, A. & Getachew, M. (2020). Effect of nitrogen rate and intra-row spacing on yield and yield components of maize at Bako, Western Ethiopia. *African Journal of Agricultural Research*, 16(10), 1464-1471.
19. Haarhoff S J and Swanepoel P A (2022). Plant population and row spacing affects growth and yield of rainfed maize in semi-arid environments. *Frontiers in Plant Science* 13, 761121
20. Hamid A M A, Dagash Y M I, Osman S, Ali Y, Farah G A, Ahmed G A M and Ali O A. (2022). Effects of sowing methods and intra-row spacing on grain yield and some agronomic characters of maize (*Zea mays*). *American International Journal of Contemporary Research*, 12(1).
21. Hammad H. M., Chawla, M. S., Jawad, R, Alhuqail A, Bakhat, H. F, Farhad and Fahad S. (2022). Evaluating the impact of nitrogen application on growth and productivity of maize under control conditions. *Frontiers in Plant Science*, 13, 1402
22. Kannan, R. L., Dhivya, M., Abinaya, D., Krishna, R. L., & Krishnakumar, S. (2013). Effect of integrated nutrient management on soil fertility and productivity in maize. *Bulletin of Environment, Pharmacology and Life Sciences*, 2(8), 61-67.
23. Kandil, E. E. Abdelsalam, N. R. Mansour, M. A. Ali, H. M. & Siddiqui, M. H. (2020). Potentials of organic manure and potassium forms on maize (*Zea mays* L.) growth and production. *Scientific Reports*, 10(1), 8752.
24. Kebede M B. (2019). Effect of inter and intra row spacing on growth, yield components and yield of hybrid maize (*Zea mays* L.) varieties at Haramaya, Eastern Ethiopia. *American Journal of Plant Sciences*, 10(9), 1548-1564
25. Khan, W, Singh, V, Sagar, A and Singh, S. N. (2017). Response of phosphorus application on growth and yield attributes of sweet corn (*Zea mays* L. *Saccharata*) varieties. *Journal of Pharmacognosy and Phytochemistry* 6(5), 2144-2146

26. Koirala S, Dhakal A, Niraula D, Bartaula S, Panthi U and Mahato M (2020). Effects of row spacings and varieties on grain yield and economics of maize. *Journal of Agriculture and Natural Resources* 3(1), 209-218
27. Kumar K V and Mehera B (2022). Effect of bio-fertilizers and potassium on growth and yield of maize (*Zea mays* L.). *The Pharma Journal* 11(3), 2348-2351.
28. Mahato D C, Dutta D, Maity L, Biswas P, Mahato B, Ghosh Cand Patra S, (2020). Response of plant spacing and balanced fertilization on growth and yield of maize (*Zea mays* L.) in red laterite zone of Purulia district of West Bengal. *Journal of Experimental Agriculture* 42, 1-6.
29. Manan, J., Singh, G., Sharma, M., & Singh, G. (2016). Effect of plant spacing on yield and yield attributes of maize hybrids. *Journal of KrishiVigyan*, 5(1), 41-45.
30. Manasa, P., Maitra, S., & Barman, S. (2020). Yield attributes, yield, competitive ability and economics of summer maize-legume intercropping system. *International Journal of Agriculture, Environment and Biotechnology*, 13(1), 33-38.
31. Meena A, Solanki R M, Parmar P M and Chaudhari S. (2022). Effect of spacing and nitrogen fertilization on growth, yield and economics of fodder maize (*Zea mays* L.). *The PharmaJournal* 11(4) 1732-1735.
32. Nagarajan M. (2019). Performance evaluation of maize crop under different irrigation regimes and spacing. *International Journal of Agriculture Sciences* 0975-3710.
33. Nsanzabaganwa, E., Das, T. K., & Rana, D. S. (2014). Nitrogen and phosphorus effects on the growth, phenology, heat and nutrients accumulation and yield of winter maize (*Zea mays*) in western Indo-Gangetic Plains. *The Indian Journal of Agricultural Sciences*, 84(5), 661-4.
34. Pal B, Hirpara, D S, Vora, V D, Vekaria, P. D, Sutaria, G. S, Akbari, K Nand Verma. (2017). Effect of nitrogen and phosphorus on yield and yield attributes of maize in South Saurashtra, India. *International Journal of Current Microbiology and Applied Sciences* 6(3), 1945-1949
35. Patil Sand Basavaraja. (2017). Effect of different sources and levels of potassium on yield, nutrient requirement and nutrient use efficiency by maize crop (*Zea mays* L.) in low K soils of Eastern Dry Zone of Karnataka, India. *International Journal of Currennt Microbiology and Applied Sciences* 6(6), 193-199
36. Paul, N. C., Paul, U., Paul, S. C., & Paul, S. K. (2019). Effects of plant spacing and nitrogen level on the green fodder yield of maize (*Zea mays* L.). *Archives of Agriculture and Environmental Science*, 4(3), 307-312.
37. Ponmozhi, C. I, Kumar, R, Baba, Y. and A. Rao (2019). Effect of integrated nutrient management on growth and yield of maize (*Zea mays* L.). *International Journal of Current Microbiology and Applied Sciences* 8(11), 2675-2681
38. Priya, S. Kaushik, M. K. Sharma, S. K. & Priyanka, K. (2014). Impact of integrated nutrient management on growth and productivity of hybrid maize (*Zea mays* L.). *Annals of Biology*, 30(1), 106-108.

39. Raghavendra M, Singh Y V, Gaiind S, Meena, M. C and Das(2018). Effect of potassium and crop residue levels on potassium solubilizers and crop yield under maize-wheat rotation. *International Journal of Current Microbiological Applied Sciences* 7, 424-435
40. Rina L, Singh V and Dawson J. (2020). Effect of row spacing with different levels of phosphorous and biofertilizer on growth and yield of maize (*Zea mays* L.). *International Journal of Conservation Science* 8(3), 1171-1173
41. Sabu A, Dawson J and Thomas T. (2021). Effect of phosphorus and potassium levels on growth and yield of baby corn (*Zea mays* L.). *The PharmaInnovation* 10(10), 1959-1963.
42. Sankadiya S and Sanodiya L. (2021). Effect of phosphorus and potassium levels on growth and yield of maize (*Zea mays* L.). *Pharma Innovation Journal* 10(10), 1347-1350
43. Salem, E. M. Kenaway, K. M. Saady, H. S. & Mubarak, M. (2021). Soil mulching and deficit irrigation effect on sustainability of nutrients availability and uptake, and productivity of maize grown in calcareous soils. *Communications in Soil Science and Plant Analysis*, 52(15), 1745-1761.
44. Sharifi R S and Namvar A. (2016). Plant density and intra-row spacing effects on phenology, dry matter accumulation and leaf area index of maize in second cropping. *Biologija*, 62(1)
45. Singh N K, Rai B, Kumar R and Singh A K (2010). Response of winter maize (*Zea mays* L.)-based intercropping system under irrigated condition of Uttar Pradesh. *Environment and Ecology* 28(3), 1688-1690
46. Singh, L., Kumar, S., Singh, K., & Singh, D. (2017). Effect of integrated nutrient management on growth and yield attributes of maize under winter season (*Zea mays* L.). *Journal of pharmacognosy and Phytochemistry*, 6(5), 1625-1628.
47. Singh, S., Singh, H., Kumar, P., Singh, V., Kumar, S., & Singh, R. (2021). Effect of NPK levels with bio-fertilizers on productivity of maize (*Zea mays* L.). *International Journal of Chemical Studies*, 9(1), 1476-1479.
48. Tan, Y. Hu, F. Chai, Q. Li, G. Coulter, J. A. Zhao, C. & Yin, W. (2020). Expanding row ratio with lowered nitrogen fertilization improves system productivity of maize/pea strip intercropping. *European Journal of Agronomy*, 113, 125986.
49. Timsina, J. Kumar Singh, V. & Majumdar, K. (2013). Potassium management in rice–maize systems in South Asia. *Journal of Plant Nutrition and Soil Science*, 176(3), 317-330.
50. Thavaprakash, N. Velayudham, K. & Muthukumar, V. B. (2005). Effect of crop geometry, intercropping systems and integrated nutrient management practices on productivity of baby corn (*Zea mays* L.) based intercropping systems. *Research Journal of Agricultural and Biological Sciences*, 1(4), 295-302.
51. Yang, F. Wang, X. Liao, D. Lu, F. Gao, R. Liu, W. & Yang, W. (2015). Yield response to different planting geometries in maize–soybean relay strip intercropping systems. *Agronomy Journal*, 107(1), 296-304.