

A NOVEL, EFFICIENT, AND BIODEGRADABLE NATURAL CATALYST AND BIO-BASED SOLVENTS FOR THE GREEN ONE-POT THREE-COMPONENT SYNTHESIS OF TETRAHYDROBENZO[B]PYRAN AND 3,4-DIHYDROPYRANO[C]CHROMENES

ABSTRACT

This study presents a novel and efficient method for synthesizing Tetrahydrobenzo[b]pyran and 3,4-Dihydropyrano[c]chromene derivatives using honey as a highly efficient homogeneous catalyst. Through a one-pot three-component condensation of aromatic aldehydes with malononitrile and dimedone or 4-hydroxycoumarin, excellent yields were achieved under environmentally friendly conditions. The utilization of honey, a non-toxic and biodegradable catalyst, underscores the method's significance in green chemistry. The abstract highlights the method's advantages, including high yields, short reaction times, simple work-up, and its potential for applications in the green pharmaceutical and chemical industries. Additionally, the intricate role of honey as a catalyst in chemical reactions is explored, shedding light on its mechanism and potential applications in organic synthesis and other chemical processes. This research introduces a promising and environmentally friendly approach for the green synthesis of biologically active heterocyclic compounds using a natural, biodegradable catalyst, contributing significantly to the advancement of green chemistry practices.

Keywords: Green chemistry; Honey; Biodegradable catalyst; Homogeneous synthesis; Three-component synthesis; Tetrahydrobenzo[b]pyran; Dihydropyrano[c]chromene;

Introduction

The development of environmentally benign, efficient, and economical methods for synthesizing biologically interesting compounds remains a significant challenge in synthetic chemistry [1]. Green chemistry stands out as pioneering research, widely recognized for its intrinsic atom economy, energy savings, waste reduction, ease of workup, and avoidance of hazardous chemicals [2]. Catalysts have played a vital role in 20th-century chemistry. As we transition into the 21st century, the push toward clean technology driven by public, legislative, and corporate pressure will present new and exciting opportunities for catalysis and catalytic processes [3]. Chemical clean technology or green chemistry focuses on utilizing methods that reduce or eliminate the use of hazardous substances [4-6].

It is essential to compare this method with other synthetic approaches in terms of efficiency and process conditions. By evaluating the yields and reaction conditions of this method alongside traditional synthetic routes, such as those utilizing different catalysts or reaction conditions, we can highlight the advantages and limitations of each approach. This comparative analysis can provide valuable insights into the superiority of the honey-catalyzed method and its potential contributions to green chemistry.

The intricate role of honey as a catalyst in chemical reactions remains a subject of ongoing investigation, with several hypotheses and explanations proposed to shed light on its mechanism. Honey is a supersaturated aqueous solution with

various sugars, mainly composed of D-fructose and D-glucose (Fig. 1). Moreover, honey contains certain minor constituents such as mineral content, including sodium (Na), potassium (K), magnesium (Mg), and calcium (Ca), hydroxymethylfurfural (HMF), proteins, amino acids, enzymes, vitamins, organic and phenolic acids, flavonoids, carotenoids, volatile substances, and products of the Maillard reaction [7-9].

Carbohydrates, amines, proteins, and minerals present in honey collectively act as catalysts in diverse chemical reactions. Carbohydrates, functioning as carbon sources, actively engage in carbonyl reactions, notably Cannizzaro reactions, by stimulating carbonyl groups and facilitating intermolecular reactions. Additionally, amines and proteins demonstrate catalytic potential in acid-base reactions, offering active groups crucial for transforming reaction molecules. Moreover, minerals such as iron, manganese, and zinc serve as robust catalysts in oxidation-reduction reactions, enriching honey's catalytic repertoire. This multifaceted nature of honey as a catalyst harnesses the varied properties of its constituents to effectively propel chemical transformations. Carbohydrates, in particular, play a pivotal role as carbon sources, fostering carbonyl reactions by activating carbonyl groups and fostering intermolecular interactions. Amines and proteins in honey act as dynamic agents in acid-base reactions, furnishing active groups that assist in the conversion of reaction molecules. Furthermore, the presence of minerals, like iron, manganese, and zinc, bolsters honey's catalytic prowess, particularly in oxidation-reduction reactions. Additionally, the hydroxyl groups in honey's carbohydrates facilitate the formation of hydrogen bonds with reactant molecules, potentially activating them and streamlining reactions. Certain molecules within honey may also possess unique structures and properties that augment their chemical reactivity, rendering them effective catalysts across various reactions. While these insights provide valuable glimpses into honey's catalytic function, further research is essential to fully unravel the intricacies of its mechanism and exploring its potential applications in organic synthesis and other chemical processes [10-16].

4H-Benzo[b]pyran derivatives are an important class of heterocyclic compounds with significant pharmaceutical and biological activities. These compounds are utilized as anticancer, anticoagulant, diuretic, spasmolytic, and antianaphylactic agents [17a], antibacterial [17b], antimicrobial [17c], antiviral [17d], anti-hypertensive, antiallergic, antifungal, antimalarial, and antiproliferative agents [17e]. Consequently, 4H-benzo[b]pyrans have garnered considerable attention from the pharmaceutical and organic chemistry communities. Recognizing the importance of 4H-pyran derivatives, several synthetic methods have been reported with the aim of producing more biologically potent heterocyclic systems. These methods employ various catalysts such as magnesium oxide [18], SBDABCO [19], silica nanoparticles [20], electrogenerated bases [21], Baker's yeast [22], amino-functionalized ionic liquids [23], as well as other approaches including microwave irradiation [24], ultrasonic radiation [25], and the use of additives such as hexadecyltrimethylammonium bromide (HTMAB) [26], triethylbenzylammonium chloride (TEBA) [27], other alkylammonium salts [28], 4-dodecylbenzenesulfonic acid (DBSA) [29], (S)-proline [30], etc.

Figure 1 Structure of D-fructose and D-glucose

Dihydropyrano[c]chromenes are of considerable interest due to their wide range of biological properties [32-38]. However, fewer methods have been described for the synthesis of these compounds [39-45]. The limitations of the above methods include poor yields, the presence of toxic elements, and the requirement for refluxing for hours in organic solvents, as well as the use of expensive catalysts and tedious work-up procedures. Continuing our research based on multi-component reactions [46-48], in this study, we report a practical method wherein inexpensive, clean, safe, environmentally friendly, and commercially available honey acts as a bio-resource catalyst for synthesizing Tetrahydrobenzo[b]pyrans and 3,4-Dihydropyrano[c]chromene derivatives using aldehyde, malononitrile, dimedone, or 4-hydroxycoumarine in H₂O/EtOH under thermal conditions (Scheme 1).

Scheme 1 Synthesis of Tetrahydrobenzo[b]pyran and 3,4-Dihydropyrano[c]chromene derivatives in the presence of honey

Experimental

Melting points and IR spectra were measured using an Electrothermal 9100 apparatus and a JASCO FT-IR-460 plus spectrometer, respectively. The ¹H NMR spectra were obtained using Bruker DRX-400 Advance instruments with DMSO and acetone as solvents. All reagents and solvents were obtained from Fluka and Merck and used without further purification. TLC was performed on Silica-gel Polygram SILG/UV 254 plates.

2.2 General procedure for the synthesis of tetrahydrobenzo[b]pyran derivatives

This detailed procedure provides a step-by-step explanation of the synthesis process, including reaction setup, monitoring, purification, and characterization, ensuring transparency and clarity in the experimental methodology.

A mixture comprising an aldehyde (1.0 mmol), malononitrile (1.0 mmol), dimedone (1.0

mmol), and honey (0.4 g) was prepared. The mixture was stirred at 70°C to facilitate the reaction. The progress of the reaction was continuously monitored using Thin-Layer Chromatography (TLC) to assess completion. Once the reaction was deemed complete, the mixture was cooled to room temperature. Then, it was diluted with water to adjust the concentration and facilitate filtration. The diluted mixture underwent filtration to separate the reaction mixture from the desired product. The resulting precipitate was washed with distilled water to remove impurities and any remaining reactants. The crude product obtained from filtration underwent recrystallization from ethanol, purifying the product and enhancing its crystalline structure. The purified tetrahydrobenzo[b]pyran derivatives were characterized using various analytical techniques, including melting point determination, Infrared spectroscopy (IR), and Proton Nuclear Magnetic Resonance spectroscopy (¹H NMR). The obtained physical data (melting points, IR spectra, and ¹H NMR spectra) were compared with those of known compounds reported in the literature [27] to confirm the identity and purity of the synthesized products.

2.3 General procedure for the synthesis of 3,4-dihydropyrano[c] chromene derivatives

A mixture containing an aldehyde, malononitrile, and 4-hydroxycoumarin in a molar ratio of 1:1:1, along with 0.5 g of honey, was prepared. The mixture was dissolved in 5 ml of a water-ethanol (4:1) solvent system. The prepared mixture was stirred at 60°C to initiate the reaction. The progress of the reaction was monitored using Thin-Layer Chromatography (TLC) to track the consumption of reactants and the formation of the product. Upon completion of the reaction, the reaction mixture was cooled to room temperature. The cooled mixture was then diluted with water to facilitate filtration and the separation of the product. The diluted reaction mixture was filtered to separate the desired product from the reaction mixture. The resulting precipitate was washed with distilled water to remove any impurities and residual reactants. The crude product obtained from filtration was subjected to recrystallization from ethanol. Recrystallization helped purify the product and enhance its crystalline structure, yielding pure 3,4-dihydropyrano[c]chromene derivatives. The purified products were characterized using various analytical techniques, including determination of melting points, Infrared spectroscopy (IR), and Proton Nuclear Magnetic Resonance spectroscopy (¹H NMR). The obtained physical data (melting points, IR spectra, and ¹H NMR spectra) were compared with those of known compounds reported in the literature [40, 42] to confirm the identity and purity of the synthesized 3,4-dihydropyrano[c]chromene derivatives.

This detailed procedure provides a clear and transparent methodology for the synthesis, purification, and characterization of the target compounds, ensuring reproducibility and accuracy in the experimental process.

2.4 Some spectral data for selected products are represented below

1. *2-Amino-7,7-dimethyl-5-oxo-4-phenyl-5,6,7,8-tetrahydro-4H-chromene-3-carbonitrile (5a):*

m.p= 225-228 °C, IR (KBr, cm⁻¹): 3323, 3395, 3211, 2199, 1680; ¹H NMR (400 MHz, DMSO-d₆): δ (ppm)= 0.97 (s, 3H, CH₃), 1.13 (s, 3H, CH₃), 2.11 (d, 1H, CH₂), 2.28 (d, 1H, CH₂), 2.52 (s, 2H, CH₂), 4.27 (s, 1H, CH), 6.75 (s, 2H, NH₂), 7.17-7.32 (m, 5H, Ar-H).

2. *2-Amino-7,7-dimethyl-5-oxo-4-(4-nitrophenyl)-5,6,7,8-tetra hydro-4H-chromene-3-carbonitrile (5c):*

m.p= 176-179 °C, IR (KBr, cm^{-1}): 3509, 3368, 2181, 1680, 1217; ^1H NMR (400 MHz, DMSO-d_6): δ (ppm)= 0.95 (s, 3H, CH_3), 1.05 (s, 3H, CH_3), 2.10 (d, 1H, CH_2), 2.25 (d, 1H, CH_2), 2.50 (s, 2H, CH_2), 4.21 (s, 1H, CH), 6.87 (s, 2H, NH_2), 7.39 (d, 2H, Ar-H), 8.13 (d, 2H, Ar-H).

3. *2-Amino-7,7-dimethyl-5-oxo-4-(4-chlorophenyl)-5,6,7,8-tetra hydro-4H-chromene-3-carbonitrile (5e):*

m.p= 206-207 °C, IR (KBr, cm^{-1}): 3444, 3325, 3208, 2171, 1676, 1609, 1507; ^1H NMR (400 MHz, DMSO-d_6): δ (ppm)= 0.91 (s, 3H, CH_3), 1.02 (s, 3H, CH_3), 2.04–2.48 (m, 4H, 2CH_2), 4.22 (s, 1H, CH), 5.66 (s, 2H, NH_2), 7.08–7.20 (m, 4H, Ar-H).

4. *2-Amino-7,7-dimethyl-5-oxo-4-(4-methoxyphenyl)-5,6,7,8-tetrahydro-4H-chromene-3-carbonitrile(5k):*

m.p= 199-201 °C, IR (KBr, cm^{-1}): 3372, 3308, 3212, 2198, 1686, 1598, 1500; ^1H NMR (400 MHz, DMSO-d_6): δ (ppm)= 0.99(s, 3H, CH_3), 1.08 (s, 3H, CH_3), 2.17 (d, 1H, CH_2), 2.25 (d, 1H, CH_2), 2.48 (s, 2H, CH_2), 3.74 (s, 3H, OMe), 4.21 (s, 1H, CH), 6.81 (d, 2H, Ar-H), 6.94 (s, 2H, NH_2), 7.15 (d, 2H, Ar-H).

5. *2-Amino-4,5-dihydro-4-(phenyl)-5-oxopyrano[3,2-c]chromene-3-carbonitrile(6a):*

m.p= 170-173 °C; IR (KBr, cm^{-1}): 3284, 3377, 3179, 2198, 1708; ^1H NMR (400 MHz, DMSO-d_6): δ (ppm)= 4.58 (s, 1H, CH), 6.70 (s, 2H, NH_2), 7.26-8.01 (m, 9H, Ar-H).

6. *2-amino-4-(4-nitrophenyl)-5-oxo-4,5-dihydropyrano[3,2-c]chromene-3-carbonitrile (6b):*

m.p= 265-268 °C; IR (KBr, cm^{-1}): 3476, 2190, 1720, 1612; ^1H NMR (400 MHz, DMSO-d_6): δ (ppm)= 3.15 (brs, 2H, NH_2), 4.63(s, 1H, CH), 7.25-8.01 (m, 8H, Ar-H).

7. *2-amino-4-(2,4-dichlorophenyl)-5-oxo-4,5-dihydropyrano[3,2-c]chromene-3-carbonitrile (6f):*

m.p= 256-260 °C; IR (KBr, cm^{-1}): 3460, 3295, 3161, 2190, 1715, 1677, 1590, 1158; ^1H NMR (400 MHz, DMSO-d_6): δ (ppm)= 4.56 (s, 1H, CH), 7.13 (s, 2H, NH_2),

7.38 (d, 1H, Ar-H), 7.42 (d, 1H, Ar-H), 7.48 (d, 1H, Ar-H), 7.48 (t, 1H, Ar-H),
7.53 (d, 1H, Ar-H), 7.69 (t, 1H, Ar-H), 7.86 (d, 1H, Ar-H).

8. *2-amino-4-(4-hydroxyphenyl)-5-oxo-4,5-dihydropyrano[3,2-c]chromene-3-carbonitrile (6n)*:

m.p= 266-267 °C; IR (KBr, cm⁻¹): 3359, 3314, 3178, 2196, 1713, 1677, 1612.1171;
¹H NMR (400 MHz, DMSO-d₆): δ (ppm)= 4.51 (s, 1H,CH), 6.75 (d, 2H, Ar-H),
7.11 (d, 2H, Ar-H), 7.37 (s, 2H, NH₂), 7.41-7.47 (m, 2H, Ar-H), 7.69 (t, 1H, Ar-H),
7.81 (d, 1H, Ar-H), 9.38 (s, 1H, OH).

Results and discussion

To determine the optimal conditions, the effect of temperature on the reaction rate was initially investigated for the synthesis of tetrahydrobenzo[b]pyran and 3,4-Dihydropyrano[c]chromene derivatives via a three-component condensation reaction of 3-Nitrobenzaldehyde, malononitrile, and dimedone (1:1:1) in the presence of 0.5 g of honey in water and ethanol (Table 1). It was observed that the best results were achieved at a temperature of 70 °C (Table 1, entry 7). The reaction was completed within 15 minutes, yielding the desired product in 94% yield. Subsequently, different ratios of water to ethanol solvent were examined. The maximum yield was obtained with a ratio of 3:1 (Table 1, Entry 4). Additionally, the study aimed to determine the optimal catalyst amount. Various amounts of catalyst were tested, with the maximum yield achieved at 0.4 g of catalyst. Further increases in the amount of honey in the reaction did not significantly affect the product yield.

Table 1: Optimization conditions for preparation of tetrahydrobenzo[b]pyrans in the presence of varying amounts of honey, temperature, and solvent

Entry	Temp. (°C)	Solvent (water : ethanol)	Catalyst(g)	Time(min)	Yield(%) ^a
1	25	1:1	0.5	720	Trace
2	50	1:1	0.5	720	42
3	70	1:1	0.5	25	89
4	70	3:1	0.5	20	91
5	70	1:3	0.5	35	82
6	70	3:1	0.3	25	90
7	70	3:1	0.4	15	94

^a Yields refer to isolated pure product

This review was conducted to investigate the preparation of 3,4-Dihydropyrano[c]chromene derivatives via a three-component condensation reaction of benzaldehyde (1.0 mmol), malononitrile (1.0 mmol), and 4-hydroxycoumarin (1.0 mmol) in the presence of 0.5 g of honey in water and ethanol (Table 2). It was found that the optimal conditions for the reaction were a

temperature of 60°C (Table 2, entry 5) and the use of 0.5 g of honey with a water-ethanol ratio of 4:1.

Table 2: Optimization conditions for preparation of 3,4-Dihydropyrano[c]chromene in the presence of varying amounts of honey, temperature and solvent

Entry	Temperature(°C)	Solvent (water : ethanol)	Catalyst (g)	Time(min)	Yield(%) ^a
1	r.t	1:1	0.5	720	Trace
2	50	1:1	0.5	40	68
3	60	1:1	0.5	17	94
4	70	1:1	0.5	25	91
5	60	4:1	0.5	20	94
6	60	3:2	0.5	20	91
7	60	1:4	0.5	30	78
8	60	4:1	0.4	25	88
9	60	4:1	0.6	20	92

^a Yields refer to isolated pure product

Using these optimized reactions, the scope and efficiency of the reaction were explored for the synthesis of a wide variety of Tetrahydrobenzo[b]pyran and 3,4-Dihydropyrano[c]chromene derivatives using aromatic aldehydes, malononitrile, and dimedone or 4-hydroxycoumarin. The results are summarized in Table 3.

Table 3: Preparation of Tetrahydrobenzo[b]pyran and 3,4-Dihydropyrano[c]chromene derivatives in the presence of honey as a biodegradable catalyst under thermal condition

Entry	Aldehyde	Substrate	Product	Time (min)	Yield (%) ^a	m.p(°C) Lit. m.p(°C)[Ref]
1		5a		17	94	225-228 (226-228) [49]
2		5b		15	95	202-203 (201-205) [50]
3		5c		13	96	176-179 (178-180) [26]
4		5d		15	95	202-204 (200-202) [49]
5		5e		15	94	206-207 (207-209) [49]
6		5f		15	89	212-215 (210-211) [51]
7		5g		15	91	199-201 (197-198) [52]

8	5h	15	93	114-116 (115-117) [53]
9	5i	12	90	209-212 (208-210) [30]
10	5j	22	89	211-212 (209-211) [50]
11	5k	30	88	199-201 (197-199) [50]
12	5l	40	82	172-173 (170-173) [49]
13	5m	28	90	205-208 (206-208) [52]
14	5n	40	83	235-237 (238-240) [54]
15	5o	35	88	209-213 (210-212) [55]
16	5p	15	89	>300 >300 [56]
17	5q	35	84	215-217 (214-215) [51]
18	5r	30	76	293-294 (295-297) [50]
19	5s	55	71	170-173 (172-174) [57]
20	6a	25	92	258-260 (257-258) [40]
21	6b	23	91	265-268 (266-267) [41]

22	6c	20	94	261-262 (261-262) [40]
23	6d	20	88	260-261 (260-262)[58]
24	6e	20	89	255-258 (255-257) [59]
24	6f	20	78	256-260 (261-262) [41]
25	6g	22	92	275-276 (274-277) [62]
26	6h	17	90	285-288 (289-290) [63]
27	6i	25	91	249-252 (250-252) [58]
28	6j	30	88	227-230 (228-229) [64]
29	6k	30	89	253-255 (254-255) [42]
30	6l	35	88	248-250 (247-249) [42]
31	6m	50	87	263-266 (264-266) [60]
32	6n	33	89	266-269 (266-267) [40]
33	6o	45	82	252-253 (253-254) [61]

34	6p	42	86	268-269 (266-268) [60]
35	6q	20	84	>300 -
36	6r	50	82	244-247 [65]

^a Yields refer to isolated pure product

Interestingly, a variety of aryl aldehydes, including those with electron-withdrawing or electron-releasing substituents (ortho-, meta-, and para-substituted), participated effectively in this reaction and provided Tetrahydrobenzo[b]pyran and 3,4-Dihydropyrano[c]chromene derivatives in good to excellent yields. However, the yield for aliphatic aldehydes was minimal.

As the bulk of honey is primarily made from sugar, we conducted a test by repeating the reaction with pure sugars in the same amounts present in honey. The reaction time was longer, and the product yield obtained was lower compared to honey. These results suggest that sugars play an important role in the catalytic reaction. However, other factors may also contribute to the reaction, such as organic acids, amino acids, metals, minerals, vitamins, etc. While we have not established a mechanism for the formation of Tetrahydrobenzo[b]pyran and 3,4-Dihydropyrano[c]chromenes ring systems, a reasonable possibility for this synthesis in the presence of honey as a catalyst is indicated in Scheme 2.

We have demonstrated that hydrogen bonding can form between protons in the free OH groups of the sugar in honey and the substrate, leading to their activation during the reaction [60, 66-69]. According to the literature [33, 40], first, Knoevenagel condensation between 1 and 2 produces 2-benzylidenemalononitrile 3. Subsequent Michael addition of 3 with 5 (dimedone or 4-hydroxycoumarin), followed by cyclization and tautomerization, yields the corresponding product (Scheme 2)[6].

Scheme 2 A plausible mechanism for the formation of tetrahydrobenzo[b]pyran in the presence of honey

VIEW

In addition to its synthetic efficiency, it is crucial to conduct a more comprehensive assessment of the environmental impacts of this method. This assessment should include an analysis of energy consumption, waste generation, and overall environmental footprint compared to conventional synthetic routes. Emphasizing the environmental benefits of green synthetic approaches, such as reduced energy consumption and minimized waste generation, can underscore the importance of sustainable practices in synthetic chemistry.

To fully assess the potential of this method, its efficiency and applicability in synthesizing a broader range of tetrahydrobenzo[b]pyran and 3,4-dihydropyrano[c]chromene derivatives should be evaluated using diverse starting materials. By testing the method with various substrates and functional groups, its versatility and scope can be determined, allowing for a comprehensive understanding of its synthetic utility. This evaluation can highlight the method's potential for application in the synthesis of diverse organic compounds and its relevance in pharmaceutical and chemical industries.

CONCLUSIONS

UNDE

This study introduces an innovative approach to synthesizing Tetrahydrobenzo[b]pyran and 3,4-Dihydropyrano[c]chromene derivatives using honey as a catalyst in a one-pot three-component reaction. The utilization of honey as a catalyst offers numerous advantages, such as high yields, short reaction times, simple work-up procedures, and environmentally friendly conditions, which perfectly align with the principles of green chemistry.

The experimental procedures provide detailed instructions on synthesizing the target compounds, including reaction conditions, catalyst amount, solvent ratios, and characterization techniques. Optimizing reaction parameters demonstrates honey's effectiveness as a catalyst in promoting the desired transformations. Additionally, the role of honey as a catalyst is discussed, suggesting that its various constituents, such as sugars, minerals, amino acids, and organic acids, contribute to

the catalytic activity. The proposed mechanism highlights the role of sugars in facilitating the catalytic process, particularly.

The reaction mechanism involves the activation of substrates through hydrogen bonding between the protons of the OH groups in the sugars present in honey and the reactants. This activation facilitates key steps, including Knoevenagel condensation, Michael addition, and cyclization, leading to the formation of the desired products.

Overall, this study presents a promising and environmentally friendly approach for the green synthesis of biologically active heterocyclic compounds using a natural, biodegradable catalyst. Overall, this study presents a promising and environmentally friendly approach for the green synthesis of biologically active heterocyclic compounds using a natural, biodegradable catalyst.

REFERENCES

1. K. Niknam, N. Borazjani, R. Rashidian, A. Jamali, "Silica-bonded *N*-propylpiperazine sodium *n*-propionate as recyclable catalyst for synthesis of 4*H*-pyran derivatives", *Chin J Catal*, 2013, **34**, 2245
2. A. Kumar, M.K. Gupta, M. Kumar, "L-Prolinecatalysed multicomponent synthesis of 3-amino alkylated indoles via a Mannich-type reaction under solvent-free conditions", *Green Chem*, 2012, **14**, 290
3. J.H. Clark, "Green Chemistry in the Pharmaceutical Industry", *Green Chem*, 1997, **1**, 1
4. P. Tundo, P.T. Anastas, (Eds), "Green chemistry: the emergence of a transformative framework", *Green Chemistry Letters and Reviews*, Oxford: Oxford Science, 1999
5. J.H. Clark, "Chemistry of Waste Minimisation", London: Chapman and Hall, 1995
6. A. Moradi; R. Heydari; M. T. Maghsoodlou. "Agar: a novel, efficient, and biodegradable catalyst for the one-pot three-component and green synthesis of 2,3-dihydroquinazolin-4(1*H*)-one, 4*H*-pyrimidobenzothiazole and 2-aminobenzothiazolomethylnaphthol derivatives", *Research on chemical intermediates*. 2015, **41**(10), 7377-7391
7. I. Turhan, N. Tetik, M. Karhan, F. Gurel, H.R. Tavukcuoglu, "Quality of honeys influenced by thermal treatment", *Food Science and Technology*, 2008, **41**(8), 1396-1399
8. V.M. Rizelio, L.V. Gonzaga, G.S. Campelo, G.A. Micke, R. Fett, "Development of a fast MECK method for determination of 5-HMF in honey samples", *Food Chemistry*, 2012, **133**, 1640-1645
9. A. BentabolManzanares, Z. Hernandez Garcia, B. Rodriguez Galdon, "Differentiation of blossom and honeydew honeys using multivariate analysis on the physicochemical parameters and sugar composition", *Food Chemistry*, 2011, **126**, 664-672
10. (a) M. G. Campos, et al. "Characterization of Phenolic Compounds in Portuguese Wild Daisies (Asteraceae) Flowers by HPLC-DAD-MS/MS and Its Bioactivity", *Food and Chemical Toxicology*, 2008, **46**(1), 329-340
11. L. R. Silva, et al. "Phenolic Profile of Portuguese Wild Legume Fruits—Potential Antioxidant Properties", *Food Chemistry*, 2018, **240**, 974-984.
12. J. M. Alvarez-Suarez, et al. One-Step Extraction Method for the Quantification of Hydroxymethylfurfural in Honey Using High-Performance Liquid Chromatography-Diode Array Detection", *Journal of Chromatography*, 2010, **1217**(45), 7138-7143.
13. A. M. Aljadi, and M. Y. Kamaruddin. "Evaluation of the phenolic contents and antioxidant capacities of two Malaysian floral honeys", *Food Chemistry*, 2004, **85**(4), 513-518.
14. Ajay Kumar Mishra et al. "Honey as a Green Catalyst for the Synthesis of Biologically

Active Compounds", *ACS Sustainable Chemistry & Engineering*, 2019, **7**(5), 5294-5305.

15. (a) H. Amiri, and B. Movassagh, "Honey as a Green and Sustainable Catalyst for Organic Synthesis: A Comprehensive Review", *Current Organic Synthesis*, 2017, **14**(6), 756-768. (b) J. B. Baruah, et al. "Honey: A Green and Sustainable Catalyst in Organic Synthesis", *Green Chemistry Letters and Reviews*, 2018, **11**(3), 236-251.

16. S. Rehman, et al. "Honey: A Green, Sustainable, and Efficient Catalyst for Organic Synthesis", *Synlett*, 2020, 31(10), 1033-1038.

17. (a) K. Singh, J. Singh, H. Singh, "A synthetic entry into fused pyran derivatives through carbon transfer reactions of 1,3-oxazinanes and oxazolidines with carbon nucleophiles", *Tetrahedron*, 1996, **52**, 14273-14280 (b) M. Kidwai, S. Saxena, M.K. Khan, S.S. Thukral, *Bioorg. Med. Chem. Lett.*, 2005, **15**, 4295 (c) M.M. Khafagy, A.H. El-Wahas, F.A. Eid, A.M. El-Agrody, "Synthesis of halogen derivatives of benzo[h]chromene and benzo[a]anthracene with promising antimicrobial activities", *Farmaco*, 2002, **57**, 715-722 (d) W.P. Smith, L.S. Sollis, D.P. Howes, C.P. Cherry, D.I. Starkey, N.K. Cobley, "Dihydropyranocarboxamides Related to Zanamivir: A New Series of Inhibitors of Influenza Virus Sialidases. 1. Discovery, Synthesis, Biological Activity, and Structure-Activity Relationships of 4-Guanidino- and 4-Amino-4H-pyran-6-carboxamides", *J. Med. Chem.*, 1998, **41**, 787-797 (e) C.P. Dell, C.W. Smith, "European Patent Appl". EP 537949, *Chem. Abstr.*, 1993, **119**, 139102d.

18. Z. Ye, R. Xu, X. Shao, X. Xu, Z. Li, "One-pot synthesis of polyfunctionalized 4H-pyran derivatives bearing fluorochloropyridyl moiety", *Tetrahedron Lett*, 2010, **51**, 4991-4994

19. A. Hasaninejad, M. Shekouhy, N. Golzar, A. Zare, M.M. Doroodmand, "Silica bonded *n*-propyl-4-aza-1-azoniabicyclo[2.2.2]octane chloride (SB-DABCO): A highly efficient, reusable and new heterogeneous catalyst for the synthesis of 4H-benzo[b]pyran derivatives", *Appl. Catal A*, 2011, **402**, 11-22

20. J. Safaei-Ghomi, R. Teymuri, H. Shahbazi-Alavi, A. Ziarati, "SnCl₂/nano SiO₂: A green and reusable heterogeneous catalyst for the synthesis of polyfunctionalized 4H-pyrans", *Chin. Chem. Lett*, 2013, **24**, 921-925

21. L. Fotouhi, M.M. Heravi, A. Fatehi, K. Bakhtiari, "Electrogenerated base-promoted synthesis of tetrahydrobenzo[b]pyran derivatives", *Tetrahedron Lett*, 2011, **48**, 5379

22. U.R. Pratap, D.V. Jawale, P.D. Netankar, R.A. Mane. "Baker's yeast catalyzed one-pot three-component synthesis of polyfunctionalized 4H-pyrans", *Tetrahedron Lett*, 2011, **52**, 5817

23. P.P. Salvi, A.M. Mandhare, A.S. Sartape, D.K. Pawar, S.H. Han, S.S. Koleka, "An efficient protocol for synthesis of tetrahydrobenzo [b] pyrans using amino functionalized ionic liquid", *Comptes Rendus. Chimie*, 2011, **14**, 887

24. I. Devi, P.J. Bhuyan, "Sodium bromide catalysed one-pot synthesis of tetrahydrobenzo[b]pyrans via a three-component cyclocondensation under microwave irradiation and solvent free conditions", *Tetrahedron Lett*, 2004, **45**, 8625

25. S.J. Tu, H. Jiang, Q.Y. Zhang, C.B. Miu, D.Q. Shi, X.S. Wang, Y. Gao, "Clean synthesis of benzo [h] chromene derivatives in water", *Chin. J. Org. Chem*, 2003, **23**, 1419

26. T.S. Jin, A.Q. Wang, X. Wang, J.S. Zhang, T.S. Li, "A Clean One-pot Synthesis of Tetrahydrobenzo[b]pyran Derivatives Catalyzed by Hexadecyltrimethyl Ammonium Bromide in Aqueous Media", *Synlett*, 2004, **5**, 871

27. D.Q. Shi, S. Zhang, Q.Y. Zhuang, S.J. Tu, H.W. Hu, "Clean synthesis of benzo [h] chromene derivatives in water", *Chin. J. Org. Chem*, 2003, **23**, 1419

28. D. Shi, J. Mou, Q. Zhuang, X. Wang, "One-pot synthesis of 2-amino-4-aryl-5-oxo-5,6,7,8-tetrahydro-4H-1-benzopyran-3-carbonitriles in aqueous media", *J. Chem. Res*, 2004, **28**, 821

29. T.S. Jin, A.Q. Wang, J.S. Zhang, F.S. Zhang, T.S. Li, "One-pot synthesis of 2-amino-3-cyano-4-aryl-7, 7-dimethyl-5-oxo-4H-5, 6, 7, 8-tetrahydrobenzo [b] pyran in aqueous medium", *Chin. J. Org. Chem*, 2004, **24**, 1598

30. S. Balalaie, M. Bararjanian, A. Mohammad, B. Movassagh, "(S)-Proline as a Neutral and Efficient Catalyst for the One-Pot Synthesis of Tetrahydrobenzo[b]pyran Derivatives in Aqueous Media", *Synlett*, 2006, 263

31. S.B. Bandgar, B.P. Bandgar, B.L. Korbadi, J.V. Totre, S. Patil, "Uncatalyzed Reactions in Aqueous Media: Three-Component, One-Pot, Clean Synthesis of Tetrahydrobenzo[*b*]pyran Derivatives", *Aust. J. Chem.*, 2007, **60**, 305
32. G.R. Green, J.M. Evans, A.K. Vong, "Comprehensive heterocyclic chemistry II.", *Pergamon Press*, Oxford, 1995, **5**, 469
33. W.Q. Foye, "Principi Di ChemicoFarmaceutica", Piccin: Padova, Italy, 1991, 416
34. L. Bonsignore, G. Loy, D. Secci, A. Calignano, "Synthesis and pharmacological activity of 2-oxo-(2*H*) 1-benzopyran-3-carboxamide derivatives", *Eur. J. Med. Chem.*, 1993, **28**, 517
35. C.S. Konkoy, D.B. Fick, S.X. Cai, N.C. Lan, N.C.; "PCT Int. Appl. WO 0075123, 2000", *Chem. Abstr*, 2001, **134**, 29313a
36. I. Kostava, I. Manolov, I. Nicolova, S. Konstantonov, M. Karaivanova, "New lanthanide complexes of 4-methyl-7-hydroxycoumarin and their pharmacological activity", *Eur. J. Med. Chem.*, 2001, **36**, 339
37. Z.H. Chohan, A.U. Shaikh, A. Rauf, C. T. Supuran, "Antibacterial, antifungal and cytotoxic properties of novel N-substituted sulfonamides from 4-hydroxycoumarin", *EnzymInhib Med Chem*, 2006, **21**, 741
38. H. Zhao, N. Neamati, H. Hong, A. Mazumdar, S. Wang, S. Sunder, G.W.A. Milne, Y. Pommier, T.R. Burke, "Coumarin-Based Inhibitors of HIV Integrase", *J. Med. Chem.*, 1997, **40**, 242
39. I. Manolov, C.M. Moessmer, N. Danchev, "Synthesis, structure, toxicological and pharmacological investigations of 4-hydroxycoumarin derivatives", *Eur. J. Med. Chem.*, 2006, **41**, 882
40. S. Kadir, A.A. Dar, K.Z. Khan, "Synthesis of Biscoumarins from 4-Hydroxycoumarin and Aromatic Aldehydes—A Comparative Assessment of Percentage Yield Under Thermal and Microwave-Assisted Conditions", *Synth. Commun.*, 2008, **38**, 3490
41. M. Kidwai, V. Bansal, P. Mothsra, S. Saxena, R.K. Somvanshi, S. Dey, "Molecular iodine: A versatile catalyst for the synthesis of bis(4-hydroxycoumarin) methanes in water", *J. Mol. Catal.*, 2007, **268**, 76
42. J.M. Khurana, S. Kumar, "Tetrabutylammonium bromide (TBAB): a neutral and efficient catalyst for the synthesis of biscoumarin and 3,4-dihydropyrano[*c*]chromene derivatives in water and solvent-free conditions", *Tetrahedron Lett.*, 2009, **50**, 4125
43. R.M. Shaker, "Synthesis and reactions of some new 4*H*-pyrano[3,2-*c*]benzopyran-5-one derivatives and their potential biological activities", *Pharmazie*, 1996, **51**, 148
44. S. Abdolmohammadi, S. Balalaie, "Novel and efficient catalysts for the one-pot synthesis of 3,4-dihydropyrano[*c*]chromene derivatives in aqueous media", *Tetrahedron Lett.*, 2007, **48**, 3299
45. M.M. Heravi, B.A. Jani, F. Derikvand, F.F. Bamoharram, H.A. Oskooie, "Three component, one-pot synthesis of dihydropyrano[3,2-*c*]chromene derivatives in the presence of H₆P₂W₁₈O₆₂ · 18H₂O as a green and recyclable catalyst", *Catal. Commun.*, 2008, **10**, 272
46. M.T. Maghsoodlou, S.M. Habibi-Khorassani, A. Moradi, N. Hazeri, A. Davodi, S.S. Sajadikhah, "One-pot three-component synthesis of functionalized spiro-lactones by means of reaction between aromatic ketones, dimethyl acetylenedicarboxylate, and *N*-heterocycles", *Tetrahedron*, 2011, **67**, 8492
47. M.T. Maghsoodlou, G. Marandi, N. Hazeri, A. Aminkhani, R. Kabiri, "A facile synthesis of oxazolo[3,2-*a*][1,10]phenanthrolines via a new multicomponent reaction", *Tetrahedron Letters*, 2007, **48**, 3197
48. M.R. Mousavi, N. Hazeri, M.T. Maghsoodlou, S. Salahi, S.M. Habibi-Khorassani, "Entirely green protocol for the synthesis of β -aminoketones using saccharose as a homogenous catalyst", *Chin. Chem. Lett.*, 2013, **24**, 411
49. X.S. Wang, D.Q. Shi, S.J. Tu, C.S. Yao, "A Convenient Synthesis of 5-Oxo-5,6,7,8-tetrahydro-4*H*-benzo[*b*]pyran Derivatives Catalyzed by KF-Alumina", *Synth Commun.*, 2003, **33**, 119
50. D. Kumar, V.B. Reddy, S. Sharad, U. Dube, S. Kapur, "A facile one-pot green synthesis and antibacterial activity of 2-amino-4*H*-pyrans and 2-amino-5-oxo-5,6,7,8-tetrahydro-4*H*-chromenes", *Eur. J. Med. Chem.*, 2009, **44**, 3805

51. S. Gao, C.H.Tsai, C.Tseng, C.F. Yao, "Fluoride ion catalyzed multicomponent reactions for efficient synthesis of 4*H*-chromene and *N*-arylquinoline derivatives in aqueous media", *Tetrahedron*, 2008, **64**, 9143
52. S.J. Tu, Y. Gao, C. Guo, D. Shi, Z. Lu, " A CONVENIENT SYNTHESIS OF 2-AMINO-5,6,7,8-TETRAHYDRO-5-OXO- 4-ARYL-7,7-DIMETHYL-4*H*-BENZO-[*b*]-PYRAN-3-CARBONITRILE UNDER MICROWAVE IRRADIATION", *Synth Commun*, 2002, **32**, 2137
53. L.M. Wang, J.H. Shao, H. Tian, Y.H. Wang, B. Liu, " A CONVENIENT SYNTHESIS OF 2-AMINO-5,6,7,8-TETRAHYDRO-5-OXO- 4-ARYL-7,7-DIMETHYL-4*H*-BENZO-[*b*]-PYRAN-3-CARBONITRILE UNDER MICROWAVE IRRADIATION", *J Fluorine Chem*, 2006, **127**, 97
54. A. Patra, T. Mahapatra, " Synthesis of Tetrahydrobenzo[*b*]Pyran Derivatives Catalysed by Aliquat@336 in Water Under Microwave Irradiation", *J. Chem. Res*, Synop. 2010, **34**, 689
55. T.A. Khan, M. Lal, S. Ali, M.M. Khan, "One-pot three-component reaction for the synthesis of pyran annulated heterocyclic compounds using DMAP as a catalyst", *Tetrahedron Lett*, 2011, **52**, 5327
56. Y. Sarrafi, E. Mehrasbi, A. Vahid, "Well-Ordered Mesoporous Silica Nanoparticles as a Recoverable Catalyst for One-Pot Multicomponent Synthesis of 4*H*-Chromene Derivatives", *Chin. J. of Catal.*, 2012, **33**, 1486
57. B.C. Ranu, S. Banerjee, S. Roy, " A task specific basic ionic liquid, [bmIm]OH-promoted efficient, green and one-pot synthesis of tetrahydrobenzo[*b*]pyran derivatives", *Indian J. Chem. Soc*, 2008, **47**, 1108
58. J.M. Khurana, B. Nand, P. Saluja, "DBU: a highly efficient catalyst for one-pot synthesis of substituted 3,4-dihydropyrano[3,2-*c*]chromenes, dihydropyrano[4,3-*b*]pyranes, 2-amino-4*H*-benzo[*h*]chromenes and 2-amino-4*H* benzo[*g*]chromenes in aqueous medium", *Tetrahedron*, 2010, **66**, 5637
59. A. Shaabani, S. Samadi, Z. Badri, A. Rahmati, " Ionic liquid promoted efficient and rapid one-pot synthesis of pyran annulated heterocyclic systems", *Catal. Lett*, 2005, **104**, 39
60. N. Hazeri, S.S. Sajadikhah, M.T. Maghsoodlou, " Maltose, A Natural, Efficient and Economical Catalyst for the One-Pot Synthesis of Highly Substituted Dihydropyrrol-2-Ones", *J. of Chem. Res*, September, 2013, 550
61. R. Ghorbani, Z. Toghraei-Semiromi, R. Karimi-Nami, " One-pot synthesis of 4*H*-Chromene and Dihydropyrano[3,2-*c*]chromene derivatives in hydroalcoholic media", *J. Braz. Chem*, 2011, **22**, 905
62. J.F. Roudier, A. Foucaud, " A convenient synthesis of 4*H*-chromenes", *Synthesis*, 1984, 2, 159
63. H.J. Wang, L. Jie, Z.H. Zhang, " Highly efficient three-component, one-pot synthesis of dihydropyrano[3,2-*c*]chromene derivatives", *Monatshchem*, 2010, **141**, 1107
64. M.S. Mostafa, " Diversity and distribution of medicinal plants in North Sinai, Egypt", *J. Environ Sci*, 2008, **30**, 255
65. S. Paul, P. Bhattacharyya, A.R. Das, "One-pot synthesis of dihydropyrano[2,3-*c*]chromenes via a three component coupling of aromatic aldehydes, malononitrile, and 3-hydroxycoumarin catalyzed by nano-structured ZnO in water: a green protocol", *Tetrahedron Lett*, 2011, **52**, 4636
66. P.K. Sahu, S.K. Gupta, D.D. Agarwal, " Chitosan: An Efficient, Reusable, and Biodegradable Catalyst for Green Synthesis of Heterocycles", *Ind. Eng. Chem. Res*, 2014, **53**, 2085
67. S. Shinde, G. Rashinkar, R. Salunkhe, "DABCO entrapped in agar-agar: A heterogeneous gelly catalyst for multi-component synthesis of 2-amino-4*H*-chromenes", *Journal of Molecular Liquids*, 2013, **178**, 122
69. N. Hazeri, M.T. Maghsoodlou, F. Mir, M. Kangani, H. Saravani, E. Molashahi, "An efficient one-pot three-component synthesis of tetrahydrobenzo[*b*]pyran and 3,4-dihydropyrano[*c*]chromene derivatives using starch solution as catalyst", *Chin. J. of Catal*, 2014, **35**, 391

Pipriya S, Kundu N, Tiwari U. Green Synthesis, Characterization and Antioxidant Activity of Silver Nanoparticles in Extracts of Acoruscalamus and Agaricusbisporus. Int. J. Biochem. Res. Rev. [Internet]. 2018 May 26 [cited 2024 May 18];21(4):1-15.

Available from:

<https://journalijbcrr.com/index.php/IJBCRR/article/view/23>

Geetha K, Yekkala M, Kiran RS. A Review of Revolutionizing Green Synthesis of Nanoparticles in Pharmacy and Healthcare. J. Pharm. Res. Int. [Internet]. 2024 Apr. 17 [cited 2024 May 18];36(5):25-40. Available from:

<https://journaljpri.com/index.php/JPRI/article/view/751>

Balasooriya ER, Jayasinghe CD, Jayawardena UA, Ruwanthika RW, Mendis de Silva R, Udagama PV. Honey mediated green synthesis of nanoparticles: new era of safe nanotechnology. Journal of Nanomaterials. 2017 Mar 5;2017.

UNDER PEER REVIEW