

Original Research Article

INSECTICIDAL ACTIVITY OF DIFFERENT DOSES OF *ACORUS CALAMUS* ESSENTIAL OIL AGAINST *SITOPHILUS ORYZAE*

Abstract

The toxicity of different doses (3, 4, 5, 6 and 7 percent) of *Acorus calamus* essential oil (EO), against insect pest *Sitophilus oryzae* was evaluated in the laboratory of the Division of Entomology, FoA, Wadura, SKUAST-Kashmir during the year 2022. The various concentrations of *Acorus calamus* essential oil tested had significant effect on the mortality of *Sitophilus Oryzae*. 12 Hours After Treatment (HAT), the highest mortality of 74.27 percent was recorded in 7 percent concentration of *Acorus A. calamus* essential oil on EO 12 Hours After Treatment (HAT) followed by concentrations of 6 (71.01%), 5 (67.15%) and 4 (39.11%) percent concentration while untreated control recorded 5.13 percent mortality respectively. However, on 24 HAT, the cumulative mortality increased to 76.35 in 7 percent, 74.11 in 6 and 5 percent, and to 43.01 in 4 percent concentrations. On 72 HAT the concentrations of 7, 6 and 5 recorded 94.11, 91.27 and 90.11 percent mortality respectively, while only 7.05 percent mortality was recorded in untreated control.

Formatted: Font: Italic

Comment [H1]: Why respectively?!

Formatted: Highlight

Keywords: *Acorus calamus*, Essential oil, Mortality, Rice weevil

1. Introduction

Stored grain insect pests can cause reductions in weight, quality, commercial value and seed viability. Seventy-five percent of these insects are coleopterans [1] and the most damaging species of storage insects are in the genera *Sitophilus* and *Tribolium* [2, 3]. *Sitophilus oryzae* L. (Coleoptera: Curculionidae), a ubiquitous pest of economic importance, is an internal feeding insect that bores into stored grain. Adult weevils feed mainly on the endosperm, reducing the carbohydrate content and the larvae feed preferentially on the germ of the grain, thus removing a large percentage of the protein and vitamins. Insects that selectively attack the germ will cause a greater loss in germination than others. The control of arthropod pests on stored products has been primarily through the use of fumigants and residual chemical insecticides to augment the more obvious approach of hygiene [4, 5]. The excessive use of conventional chemical insecticides has resulted in a number of serious problems, e.g. resistance to the chemical insecticides, elimination of economically beneficial

insects, persistence in the environment, toxicity to humans and wildlife and higher cost of crop production [6]. Numerous insects and mites have developed the ability to withstand nearly all pesticides used for their management due to cross-resistance and multiple resistance mechanisms [7]. Recognition of the deleterious effects of pesticides has spurred the exploration of alternative, less intrusive management approaches, such as employing the use of essential oils.

Many medicinal plants and spices have been used as pest control agents [8, 9]. Farmers and researchers often claim the successful use of plant materials in insect pest control, including ash [10, 11], vegetable oils [12], plant extracts [13] and botanical powders [14]. It has been reported that certain plant preparations and traditional methods are much safer than chemical insecticides [15]; therefore, these materials should be explored to protect stored products against pest infestation. *Acorus calamus* L. (Araceae), commonly known as “sweet flag”, is a well-known ethnomedicinal and ethnobotanical plant worldwide. *Acorus* rhizome possesses a wide range of pharmacological activities, such as sedative, central nervous system depressant, behaviour modifying, anticonvulsant, memory enhancing, anti-inflammatory, antioxidant, antispasmodic, cardiovascular, hypolipidemic, and immunosuppressive. *Acorus calamus* L. essential oil have also been used as fumigant against stored food products [16]. Keeping in view the insecticidal properties of essential oil of *Acorus calamus*, the present study was carried out to evaluate its various concentrations against rice weevil.

2. Material and methods

2.1 Raising of plant material

The seedlings of *A. calamus* which were raised in polybags were procured from Regional cum Facilitation Centre North Zone II located at Faculty of Agriculture, Wadura Sopore and transplanted in the field with plant to plant and row to row distance of 15cm and 30cm, respectively. The rhizomes of mature plants were harvested, shade dried and ground to a fine powder and sieved through 30mm mesh. The powdered material was put to hydrodistillation for extraction of essential oil using Clevenger’s apparatus. The essential oil of *A. calamus* thus obtained was collected in the vials and stored in refrigerator for further use.

2.2 Preparation of different concentrations of essential oil of *A. calamus*

Comment [H2]: Amber color vials ?

Acetone was used as solvent for preparation of different concentrations. The doses 30, 40, 50, 60 and 70 μ l corresponding to 3, 4, 5, 6 and 7 per cent were used for determining toxicity of *A. calamus* essential oil against *Sitophilus oryzae*.

Comment [H3]: hadn't you treated the seeds of the control with acetone since you had used it for dilution?

2.3 Rearing of *Sitophilus oryzae*

The rearing of *S. oryzae* was carried in the laboratory of Division of Entomology, FoA, Wadura, SKUAST-Kashmir. The rice grains infested with *S. oryzae*, was brought from different godowns/shops of Kashmir. Rice grains were taken in 5 litre capacity rearing jars and placed in hot air oven at 60°C for 24 hours for sterilization. The adult insects (both male and female) were collected randomly from the infested grains with the help of aspirator and transferred to the respective sterilized food placed in jars (5 litre capacity). The jars were covered with muslin cloth in order to supply adequate humidity to the grains and kept in B.O.D incubator at 28.5 \pm 2°C and 65 \pm 5% relative humidity to ensure proper egg laying and maintaining culture of rice weevil for further investigation. Mortality per cent was recorded after 12, 24, 36, 48, 60 and 72 hours after treatment. Test insects were considered dead if appendages do not move when prodded with a fine brush.

$$\text{Mortality (\%)} = \frac{\text{No. of insects dead} \times 100}{\text{Total No. of insects fumigated}}$$

2.4 Data analysis

The data recorded in different treatments were subjected to analysis of variance (ANOVA) using R software.

3. Results and Discussion

The insecticidal activity of Sweet flag against rice weevil is shown in Table 1. The highest mortality of 74.27 percent was recorded in 7 percent concentration of Sweet Flag on 12 Hours After Treatment (HAT) followed by 6 (71.01%), 5 (67.15%) and 4 (39.11%) percent concentration while untreated control recorded 5.13 percent mortality respectively (Table 1). However, on 24 HAT the cumulative adult mortality increased to 76.35 in 7 percent, 74.11 in 6 and 5 percent, 43.01 in 4 percent concentration (Table 1). On 72 HAT the concentrations of 7, 6 and 5 recorded 94.11, 91.27 and 90.11 percent mortality respectively, while only 7.05 percent mortality was recorded in untreated control.

Table 1: Effect of different concentrations of *Acorus calamus* essential oil against rice weevil

Concentration (%)	Mortality (%) (HAT)						Mean
	12	24	36	48	60	72	
3	28.01 ^k (5.34)	33.03 ^{jk} (5.79)	39.11 ^{ijk} (6.29)	41.01 ^{ijk} (6.44)	42.13 ^{hij} (6.53)	43.05 ^{hij} (6.60)	37.72±2.42 (6.18)
4	39.11 ^{ijk} (6.29)	43.01 ^{ijk} (6.60)	45.03 ^{hij} (6.75)	46.13 ^{hi} (6.83)	47.11 ^{ghi} (6.90)	49.15 ^{fgh} (7.05)	44.92±1.43 (6.74)
5	67.15 ^{efg} (8.22)	74.11 ^{cdef} (8.64)	81.13 ^{abcde} (9.03)	88.03 ^{abc} (9.41)	89.01 ^{abc} (9.46)	90.11 ^{ab} (9.52)	81.59±3.80 (9.06)
6	71.01 ^{def} (8.46)	74.11 ^{cdef} (8.64)	80.03 ^{abcde} (8.97)	88.17 ^{abc} (9.42)	88.35 ^{abc} (9.43)	91.27 ^{ab} (9.58)	82.16±3.42 (9.09)
7	74.27 ^{cdef} (8.65)	76.35 ^{bcde} (8.77)	81.17 ^{abcde} (9.04)	86.03 ^{abcd} (9.30)	89.01 ^{abc} (9.46)	94.11 ^a (9.73)	83.49±3.11 (9.16)
Control	5.13 ^l (2.37)	5.05 ^l (2.36)	4.05 ^l (2.13)	6.01 ^l (2.55)	4.03 ^l (2.13)	7.05 ^l (2.75)	5.22±0.47 (2.39)
Mean	47.45±11.1 (6.92)	50.94±11.85 (7.17)	55.09±12.84 (7.46)	59.23±13.81 (7.73)	59.94±14.26 (7.77)	62.46±14.4 (7.93)	

HAT = Hours After treatment

* Values in parenthesis are square root transformed values

Our results are in accordance with the findings of [17, 18 and 19], who observed that concentration of *Acorus calamus* essential oil had significant effect on mortality of *Sitophilus Oryzae*.

Formatted: Highlight

Comment [H4]: This discussion is incomplete, articles 17 and 18 focus on the toxic effect of *Acorus calamus* on other pests. They needed to clarify this and say how the results of these articles are similar to yours. Article 19 is interested in the study of The biology of rice weevil, *S. oryzae* on different hosts viz., sorghum, wheat, maize, paddy, barley and oat under laboratory conditions.. it is so far from your study.




Fig 1: Insecticidal activity of *Acorus calamus* essential oil against rice weevil

4. Conclusion

The study concludes that various concentrations of *Acorus calamus* essential oil had significant effect on mortality of *Sitophilus Oryzae* and the essential oil of *Acorus calamus* can be useful in controlling the stored product insect by surface treatment in godowns as an alternative to synthetic insecticides, further the efficacy of the botanical can be tested on other stored product insects also.

References

1. Viñuela E, ADAN A, Del Estal P, Marco V, Budia F. Plagas de los productos almacenados. Hojas divulgadoras-Ministerio de Agricultura, Pesca y Alimentación. 1993(1):1-32.
2. Khan AR, Selman BJ. On the mortality of *Tribolium castaneum* adults treated sublethally as larvae with pirimiphos methyl, *Nosema whitei* and pirimiphos methyl—*N. whitei* doses. *Entomophaga*. 1988 Sep;33:377-80.
3. Pinto Jr AR, Furiatti RS, Pereira PV, Lazzari FA. Avaliação de inseticidas no controle de *Sitophilus oryzae* (L.) (Coleoptera: Curculionidae), e *Rhyzopertha dominica* (Fab.) (Coleoptera: Bostrichidae) em arroz armazenado. *Anais da Sociedade Entomológica do Brasil*. 1997;26:285-90.
4. Brooker, D. B., Bakker-Arkema, F. W., & Hall, C. W. (1992). *Drying and storage of grains and oilseeds*. Springer Science & Business Media.
5. Adane K, Moore D, Archer SA. Preliminary studies on the use of *Beauveria bassiana* to control *Sitophilus zeamais* (Coleoptera: Curculionidae) in the laboratory. *Journal of Stored Products Research*. 1996 Apr 1;32(2):105-13.
6. Khan AR, Selman BJ. *Nosema* spp. (Microspora: Microsporida: Nosematidae) of stored-product Coleoptera and their potential as microbial control agents.
7. Metcalf RL. Changing role of insecticides in crop protection. *Annual review of Entomology*. 1980 Jan;25(1):219-56.
8. Lale NE. A laboratory study of the comparative toxicity of products from three spices to the maize weevil. *Postharvest Biology and Technology*. 1992;2(1):61-4.
9. Isman MB. Leads and prospects for the development of new botanical insecticides. *Rev. Pestic. Toxicol.* 1995;3:1-20.
10. Ofuya TI. Use of wood ash, dry chilli pepper fruits and onion scale leaves for reducing *Callosobruchus maculatus* (Fabricius) damage in cow-pea seeds during storage. *The Journal of Agricultural Science*. 1986 Oct;107(2):467-8.
11. Ajayi O, Arokoyo JT, Nezan JT, Olayinka OO, Ndirmbula BM, Kannike OA. Laboratory assessment of the efficacy of some local plant materials for the control of storage insect pests. *Samaru Journal of Agricultural Research*. 1987;5(1-2):81-6.
12. Schoonhoven AV. Use of vegetable oils to protect stored beans from bruchid attack. *Journal of economic Entomology*. 1978 Apr 1;71(2):254-6.

13. Chiasson H, Vincent C, Bostanian NJ. Insecticidal properties of a Chenopodium-based botanical. *Journal of economic entomology*. 2004 Aug 1;97(4):1378-83.
14. Gupta L, Srivastava M. Effect of Withaniasomnifera extracts on the mortality of *Callosobruchus chinensis* L. *Journal of Biopesticides*. 2008;1(2):190-2.
15. Yankanchi SR, Patil SR. Field efficacy of plant extracts on larval populations of *Plutellaxylostella* L. and *Helicoverpaarmigera* Hub. and their impact on cabbage infestation. *Journal of Biopesticides*. 2009;2(1):32-6.
16. Ebadollahi A, Safaralizadeh M, Pourmirza A. Fumigant toxicity of *Lavandula stoechas* L. oil against three insect pests attacking stored products. *Journal of Plant Protection Research*. 2010;50(1).
17. Mansoor-ul-Hasan MS, Ullah E, Ahmad F, Wakil W. Insecticidal activity of different doses of *Acorus calamus* oil against *Trogoderma granarium* (Everts). *Pak. J. Agri. sci. Val*. 2006;43:1-2.
18. Abhijith, N., Mohan, S. and Nelson, S.J., 2018. Bioassay to detect the toxicity of sweet flag (*Acorus calamus*) 6 EC formulation against pests of stored spices viz., rusty grain beetle, *C. ferrugineus* (Stephens) cigarette beetle, *L. serricornis* (F.) and red flour beetle, *T. castaneum* (Herbst). *Journal of Entomology and Zoology Studies*, 6(5), pp.1921-1924.
19. Manjhu A, Lekha MK, Chhangani G, Kumar K. Host preference studies of rice weevil, *Sitophilus oryzae* L. on various cereals. 2022; 11(1): 1363-1367