

# Original Research Article

## Genetic Diversity and Association Analysis for Agronomic and Yieldrelated Traits in Interspecific Progenies of Pigeonpea (*Cajanus cajan*)

### ABSTRACT

The limited genetic diversity in existing cultivars emphasizes the need for incorporating potential exotics and wild relatives in breeding programs. This study focussed on assessing genetic variation and correlations among yield traits in  $F_8$  and  $F_9$  progenies resulting from nine interspecific crosses, involving three cultivated parental lines and three wild species. Results indicate significant genotypic variation for all traits studied. Phenotypic and genotypic coefficients of variation varied across agronomic parameters. Heritability estimates ranged from 22.61% (Primary branches per plant) to 89.35% (Days to maturity) for different traits, emphasizing the potential for selection in breeding programs. Positive correlations were observed among various traits in both progeny generations, providing insights into the interplay of genetic factors influencing pigeonpea traits. The cluster analysis revealed that, 54 interspecific cross derived lines with three parents were grouped into four clusters, indicating the greater diversity among the lines for the selection of yield related traits for pigeonpea crop improvement.

**Keywords:** Cluster analysis, Correlation, Genetic diversity, Pigeonpea, Variability

### 1. INTRODUCTION

Pigeonpea (*Cajanus cajan* (L.) Millspaugh) holds significant economic importance as a grain legume crop, particularly in the developing countries situated in tropical and subtropical regions worldwide [1]. This crop is recognized for its nutritional value, containing approximately 20-22% protein, dietary fiber, as well as essential vitamins and micronutrients such as zinc (Zn) and iron (Fe). Consequently, pigeonpea plays a crucial role in ensuring food security and contributing to a balanced diet. Its multifaceted applications, including utilization as food, fodder, fuel, for soil conservation, integrated farming systems, and symbiotic nitrogen fixation, underscore its vital role in subsistence agriculture[2]

Globally, pigeonpea cultivation spans an area of 6.09 million hectares, yielding 5.01 million tonnes with a productivity rate of 822 kg/ha [3]. Notably, India stands as the primary contributor, accounting for 90% of the global pigeonpea production. In India, pigeonpea is cultivated over 5.01 million hectares, resulting in a production of 3.89 million tonnes and a productivity of 859 kg/ha. Despite these substantial production figures, the national harvest falls short of meeting the demands

of India's growing population, necessitating the import of approximately 1.2 million tonnes annually from African countries such as Mozambique, Tanzania, Uganda, and Kenya. This import practice incurs a cost of approximately USD 116.57 million [4].

The majority of currently available pigeonpea cultivars exhibit a limited genetic diversity, necessitating prompt remedial actions through the inclusion of potential exotics and wild relatives in breeding programs. Interspecific hybridization has proven effective in enhancing crop plants by facilitating the transfer of desirable genes and chromosomes from wild to cultivated species. Notably, *Cajanus scarabaeoides* has been successfully employed as a donor parent at the ICRISAT Centre to transfer high-protein genes to pigeonpea[5].

*Cajanus cajanifolius*, identified as the putative progenitor of pigeonpea, imparts resistance to bruchid and pod borer, as well as tolerance to drought, seed hardness, and elevated levels of both protein and methionine. Considering these factors, the present study aims to augment our understanding by investigating the  $F_8$ , and  $F_9$  progenies resulting from nine inter-specific crosses, to assess the extent of genetic variation and correlations among yield traits in the  $F_8$  and  $F_9$  generation progenies derived from these inter-specific crosses.

## 2. MATERIAL AND METHODS

The experimental investigation was carried out at the Norman E Borlaug Crop Research Centre, Govind Ballabh Pant University of Agriculture and Technology, Pantnagar, Uttarakhand (29°N latitude; 79.30°E longitude; 243.84 meters above mean sea level). This study encompassed the examination of progenies resulting from nine inter-specific crosses originating from cultivated parental lines, namely, ICPL 84023, Pant A134, and UPAS120, and wild species, including *Cajanus scarabaeoides*, *Cajanus acutifolius*, and *Cajanus cajanifolius*. The crosses were executed earlier and advanced through pedigree method of selection. The lines were available at  $F_8$  and  $F_9$  were used for the study in the year 2009-10 and 2010-11 respectively. (Table 1). In the kharif seasons of 2009-10 and 2010-11, seeds of  $F_8$  and  $F_9$  progenies were sowed, along with three cultivated parent lines, in a randomized complete block design, wherein 54 inter-specific progeny lines and the three cultivated parent lines were sowed across three replications. Field data were recorded (90 cm x 30 cm, 3 m plot size) with on five randomly selected plants within each plot.

The observations on the number of days to 50% flowering, number of days to maturity, Plant height, number of primary branches per plant, number of pods per plant and Seed yield per plant were recorded at appropriate growth stages. In the analysis of variance within the experimental traits, the coefficients of variability, heritability, and simple correlation were determined through the utilization of mean data encompassing all parameters. Statistical analysis was conducted to assess the significance of disparities among the means of diverse characters. The procedures employed for the analysis of variance adhered to the framework outlined by [6]. Phenotypic, genotypic, and environmental coefficients of variation were computed by expressing the ratio of the respective standard deviation to the overall mean of the characters in percentage terms, following the

methodology proposed by [7]. The calculation of heritability in the broad sense involved determining the ratio of genotypic variance to phenotypic variance, expressed as a percentage [8]. The anticipated genetic advance under selection was computed. Correlation coefficients at both the phenotypic and genotypic levels, relating to all conceivable pairs of characters, were estimated using the methodology presented by Searle [9].

### 3. RESULTS AND DISCUSSIONS

**Analysis of Variance:** Analysis of variance was conducted for progenies of nine interspecific crosses and their three cultivated parents, revealing that the mean square attributable to genotypes was profoundly significant for all examined traits at a 5% level of significance. This observation signifies the existence of substantial variability within the studied material. The mean squares associated with genotypes, replications, as well as the general mean, standard error of the mean, critical difference at a 5% level (CD at 5%), and coefficient of variation (CV) are comprehensively presented in [Table 2](#) and [Table 3](#).

**Phenotypic and Genotypic coefficient of variation:** During the kharif seasons of 2009-10 and 2010-11, various agronomic traits were assessed to determine the phenotypic and genotypic variations within the studied crop. In the kharif season 2009, the coefficient of phenotypic variation (PCV %) reached its highest value for pods per plant (34.25%), succeeded by seed yield per plant (31.64%), and primary branches per plant (22.97%), while the minimum PCV % was observed for days to maturity (8.05%). Similar trends were observed in the subsequent kharif 2010, with the maximum PCV % recorded for pods per plant (35.92%), followed by seed yield per plant (33.74%) and primary branches per plant (25.14%), and the minimum PCV % registered for days to maturity (7.35%) (Supplementary [Table S1](#) and Supplementary [Table S2](#)).

In parallel, the assessment of genetic coefficient of variation (GCV %) during the kharif season 2009 indicated the highest values for pods per plant (29.55%), seed yield per plant (24.81%), and primary branches per plant (17.06%), while the lowest GCV % was observed for days to maturity (6.26%). Corresponding observations were made in the kharif 2010-11, with maximum GCV % values noted for pods per plant (27.58%), seed yield per plant (20.83%), and primary branches per plant (11.95%), and the minimum GCV % observed for days to maturity (6.94%) (Supplementary [Table S1](#) and Supplementary [Table S2](#)).

Additionally, the evaluation of genotypic coefficient of variation (GCV %) in the kharif season 2009 revealed the highest values for seed yield per plant (24.81%), pods per plant (29.55%), and primary branches per plant (17.06%), with the lowest GCV % observed for days to maturity (6.26%). Analogous results were obtained in the kharif season 2010, where the maximum GCV % was recorded for seed yield per plant (20.83%), followed by pods per plant (27.58%), while the minimum GCV % was noted for days to maturity (6.94%) (Supplementary [Table S1](#) and Supplementary [Table S2](#)).

Estimation of heritability and Genetic advance for various agronomic and yield traits: Results from the kharif seasons of 2009-10 and 2010-11 indicate varying heritability estimates for different agronomic traits. In the kharif 2009-10 dataset, the trait with the highest heritability was pods per plant (74.44%), followed by seed yield per plant (61.49%), and day to maturity (60.51%). Conversely, plant height exhibited the lowest heritability at 38.76%. In the subsequent kharif season of 2010-11, heritability values ranged from 22.61% for primary branches per plant to 89.35% for days to flowering. Moderate heritability was observed for seed yield per plant (38.13%), pods per plant (58.94%), and plant height (45.20%) during this period (Supplementary Table S1 and Supplementary Table S2). Specifically, during kharif 2009-10, the Genetic advance as % of mean highest value was recorded for pods per plant (52.53%), followed by seed yield per plant (40.08%) and primary branches per plant (26.07%). Conversely, days to maturity exhibited the lowest value at 16.94%. In the subsequent kharif season of 2010-11, the observed results demonstrated the highest genetic advance mean for pods per plant (43.62%), followed by days to 50% flowering (28.03%), seed yield per plant (26.68%), with the lowest value observed for days to maturity (13.54%) (Supplementary Table S1 and Supplementary Table S2).

Genotypic and phenotypic correlation coefficients: In the  $F_8$  progeny generation, a significant and positive correlation was observed between the duration of flowering and variables such as plant height, pods per plant, and the number of primary branches per plant. However, there was no statistically significant correlation observed between plant height and seed yield per plant. Conversely, for the remaining traits, highly significant and positive correlations were identified. In the subsequent  $F_{10}$  progeny generation, all studied characteristics exhibited a highly significant and positive correlation, as detailed in Supplementary Table S3 and Supplementary Table S4.

Cluster analysis for interspecific cross progenies: The cluster analysis revealed that, 54 interspecific progenies and three parents were grouped into four clusters (Fig. 1). The first cluster (I) consists of 26 lines, for the cluster average plant height about 159.58 cm, number of primary branches per plant 11.85, number of pods per plant 122.32, days to flowering 85.24, days to maturity 176.85, grain yield per plant 26.91 g. This cluster was mainly formed from the lines derived from, UPAS 120 with all three wild species and few of PA 134 with all three wild species. In the second cluster (II), 10 lines were grouped together with average plant height of 151.19 cm, number of primary branches per plant was 13.53, number of pods per plant was 157.04, days to flowering was 78.47, days to maturity about 174.50, seed yield per plant was 34.55 g. This cluster mainly contains the genotype from PA 134 with all three wild species except G44 which is derived from UPAS 120 crossed with *C. acutifolius*. The third cluster (III) consists of 18 lines with one parental line (ICPL 84023), having average plant height of 148.76 cm, number of primary branches per plant 8.92, number of pods per plant was 86.66, days to flowering was 70.53, days to maturity about 156.65, seed yield per plant was 18.79 g. The III cluster consists of lines only from ICPL 84034 with all three wild species along with the parent ICPL 84023. The last cluster (IV) consist of two parental lines, PA 134 and UPAS 120, with average plant height of 179.55 cm, number of primary branches per plant was 10.29, number of pods per plant was

185.91, days to flowering was 74, days to maturity about 174.83, seed yield per plant was 20.12 g (Table 4).

Lack of genetic variability in plant varieties results in naturally low yielding potential; inefficient plant types also have low yielding potential; there are no suitable ideotypes for various cropping systems; poor harvest index; low crop management; increased weed competition; and susceptibility to biotic and abiotic stresses are the main obstacles to achieving higher yields. As like in our study, high variability for different yield characters was also reported in pigeonpea from other studies (10-14). Heritability estimates of yield traits shows moderate to high heritability for pods per plant, seed yield and days to maturity in both  $F_8$  and including day to maturity, days to 50% flowering in  $F_9$ . High heritability for different characters were reported in pigeonpea by different studies [15-18]. Yield contributing characters had moderate heritability with moderate to high genetic advance as per cent mean were exhibited by pods per plant, seed yield per plant in  $F_8$  progenies and in  $F_9$  it was found only in seed yield per plant. This indicates preponderance of additive gene action for these traits. In rest of the characters, in both the generations the estimates of low heritability with low genetic advance as % mean indicates the preponderance of non-additive gene action and hence selection may not be effective for these characters. The disagreement of the previous reports with the present investigation on heritability for some characters could be ascribed to the population and the environment in which the estimates of heritability for some characters could be ascribed to the population and the environment in which the estimates of heritability were taken as the heritability estimates are subjected to these conditions. In the  $F_8$  progeny generation, a robust and positive correlation was evident for all examined traits, except in the case of plant height and seed yield per plant. Notably, a highly significant and positive correlation was identified for all traits in  $F_9$  progeny generation. Similar findings were also reported by different studies [19-24] in past. The cluster analysis among the pigeonpea hybrids [25], land races [26], germplasm lines [28] were reported earlier, however the clustering patterns of interspecific pre-bred lines carried out in this study is unique. The lines clustered were distinguishable based on the modern cultivars involved in the cross. Further studies using molecular tools will be helpful and through a light on combining ability and gene introgression patterns in the interspecific crosses of pigeonpea.

#### **4. CONCLUSION**

Consequently, the findings suggest the presence of considerable genetic diversity for the grain yield and its related traits, including seed yield per plant and the number of pods per plant, demonstrating substantial genetic advancement. Notably, a noteworthy and positive correlation was observed between seed yield per plant and the absence of primary branches, as well as the number of pods per plant. Enhancements in these traits are anticipated to positively influence seed yield. The results of the cluster analysis showed that four clusters comprising 54 interspecific cross-derived lines with three parents were formed, suggesting a higher degree of diversity among the lines for the selection of yield-related traits for the improvement of the pigeonpea crop.

#### **REFERENCES**

1. Varshney RK, Chen W, Li Y, Bharti AK, Saxena RK, Schlueter JA, Donoghue MT, Azam S, Fan G, Whaley AM, Farmer AD. Draft genome sequence of pigeonpea (*Cajanus cajan*), an orphan legume crop of resource-poor farmers. *Nat Biotechnol.* 2012;**30**(1): 83.
2. Shruthi HB, Hingane AJ, Reddi MS, Kumar SCV, Prashanthi L, Reddy BBV, Rathore A. Genetic Divergence for Yield, Physiological and Quality Traits in Super-Early Pigeon pea (*Cajanus cajan* (L.) Millsp.). *Int J Curr Microbiol Appl Sci.* 2020;9(1):2422-2433.
3. Food and Agricultural Organization.FAOSTAT database. <http://www.fao.org>. 2020.
4. Connect2India. Pigeonpea import to India. <http://connect2india.com/global/Pigeon-Pea-import-to-india/1>. 2021.
5. Saxena KB, Singh L, Reddy MV, Singh U, Lateef SS, Sharma SB, Ramanandan P. Intra-species variation in *Atylosiascarabaeoides*(L.) Benth and wild relatives of Pigeon Pea [*Cajanus cajan* (L.) Mill sp.]. *Euphytica.*1990;19:185-191.
6. Panse VG, Sukhatme PV. *Statistical Methods for Agricultural Workers.* ICAR Publication, New Delhi.1985.
7. Burton GW, Devane DE. Estimating heritability in tall fescue (*Festuca arundinacea*) from replicated clonal material. *Agronomy J.*1953;45(10): 478-481.
8. Allard RW.Principles of Plant Breeding. New York: John Wiley and Sons;1960.
9. Searle SR. Phenotypic, genotypic and environmental correlations. *Biometrics.* 1961; 17:474–80.
10. Balyan HS, Sudhakar MV. Variability, character association and path coefficient studies on genotypes of early maturing group in pigeonpea [*Cajanus cajan*(L.) Millsp.]. *Madras Agri J.* 1985;72(3):168-172.
11. Pundir RPS, Singh RB. Possibility of genetic improvement of pigeonpea[*Cajanus cajan*(L.) Millsp.] utilizing wild gene sources. *Euphytica.*1987;36:33-37.
12. Dahat DV, Deshmukh RB, Patil JV.Genetic variability and character association in pigeonpea. *Int J Agr Res.*1997;31(2):82-86.
13. Patel JA, Patel DB, Zaveri PP, Pathak AR. 1988. Path coefficient studies in pigeonpea. *Int Pigeonpea New.* 1988;7:5-6.
14. Venkateswarlu O. Genetic variability in pigeonpea [*Cajanus cajan* (L.) Millsp.]. *Legume res.* 2001;24(3): 205-206.
15. Jagshoran. Studies in genetic variability for some quantitative character in pigeonpea [*Cajanus cajan* (L.) Millsp.].*Madras Agric J.*1983. 70;146-148.
16. Saxena AK, Kataria VP. Study of genetic variability for yield and its component characters in pigeonpea. *Bhartiya Krishi Anusandhan Patrika.*1993;8(3-4): 189-192.
17. Aher RP, Thombre BB, Dahat DV. Genetic variability and character association in pigeonpea. *Legume Res.* 1998;21(1):41-44.
18. Baskaran K, Muthiah AR. Variability studies in pigeonpea [*Cajanus cajan* (L.) Millsp.]. *Res crop.* 2007;7(1):249-252.
19. Sinha SC, Srivastava AN, Singh KN. Interrelationship of some quantitative traits in pigeonpea. *Int PigeonpeaNewsl.*1987;6:21-24.

20. Balkrishna K, Natarajaratnam N. Association of yield attributes in pigeonpea. Madras Agri J.1989;76(6):349-350.
21. Ganeshamurti K, Durairaj MS.Character association on pigeonpea [*Cajanus cajan* (L) Mill sp.].Madras Agric J.1990;77(5-6):201-204.
22. Henry A, Krishna GVSR. 1990. Correlation and path coefficient analysis in pigeonpea. Madras Agric J. 1990;77:440-442.
23. Salunke DK, Chavan JK, Kadam SS.Pigeonpea as important food source. Crit Rev Food Sci Nutr. 1986;23(2): 103-141.
24. Deshmukh RB, Rodge RG, Patil JV, Sahane DV.Genetic variability and character association in pigeonpea under different cropping systems. J Maharashtra Agri Univ. 2000.25(2):176-178.
25. Sandeep S, Sujatha K, Minnie CM, RaniCS.Genetic analysis of pigeonpea [*Cajanus cajan* (L.) millsp.] hybrids for yield and yield attributes. Legume Res. 2022;45(10):1216-1222.
26. Reddy SD, Jayamani P. 2019. Genetic diversity in land races of pigeonpea (*Cajanus cajan*(L.) Millsp.). Electron J Plant Breed. 2019 10(2):667-672.
27. Sujit MK, Belsariya N, Pandey VK. 2023 Studies on genetic diversity for yield and yield attributing traits in pigeonpea [*Cajanus Cajan* (L.) Millsp] germplasm. J Pharm Innov. 2023;12(3):5272-5277.

**Table 1.List of interspecific crosses and their cultivated parents**

S. No.	Cultivated Parents	
1	ICPL84023	Determinate, early maturing line (G55)
2	PA134	Indeterminate, early maturing line (G56)
3	UPAS120	Indeterminate, early variety (G57)
S. No.	Interspecific crosses	
1	ICPL 84023 x <i>C. scarabaeoides</i> (G1-G6)	
2	ICPL 84023 x <i>C. acutifolius</i> (G7-G12)	
3	ICPL 84023 x <i>C. cajanifolius</i> (G13-G18)	
4	PA134 x <i>C. scarabaeoides</i> (G19-G24)	
5	PA134 x <i>C. acutifolius</i> (G25-G30)	
6	PA134 x <i>C. cajanifolius</i> (G31-G36)	
7	UPAS 120 x <i>C. scarabaeoides</i> (G37-G42)	
8	UPAS 120 x <i>C. acutifolius</i> (G43-G48)	
9	UPAS 120 x <i>C. cajanifolius</i> (G49-G54)	

**Table 2. Analysis of variance for agronomic and yield related traits in F<sub>8</sub> progenies of interspecific crosses and cultivated parents.**

Source of variation	d.f.	Days to 50% flowering	Days to maturity	Plant height (cm)	Primary branches/plant	Pods per plant	Seed yield per plant
Replication	2	100.32	66.27	1590.55	24.45	2373.95	107.19
Treatment	56	307.96**	408.38**	732.45**	12.90**	4670.46**	126.54**
Error	112	73.32	72.95	252.81	2.75	479.53	21.85
GM±		76.72	168.79	158.74	10.78	126.46	23.80
SEM		4.94	4.93	9.17	0.95	12.64	2.69
CD at 5%		13.82	13.78	25.66	2.67	35.34	7.54
CV (%)		11.16	5.06	10.01	15.38	17.31	19.63

\* Significant at 5% level of probability \*\* Significant at 1% level of probability

**Table 3. Analysis of variance for agronomic and yield related traits in F<sub>9</sub> progenies of inter-specific crosses and cultivated parents.**


Source of variation	d.f.	Days to 50% flowering	Days to maturity	Plant height (cm)	Primary branches/plant	Pods per plant	Seed yield per plant
Replication	2	3.03	650.91	3948.5	52.19	8326.38	382.60
Treatment	56	409.27**	507.87**	1596.74**	13.06**	5635.59**	130.51**
Error	112	63.99	63.98	509.07	6.95	1061.92	47.18
GM±		73.95	170.35	194.34	11.92	141.56	25.34
SEM		4.61	4.62	13.02	1.52	18.81	3.96
CD at 5%		12.91	12.92	36.41	4.25	52.59	11.08
CV (%)		10.81	4.69	11.60	22.12	23.01	27.10

\* Significant at 5% level of probability \*\* Significant at 1% level of probability

**Table 4. Details of five clusters with traits, Plant height (PH), Number of primary branches per plant (NP), Number of Pods per plant (NP), Days to 50% flowering (DF), Days to maturity (DM) and Seed yield per plant (SY).**

Cluster	PH	NB	NP	DF	DM	SY	Genotypes
---------	----	----	----	----	----	----	-----------

I	159.58	11.85	122.32	85.24	176.85	26.91	G54, G46, G40, G47, G35, G43, G34, G51, G53, G52, G50, G19, G26, G39, G48, G37, G33, G25, G42, G27, G45, G38, G49, G31, G32, G41
II	151.19	13.53	157.04	78.47	174.50	34.55	G20, G21, G22, G23, G24, G28, G29, G30, G36, G44
III	148.76	8.92	86.66	70.53	156.65	18.79	G1, G2, G3, G4, G5, G6, G7, G8, G9, G10, G11, G12, G13, G14, G15, G16, G17, G18, G55
IV	179.55	10.29	185.91	74	174.83	20.12	G56, G57



**Fig. 1. Cluster analysis for 54 lines derived from 9 interspecific cross progenies of pigeonpea along with the parents**

**Table S1. Range, general mean, heritability, genetic advance PCV and GCV in F<sub>8</sub> progenies and cultivated parents for yield and yield related traits.**

Character	Range	General mean	PCV%	GCV%	ECV%	Heritability (%)	Genetic advance	Genetic advance as % of mean
Days to 50% flowering	51.66-96.33	76.72±4.94	16.04	11.52	11.16	51.61	13.08	17.05

Days to maturity	142.0-190.3	168.79±4.93	8.05	6.26	5.06	60.51	16.94	10.04
Plant height (cm)	137.0-199.5	158.74±9.17	12.79	7.96	10.01	38.76	16.21	10.21
Primary branches per plant	6.93-14.38	10.78±0.95	22.97	17.06	15.38	55.16	2.81	26.07
Pods per plant	72.82-215.82	126.46±12.64	34.25	29.55	17.31	74.44	66.43	52.53
Seed yield per plant (g)	11.55-37.47	23.80±2.69	31.64	24.81	19.63	61.49	9.54	40.08

**Table S2. Range, general mean, heritability, genetic advance PCV and GCV in F<sub>9</sub> progenies and cultivated for yield and yield related traits.**

Character	Range	General mean	PCV%	GCV%	ECV%	Heritability (%)	Genetic advance	Genetic advance as % of mean
Days to 50% flowering	50.0-96.0	73.95±4.61	13.49	12.72	44.76	88.99	20.73	28.03
Days to maturity	145.0-187.66	170.35±4.62	7.35	6.94	2.40	89.35	23.06	13.54
Plant height (cm)	130.7-223.1	194.34±13.02	14.55	9.79	10.77	45.20	26.65	13.71
Primary branches per plant	7.39-15.46	11.92±1.52	25.14	11.95	22.12	22.61	1.39	11.66
Pods per plant	80.05-212.31	141.56±18.81	35.92	27.58	23.01	58.94	61.75	43.62
Seed yield	12.29-	25.34±3.96	33.74	20.83	26.53	38.13	6.76	26.68

per plant 40.003  
(g)

**Table S3. Phenotypic and genotypic correlation coefficients for yield and yield contributing traits in F<sub>8</sub>.**

	Days to 50% flowering	Day to maturity	Plant height (cm)	Primary branches per plant	Pods per plant	Seed yield per plant
Days to 50% flowering	$r_p$	0.604**	0.211*	0.450**	0.270*	0.473**
	$r_g$	0.990**	0.470**	0.887**	0.398**	0.810**
Day to maturity	$r_p$		0.325**	0.533**	0.512**	0.525**
	$r_g$		0.656**	0.909**	0.720**	0.806**
Plant height (cm)	$r_p$			0.255*	0.534**	0.066
	$r_g$			0.253*	0.964**	0.102
Primary branches per plant	$r_p$				0.509**	0.775**
	$r_g$				0.508**	0.929**
Pods per plant	$r_p$					0.577**
	$r_g$					0.393**

\* Significant at 5% level of probability \*\* Significant at 1% level of probability

**Table S4. Phenotypic and genotypic correlation coefficients for yield and yield contributing traits in F<sub>9</sub>.**

	Days to 50% flowering	Day to maturity	Plant height (cm)	Primary branches per plant	Pods per plant	Seed yield per plant
Days to 50% flowering	$r_p$	0.757**	0.549**	0.566**	0.279**	0.625**
	$r_g$	0.797**	0.790**	0.999**	0.285**	0.919**
Day to maturity	$r_p$		0.558**	0.422**	0.466**	0.454**
	$r_g$		0.858**	0.831**	0.567**	0.674**
Plant height (cm)	$r_p$			0.551**	0.443**	0.448**
	$r_g$			0.999**	0.640**	0.751**
Primary branches per plant	$r_p$				0.519**	0.707**
	$r_g$				0.363**	0.808**
Pods per	$r_p$					0.649**

---

plant

$r_g$

0.317\*\*

---

\* Significant at 5% level of probability    \*\* Significant at 1% level of probability

UNDER PEER REVIEW