

# **Effect Of Panchagavya On Growth, Yield And Quality Of Zucchini (*Cucurbita pepo*) Under Protected Condition**

## **ABSTRACT**

An experiment was conducted to study the effect of Panchagavya on Growth, Yield and Quality of Zucchini (*Cucurbita pepo*) in Protected Condition at Sam Higginbottom. University Of Agriculture Technology And Sciences, Prayagraj during summer 2023. The Treatment Performed best in terms of Plant Height (19.83 cm) and Number of node (12.00) after Foliar spray of Panchagavya (6%) at 30 DAS. Days to emergence of 1<sup>st</sup> male flower/ plant (28.87) after drenching of Panchagavya (9%) at 30 DAS. Days to emergence of 1st female flower/plant (34.03) after drenching of Panchagavya (9%) at 15 DAS. Days to 50% female flower/plant (34.67) Foliar spray of Panchagavya (6%) at 30 DAS. Days to 1st fruit setting/plant (43.00) after foliar spray of Panchagavya (9%) at 15 DAS. The Number of fruits/plant (4.67) after Foliar spray of Panchagavya (6%) at 15 DAS and the average weight of fruit (197.50g) after Foliar spray of Panchagavya (3%) at 15 DAS. Fruit length and Girth (25.00 Cm) and (16.00mm) after Drenching of Panchagavya (9%) at 30 DAS . Fruit Yield/plant (1.113 kg) after Foliar spray of Panchagavya (3%) at 15 DAS and Fruit yield per 250 m<sup>2</sup> was (3.71q). Fruit quality TSS°brix (3.22) after Foliar spray of Panchagavya (9%) at 30 DAS. The highest benefit cost ratio was found after Foliar spray of Panchagavya (3%) at 15 DAS (2.42). Result indicates that after Foliar spray of Panchagavya (3%) at 15 DAS were recommended for plant.

**Key word- Panchagavya, Zucchini, Organic Farming, Protected Condition.**

## **INTRODUCTION**

Zucchini (*Cucurbita pepo* L.) is a fruit belonging to the *Cucurbita* genus and the *Cucurbitaceae* family. This family is made up of 22 species; the 5 most cultivated being *C. maxima*, *C. moschata*, *C. pepo*, *C. ficifolia*, and *C. argyrosperma*. According to the Ministry of Agriculture, Fisheries, and Food, the production of zucchini in Spain is 591,341 tons (t) and a cultivation area of 11,037 hectares (ha).<sup>1</sup> Zucchini is one of the lowest caloric content vegetables (14 kcal/100 g) and has the highest water content (96.5%). Its consumption covers the needs of vitamins and minerals, especially

vitamin C and potassium, vitamin C being the most significant. The presence of mucilage gives it emollient properties on the digestive system and as an easy-to-digest food, it is suitable for those with digestive problems.

Heavy use of chemicals in agriculture has weakened the ecological base in addition to degradation of soil, water resources and quality of the food. At this juncture, a keen awareness has sprung on the adoption of "organic farming" as a remedy to cure the ills of modern chemical agriculture (Kannaiyan, 2000).

Panchagavya is a term used in Ayurveda fermented product made from five ingredients obtained from cow, such as milk, urine, dung, curd and clarified butter (Amalraj et al., 2013). Role of foliar applied panchagavya in production of many plantation crops has been well documented in India (Selvaraj, 2003). Panchagavya is a popular foliar nutrition prepared by organic growers of Tamil Nadu as an indigenous material and used widely for agricultural

horticultural and crops (Swaminathan et al., 2007). The soil fertility is enhanced by an increase in organic substances and micro-nutrients, which are important for promoting the growth of crops and the reproduction of microorganism that improve the texture of the soil. Physico-chemical properties of Panchagavya revealed that they possess almost all the major nutrients, micro nutrients and growth hormones (IAA & GA) required for crop growth. Predominance of fermentative microorganisms like yeast and lactobacillus might be due to the combined effect of low pH, milk products and addition of jaggery/sugarcane juice as substrate for their growth. It acts as an organic growth-promoter and immunity booster for plants, Yield enhancement by 18% and in few cases like Cucumber.

The present investigation was conducted to examine the effect of foliar and drenching application of panchagavya on different parameters, growth yield and quality of Zucchini.

## Materials and Methods

This experiment was carried out during in 2023 at Horticulture Research Farm, SHUATS, Prayagraj, Uttar Pradesh, India which is located at 25.28 °N latitude, 81.54 °E longitude and 98 m altitude above the mean sea level. This area is situated on the right side of the river Yamuna by the side of Prayagraj Rewa Road about 5 km away from Prayagraj, city. The experiment was done in the polyhouse. The highest O<sub>2</sub> and CO<sub>2</sub> concentration was 20.2% and 480 ppm respectively. In the polyhouse the maximum temperature was 34.7 °C. The maximum light intensity inside and outside the polyhouse was 63600 lux and 110600 lux.

The temperature reached up to 48°C in summer and in winter it goes down to as low as 2-3 °C. The experiment was laid out in a randomized block design with 12 treatments and three replications. The crop was grown in a naturally ventilated polyhouse.

The seeds of Zucchini (F1 Hybrid) were collected from Plantsman F1 hybrid variety Rajbaha road Patiala (Punjab) India. The sowing of seed performed on 1<sup>st</sup> May 2023 at 45 cm × 60 cm spacing on the raised bed. The plants were grown over the bed. The necessary recommended cultural practices like Panchagavya application, irrigation and weeding.

## Statistical Analysis

The data recorded during the course of the investigation were subjected to statistical analysis as per the method of analysis of variance [2]. This investigation was done in the RBD statistical method. The significance

and non-significance of the treatment effect were judged with the help of 'f' value (variance ratio) and compared with the table value at a 5% level of significance. If the calculated value exceeds then the value, the effect of considered to be significant. The significant difference between the means was tested against the critical difference at a 5% level of significance.

### **Treatments**

The present investigation had following treatments T0-control, T1-Drenching of Panchagavya(3%)at15DAS, T2 - Drenching of Panchagavya(3%)at30DAS, T3 - Drenching of Panchagavya(6%) at 15 DAS, T4- Drenching of Panchagavya(6%) at 30 DAS, T5- Drenching of Panchagavya(9%)at15DAS, T6- Drenching of Panchagavya(9%)at30DAS, T7 - Foliar spray of Panchagavya (3%) at 15 DAS, T8 - Foliar spray of Panchagavya (3%) at 30 DAS, T9 - Foliar spray of Panchagavya (6%) at 15 DAS, T10 - Foliar spray of Panchagavya (6%) at 30 DAS, T11 - Foliar spray of Panchagavya (9%) at 15 DAS, T12 - Foliar spray of Panchagavya (9%) at 30 DAS.

### **RESULTS AND DISCUSSION**

Observations were recorded for growth parameters which are Plant height (cm),

Number of node, Number of male flower, Number of female flower, Flowering of 50% female flower, first fruit setting, Number of fruits/plant and the average weight of fruit(gm), Fruit length(cm) and Girth(mm), Fruit Yield(250sq.m). Fruit quality(TSS°brix). In economics parameters are cost of cultivation (Rs/250sq. m), Gross Return (Rs/250sq. m), Net Return (Rs/250sq. m) and Benefit Cost Ratio (Rs/250sq. m).

### **Growth parameters**

#### **Plant height (cm)**

The Maximum plant height at 30 DAS was recorded in the Treatment 10 (19.83 cm) given in (Table 1). This might be due to application of Panchagavya in different concentration and due to the favourable climatic conditions and sufficient accumulation of photosynthesis in the polyhouse condition

#### **Number of nodes/plant**

The Maximum number of node at 30 DAS was recorded in the Treatment 10 (12.00cm) given in (Table 1). This might be due to application of Panchagavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition

**Table .1 Height of Plant (cm) and number of node/plant as affected by application of Panchagavya in different concentration in protected condition.**

Symbol	Treatments	Plant height/plant (cm) 30 DAS	Node/plant 30 DAS
T0	Control	10.42	9.47
T1	Drenching of Panchagavya(3%)at15DAS	13.17	10.67
T2	Drenching of panchagavya(3%)at30DAS	12.67	9.83
T3	Drenching of Panchagavya(6%) at 15 DAS	11.50	10.00
T4	Drenching of Panchagavya(6%)at 30 DAS	14.00	9.30
T5	Drenching of Panchagavya(9%)at15DAS	13.80	8.17
T6	Drenching of Panchagavya(9%)at30DAS	12.17	10.00
T7	Foliar spray of Panchagavya (3%) at 15 DAS	13.50	9.00
T8	Foliar spray of Panchagavya (3%) at 30 DAS	11.67	9.50
T9	Foliar spray of Panchagavya (6%) at 15 DAS	9.50	8.00
T10	Foliar spray of Panchagavya (6%) at 30 DAS	19.83	12.00
T11	Foliar spray of Panchagavya (9%) at 15 DAS	15.97	10.33
T12	Foliar spray of Panchagavya (9%) at 30 DAS	14.17	10.00
	F TEST	S	S
	SE(m) ±	0.33	0.25
	C.D. at 5%	0.97	0.72
	C.V. (%)	4.11	4.36

### Earliness parameter

#### Days to emergence of 1<sup>st</sup> male flower/plant

The first male flower initiation were recorded in the Treatment 6 (28.87) given in (Table 2). This might be due

to application of Panchagavya in different concentration and male flower come early reason high temperature, Better environmental conditions and available nutrients seems to have brought quick changes in plant growth and development.

#### Days to emergence of 1<sup>st</sup> female flower/plant

The first Female flower initiation were recorded in the Treatment 5 (34.03) given in (Table 2). This might

be due to application of Panchagavya in different concentration, Better environmental conditions and available nutrients seems to have brought quick changes in plant growth and development.

#### **Days to 50% flowering/plant**

Days of 50% Flowering female flower initiation were recorded in the Treatment 10 (34.67) given in (Table 2). This might be due to application of Panchagavya in different concentration, Better environmental conditions and available nutrients

seems to have brought quick changes in plant growth and development.

#### **Days to 1st fruit setting/plant**

Days to first fruit setting were recorded in the treatment 11 (43.00) given in (Table 2). This might be due to application of Panchgavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition.

**Table .2 Days to emergence of 1<sup>st</sup> male flower/plant, Days to emergence of 1<sup>st</sup> female flower/plant, Days to 50% flowering/plant and Days to first fruit setting/plant as affected by application of Panchagavya in different concentration in protected condition.**

Symbol	Treatments	Days to emergence of 1st male flower /plant	Days to emergence of 1st female flower/plant	Days to 50% female flower/plant	Days to 1st fruit setting/per plant
T0	Control	33.67	35.83	34.67	42.50
T1	Drenching of Panchagavya(3%)at15DAS	34.67	36.83	36.00	43.83
T2	Drenching of panchagavya(3%)at30DAS	33.33	36.00	35.00	43.17

T3	Drenching of Panchagavya(6%) at 15 DAS	31.17	38.50	37.67	45.17
T4	Drenching of Panchagavya(6%)at 30 DAS	30.93	39.33	37.67	45.50
T5	Drenching of Panchagavya(9%)at15DAS	31.17	34.03	37.67	45.17
T6	Drenching of Panchagavya(9%)at30DAS	28.87	37.33	36.00	44.33
T7	Foliar spray of Panchagavya (3%) at 15 DAS	33.83	37.50	37.00	44.67
T8	Foliar spray of Panchagavya (3%) at 30 DAS	34.67	37.50	39.33	44.17
T9	Foliar spray of Panchagavya (6%) at 15 DAS	34.33	37.00	35.67	43.33
T10	Foliar spray of Panchagavya (6%) at 30 DAS	33.17	36.50	34.67	43.17
T11	Foliar spray of Panchagavya (9%) at 15 DAS	33.67	36.83	36.33	43.00
T12	Foliar spray of Panchagavya (9%) at 30 DAS	34.67	38.83	40.67	43.33
	F TEST	S	S	S	S
	SE(m) ±	0.78	0.91	0.88	0.51
	C.D. at 5%	2.30	2.68	2.57	1.51
	C.V. (%)	4.11	4.27	4.12	3.74

### 3.Yield parameter:

#### No. of fruits per plant

The Maximum number of fruits per plant were recorded in the treatment 9 (4.67) given in (Table 3). This might be due to application of Panchagavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better

growth of plants can be expected under protected condition.

#### Avg. weight/Fruits(g)

The Maximum average weight of 1 fruit(g) were recorded in the treatment 7 (197.50 g) given in (Table 3).This might be due to application of Panchagavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better

growth of plants can be expected under protected condition.

**Fruit length (cm)**

The Maximum Fruit diameter were recorded in the treatment 6 (25.00 cm) given in (Table 3).This might be due to application of Panchagavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition.

**Fruit girth (mm)**

The Maximum Fruit diameter were recorded in the treatment 6 (16.00 mm) given in (Table 3).This might be due to application of Panchagavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition.

**Fruit yield/plant (kg)**

The Maximum Fruit yield/plant were recorded in the treatment 7 (1.113 kg) given in (Table 3).This might be due to application of Panchagavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition.

**Yield 250 sqm (Quintal)**

The Maximum Fruit yield Yield 250 sqm (q) were recorded in the treatment7 (3.71q) given in (Table 3).This might be due to application of Panchagavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition.

**Table.3 Number of fruits per plant, Avg. weight/Fruits(g), Fruit length (cm), Fruit girth (mm),Fruit yield /plant (Kg) and Yield 250 m<sup>2</sup> (Quintal) as affected by application of Panchagavya in different concentration in protected condition.**

Symbol	Treatments	Number of fruit/ plant	Avg weight/ Fruit(g)	Fruit length (cm)	Fruit girth (mm)	Fruit yield /plant (Kg)	Fruit yield per 250 m <sup>2</sup> (q)
T0	Control	3.50	138.00	20.00	12.00	0.687	2.32

T1	Drenching of Panchagavya(3%)at15DAS	4.00	134.50	20.00	11.00	0.713	2.37
T2	Drenching of panchagavya (3%)at30DAS	4.11	153.20	22.00	11.50	0.860	2.86
T3	Drenching of Panchagavya(6%) at 15 DAS	4.00	163.00	21.00	14.00	0.920	2.38
T4	Drenching of Panchagavya(6%)at 30 DAS	4.00	134.50	23.00	13.00	0.667	2.24
T5	Drenching of Panchagavya(9%)at15DAS	3.33	133.50	20.00	15.00	0.617	2.25
T6	DrenchingPanchagavya(9%)at30 DAS	3.83	140.50	25.00	16.00	0.747	2.48
T7	Foliar spray of Panchagavya (3%) at 15 DAS	4.50	197.50	23.00	14.00	1.113	3.71
T8	Foliar spray of Panchagavya (3%) at 30 DAS	4.17	174.50	21.00	10.00	1.043	3.27
T9	Foliar spray of Panchagavya (6%) at 15 DAS	4.67	133.50	20.00	11.00	0.800	3.03
T10	Foliar spray of Panchagavya (6%) at 30 DAS	3.67	147.50	21.00	11.00	0.703	2.45
T11	Foliar spray of Panchagavya(9%) at 15 DAS	3.67	125.40	22.00	12.00	0.623	2.07
T12	Foliar spray of Panchagavya (9%) at 30 DAS	3.33	160.50	21.00	11.00	0.753	2.48
	F TEST	S	S	S	S	S	S
	SE(m) ±	0.17	2.97	0.52	0.26	0.02	0.11
	C.D. at 5%	0.50	8.71	1.54	0.77	0.05	0.33
	C.V. (%)	7.62	3.45	4.22	3.67	3.45	7.50

### Fruit quality

#### TSS°Brix

The Maximum TSS (°Brix) were recorded in the treatment 11 and 12 (3.22°Brix) given in

(Table 4). This might be due to application of Panchagavya in different concentration. The temperature can be controlled and regulated under protected condition, therefore healthy and better growth of plants can be expected under protected condition.

**Table.4 TSS (°Brix)as affected by application of Panchagavya in different concentration in protected condition.**

Symbol	Treatments	Fruit quality TSS°brix
T0	Control	3.20
T1	Drenching of Panchagavya(3%)at15DAS	3.20
T2	Drenching of panchagavya(3%)at30DAS	3.12
T3	Drenching of Panchagavya(6%) at 15 DAS	3.12
T4	Drenching of Panchagavya(6%)at 30 DAS	3.12
T5	Drenching of Panchagavya(9%)at15DAS	3.12
T6	Drenching of Panchagavya(9%)at30DAS	3.20
T7	Foliar spray of Panchagavya (3%) at 15 DAS	3.20
T8	Foliar spray of Panchagavya (3%) at 30 DAS	3.20
T9	Foliar spray of Panchagavya (6%) at 15 DAS	3.11
T10	Foliar spray of Panchagavya (6%) at 30 DAS	3.11
T11	Foliar spray of Panchagavya (9%) at 15 DAS	3.22
T12	Foliar spray of Panchagavya (9%) at 30 DAS	3.22
	F TEST	NS
	SE(m) ±	0.07
	C.D. at 5%	NS
	C.V. (%)	4.06

### Conclusion

From the present investigation it is concluded that the T10 Performed best in terms of growth (Plant Height (19.83 cm) and Number of node/plant (12.00).The T6 is best performed in terms of days to emergence of 1<sup>st</sup> male flower/plant (28.87) and T5 is best performed in terms of days to

emergence of 1<sup>st</sup>female flower/plant (34.03).The T10 is best performed in terms of Days to 50% female flower/plant(34.67) , T11 performed best in term of days to 1<sup>st</sup>fruit setting/plant (43.00) and the T9 best performed in terms of Number of fruits/plant (4.67),the T7Performed best in terms of average weight of fruit (197.50g). The T6 performed best in term of Fruit

length and Girth (25.00 Cm) and (16.00mm). The T7 performed best in term of Fruit Yield/plant (1.113 kg) and Fruit yield per 250m<sup>2</sup> (3.71q).The T11 and 12 performed best in term of Fruit quality TSS°brix (3.22). whereas, the T7 reorded highest benefit cost ratio (2.42).Therefore it was recommended that, foliar spray of Panchagavya 3% at 15 DAS were best for Zucchini plant.

## References

- Pysarenko, V., Pischalenko, M., Chaika, T., Lohvynenko, V., Krupka, N., Koroleviat, Y., & Kirieiev, Y. (2023). The impact of bio-stimulators on zucchini plants under pre-sowing seed treatment.
- Kumar PC, Gobi R, Stalin P, Sathiyamurthi S, Balasubramanian A. Effect of integrated nutrient management on growth and yield of irrigated groundnut (*Arachis hypogaea* L.). *International Journal of Innovative Research in Technology*. 2023;9(9):2349-6002.
- Prakash S, Kumar M, Kumar V, Kumar A. Effect of bio-stimulant on growth and yield of onion (*Allium cepa* L.) cv.NHRDF Red-3. *AMA*. 2022;53(11):10265-1027
- Saini Y, Patel PH, Gangadhara T, Chaudhari P, Bijarnia A. Effect of foliar application of Panchagavya and banana pseudostem sap on the growth, seed yield and economics of fenugreek. *The Indian Society of Agricultural Science*. 2022; 199
- Kumar RR, Shyam A, Adhithya S, Levin Anbu Gomez LA. Effect of Panchagavya on *Sesamum indicum*. *Journal of Research in Applied Science and Engineering Technology*. 2022;46943. DOI:10.22214/ijraset.2022.46943
- Rensang K, Dr. Singh G, Dhaked, Dr. Meghwal ML, Kent B. Effect of organic manures on growth and yield of linseed (*Linum usitatissimum* L.). *International Journal of Advanced Technology in Engineering and Science*. 2022;10(11).
- Rieznik, S., Havva, D., Butenko, A., Novosad, K. 2021. Biological activity of chernozems typical of different farming practices. *Agraarteadus*, 32(2) P. 307-313. DOI: 10.15159/jas.21.34
- Kumar CS, Singh G. Effect of Panchagavya on growth and yield: A review. *International Journal of Current Microbial Applied Science*. 2020; 9:617-624.
- Kumar P, Kaur C, Sethi S, Kaur Jambh H. Effect of extruded finger millet on dough rheology and functional quality of pearl millet- based unleavened flatbread. *Cereal Chemistry*. 2020;97(5): 991-998. DOI:10.1002/cche.10321
- Chongre S, Mondal R, Biswas S, Munshi A. Effect of liquid manure on growth and yield of summer green gram (*Vigna radiata* L.). *Current Journal of Applied Science and Technology*. 2019;38(6):1-7
- Khanna K, Jamwal VL, Kohli SK, Gandhi SG, Ohri P, Bhardwaj R, Ahmad P. Plant growth promoting rhizobacteria induced Cd tolerance in *Lycopersicon esculentum* through altered antioxidative defense expression. *Chemosphere*. 2019;217:463-474.

Joshi, H., Somdutt, Choudhary, P. and Mundra, S.L. (2019). Role of Effective Microorganisms (EM) in Sustainable Agriculture. *International Journal of Current Microbiology and Applied Sciences*. 8(3): 172-181.

Hameedi, A., Thakur, K.S., Kansal, S., Mehta, D.K., Yousafzai, A. and Mohammadi, M.H. (2018). Effect of organic nutrient sources on growth, yield and quality of bell pepper (*Capsicum annuum* L.) under mid hill condition of Himachal Pradesh. *International Journal of Multidisciplinary Research and Development*. 5: 135-138.

Safiullah, K., Durani, A., Durrani, H. and Ansari, M.A. (2018). Effect of solid and liquid organic manures on growth, yield and economics of sweet corn (*Zea mays* L. var. Saccharata Sturt) under south Gujarat condition. *International Journal of Pure Applied Bioscience*. 6(2): 567-574.

Patel DM, Patel IM, Patel BT, Singh NK, Patel CK. Effect of Panchagavya and jivamrut on yield, chemical and biological properties of soil and nutrients uptake by kharif groundnut (*Arachis hypogaea* L.). *International Journal of Chemical Studies*. 2018;6(3):804-809.

Boraiah, B., Devakumar, N., and Palanna, K.B. 2018. Yield and Quality Parameters of Capsicum (*Capsicum annuum* L. var. grossum) as Influenced by Organic Liquid Formulations. *International Journal of Current Microbiology and Applied Sciences*. 6(1): 333-338).

Safiullah, K., Durani, A., Durrani, H. and Ansari, M.A. (2018). Effect of solid and liquid organic manures on growth, yield and economics of sweet corn (*Zea mays* L. var. Saccharata Sturt) under south Gujarat

condition. *International Journal of Pure Applied Bioscience*. 6(2): 567-574.

Kumar D, Massey JX, Sharma SK, Mundra SL, Yadav SK. Vermiwash prepared from different combination of organic sources to improve growth and yield of blackgram [*Vigna mungo*(L.) Hepper] for organic agriculture. *Indian Journal of Agricultural Research*; 2017. DOI: 10.18805/IJARE.A 5630

Panchal P, Patel PH, Patel AG, Desai A. Effect of Panchagavya on growth yield and economics of chickpea (*Cicer arietinum*). *International Journal of Chemical Studies*. 2017;5(2):265-26

Spehia, R.S., Meera Devi, Singh, J., Sharma, S., Negi, A., Singh, S., Chauhan, N., Sharma, D. and Sharma, J.C. (2018). Lettuce growth and yield in Hoagland solution with an organic concoction. *International Journal of Vegetable Science*. 24(6): 557-566.

Boraiah, B., Devakumar, N., Shubha, S. and Palanna, K.B. (2017). Effect of panchagavya, jeevamrut and cow urine on beneficial micro-organisms and yield of capsicum [*Capsicum annuum* (L.) var. grossum]. *International Journal of Current Microbiology and Applied Sciences*. 6: 3226-3234

Chandrakala, M., Hebsur, N.S., Bidari, B.I. and Radder, B.M. (2007). Effect of FYM and fermented liquid manures on nutrients uptake by chilli (*Capsicum annum* L.) and soil nutrient status at harvest. *Journal of Asian Horticulture*. 4: 19-24.

Potkile, S.N., Bhale, V.M., Deshmukh, J.P., Dandge, M.S. and Choudhary, A.A. (2017). Nutrient management through organic sources in soybean-wheat cropping sequence under irrigated condition. *International*

Journal of Pure Applied Bioscience.  
5(5):1035-1041.

Ravishanker, N., Sharma, S.K., Singh, D.K.,  
Panwar, A.S. and Kumar, V. (2017).  
Organic Farming in India: Production issues  
and strategies; In: Organic farming crop  
production guide, network project on  
organic farming, ICAR, Indian Institute of  
Farming System Research, Modipuram,  
Meerut, Uttar Pradesh, India. pp-1-17.

Suchitra Rakesh, S. Poonguzhali, B.  
Saranya, S. Suguna and Jothibas, K. 2017.  
Effect of Panchagavya on Growth and Yield  
of *Abelmoschus esculentus* cv. Arka  
Anamika. International Journal of Current  
Microbiology and Applied Sciences.6(9):  
3090-3097.