

EFFECT OF seed storage conditions and duration STORAGE CONDITIONS AND DURATION OF STORAGE ON SEED GERMINATION OF *TETRAPLEURA TETRAPTERA* (Schum and Thonn) Taub AND *ANNONA MURICATA* (Linn.)

ABSTRACT

Aidan fruit (*Tetrapleura tetraptera*) and *Soursop* (*Annona muricata*) are medicinal plant species with great medicinal values. Despite their usefulness, preservation of their seed from harvest time to the time when they are required for planting has not been given adequate attention. Therefore, this study investigated the effect of seed storage conditions and duration of storage on seed germination of *T. tetraptera* and *Annona muricata*. Four storage conditions viz: cold room (-20 °C), refrigerator (5°C), room air conditioned (22°C) and laboratory (2°C) were used and the experiment was carried out in Ekiti State University, Ado - Ekiti for 6 months. Germination test was conducted on the stored seeds of the two species at the end of every month for the six months storage period at temperature ofunder.....light conditions for.....days In *Tetrapleura tetraptera*, highest percentage germination (50%) was recorded in seeds stored in the air conditioned room at 1 month. This was followed by seeds store in the refrigerator with 47% at 2 month. Seeds stored in the room air conditioned had percentage germination of 39% at 2 months, 30.50% at 3 months and 10.70% at 4 month of the experiment. No further germination in seeds under this treatment was observed at the last two months of the experiment. While seeds stored in refrigerator recorded 5.20% at 1 months of the experiment. Seeds stored in cold room and those in laboratory recorded no emergence throughout the period of the experiment. Also in *Annona muricata*, Highest percentage germination (58.00%) was observed in seeds stored in the room air conditioned at 1 month of the experiment, this was followed by the seeds in this same temperature at two month of the experiment with mean plumule emergence value of 39.00%, seeds under this temperature (air condition) did not respond to germination at 5th and 6th months of the experiment. Seeds of *Annona muricata* stored in the cold room responded to germination with unappreciable value of 2.05% only at third month of the experiment. Only 1.6% of the seeds stored in the laboratory emerged at 5th month of experimental time, seeds stored in the refrigerator did not respond for the first two months, last fifth and sixth month of the experiment but had 15.53% and 35.00% plumule emergence at third and fourth month of the experiment respectively. Thus, storage of seeds of *T. tetraptera* and *A. muricata* in air conditioned room with temperature ranges between 16 and 26°C for short period might enhance germination in these species.

Commented [MO1]: What were the germination conditions in relation to temperature and light?, which media were used, sand or paper? What was the duration of germination test

Commented [MO2]: Good trend, we expect seed germination to decrease over time

Commented [MO3]: Cold room has the lowest temperature, if the initial seed quality is good, we expect that the seeds should have better quality compared to other methods. Remember lowering the storage temperature while keeping other factors constant, extends the storage life of the seed

Commented [MO4]: In a seed storage experiment, there must be a baseline germination data ie germination at 0 month of storage/ seed germination before storage. This helps in tracking seed deterioration over time during storage

Commented [MO5]: Revisit your results before making this conclusion. You are concluding that storage at 22oC ie air conditioned is better than storage at -22oC and 2oC, scientifically this is not true. Seed stores better at lower temperature and vice versa.

What were your initial germination percentages before storage? If we have this data, we can now tell how each of the storage conditions affects this initial germination. Example, If initial germination is 90%, then how does this value decrease during the six months of storage under different conditions for the two species

Keywords ~~Medicinal plant~~, Storage Conditions, Duration of Storage, *Tetrapleura tetraptera*, *Annona muricata*, [seed germination](#)

INTRODUCTION

Tetrapleura tetraptera (Schum and Thonn.) Taub. Commonly known as Aridan in South Western Nigeria is a medicinal plant of the family Mimosaceae. The plants is locally known as Uyayak in Ibiobio; Edeminang in Efik; Osakirisa or Oshosho in Igbo and Dawo in Hausa (all in Nigeria) (Ojewole and Adewumi, 2004). Its fruit is used for the management of convulsion, leprosy, inflammation, and rheumatism (Okwu, 2003; Ojewole and Adewumi, 2004; Jamwal *et al.*, 1972). [Annona muricata](#), called soursop due to the sweet and sour flavor of the large fruit, is a lowland tropical fruit-bearing tree in the Annonaceae family. Related species include cherimoya (*A. cherimola*) and sugar-apple (*A. squamosa*). Other common names include graviola and guanábana (sometimes shortened to guanába). The genus name, 'Annona' is from the Latin word 'anon', meaning 'yearly produce', referring to the production of fruits of the various species in this genus (NRCS, 2008). *Annona muricata* is a small, upright, evergreen tree. Its young branches are hairy.

Commented [MO6]: What are the medicinal properties in the two species?

Formatted: Font: Bold, Font color: Text 1

The post-harvest processes of medicinal plants have great importance in the productive chain, ~~and its~~ [directly](#) influence ~~on~~ the quality and quantity of the active principles. Among the post-harvest processes, drying and storage are essential to maintain the product with the physical and chemical characteristics closer to those found in the fresh plant, and failures in any of these steps influence the quality of the final product. Drying, if not performed properly, can compromise the content of active ingredients while, incorrect storage can lead to loss of material, whether for physical or biological reasons (Martinazzo, 2006; Martinazzo *et al.*, 2007). Medicinal plants are often stored for long periods, before being used as raw material for the manufacture of various products (Masand *et al.*, 2014). However, improper storage can result in physical, chemical and microbiological changes (Peter, 2006; Mayuoni-Kirshinbaum, *et al.*, 2013).

Seed viability, ~~is~~ [is](#) the ability of the embryo to germinate, ~~and it~~ [is](#) affected by a number of factors including temperature, light, oxygen, water and species type. Germinability which is determined by germination percentage is the proportion of seeds that germinate from seeds subjected to the right conditions for growth while the germination rate is the speed with which the seeds germinate and is affected by seed viability, dormancy and environmental effects that

impact on the seed and seedling (Zamora, 2014). This is given as a percentage of germination over a certain amount of time, for example 90% germination within 20 days. Germination energy is the proportion of seeds in a given sample which germinate within the time of peak germination, generally taken as the highest number of germinations in a 24-hour period (Mrda et al. 2011). The interest in germination energy is based on a theory that only those seeds which germinate rapidly and vigorously under favourable conditions are likely to produce vigorous seedlings in field conditions, whereas weak seeds or delayed germination is often fatal (Aldhous 1972).

Simple techniques have been adopted to maintain the seed viability in both domesticated and wild sources (Ellis et al. 1992, Vertucci and Roos 1991). Seed deterioration and loss of viability, is a natural phenomenon occurring during storage (Nasreen et al. 2000, Schmidt. 2002) and inappropriate storage ~~condition-medium~~ such as room temperature storage (Hezewijk et al. 1993, Müller et al. 2011) has often resulted in low seed germination. Several factors, including temperature, nature of the seeds, seed moisture content, ~~and~~ relative humidity influence the seed longevity during storage (Onyekwelua and Fayose 2007, Pradhan and Badola 2008). Proper storage conditions, may effectively retain substantial viability in seeds over a considerable storage period (Butalo and Badola 2004, Chen et al. 2007, Pradhan and Badola 2008).

A long-term storage of seeds, especially under unfavourable conditions, leads to loss of viability. The nature of this physiological damage is variable, e.g. short-term deterioration in the field is different from long term deterioration during storage, which in turn is different from mechanical damage (Mrda et al. 2011). All parts of the seed such as the seed coat, which is of maternal origin and acts as a physical and chemical barrier, and the embryo are susceptible to physiological damage. Seeds are grouped based on their storage characteristics as orthodox and recalcitrant. Orthodox seeds are those that can be dried to moisture contents of 10% or less, in this condition they can successfully be stored at subfreezing temperatures (Onyekwelua and Fayose 2007). Recalcitrant seeds are seeds that do not survive drying and freezing during exsitu conservation and cannot resist the effects of drying or temperatures less than 10°C; thus, they cannot be stored for long periods like the orthodox seeds because they can lose their viability. Plants that produce recalcitrant seeds include avocado, mango, mangosteen, lychee, cocoa, rubber tree, some horticultural trees, and several plants used in traditional medicine such as species of *Virola* and *Pentaclethra* (Pradhan and Badola 2008).

Seeds of different plant species lose viability at varying degrees. For example, onion seed is very difficult to store while barley seed maintains good germination under a variety of storage conditions (Zamora 2014). Seeds and especially oilseeds including *E. bussei* contain large quantities of carbon stored as carbohydrate, lipid and protein (Bewley and Black 1985). The chemical composition of oilseeds which are rich in lipids causes specific oxidative processes to occur during storage and these are said to cause limited longevity depending on their specific chemical composition (Balešević et al. 2007).

Commented [MO7]: The background provided is good. However, the authors have not brought out the storage challenges of the two species under study. Has there been previous reports about seed storage of the two species in relation to temperature and relative humidity or duration of storage?

MATERIALS AND METHODS

Seed Source

Fresh fruits of *Tetrapleura tetraptera* were obtained from Oba's market, Ado-Ekiti, Ekiti State, Nigeria and were taken to the herbarium of the Department of Plant Science and Biotechnology, AdoEkiti, Nigeria for authentication. While the mature fruits of *Annona muricata* were collected from the parent plant from its natural habitat in Ipoti-Ekiti located in Ijero Local Government Area of Ekiti State.

Formatted: Font: Italic

Viability Test

Seeds were subjected to viability test by using 2,3,5 Triphenyl Tetrazolium Tetrasolium chloride according to Kruse et al., (2004).

Commented [MO8]: Describe the maturity status of the fruit. This can be done using morphological markers. The maturity status of the fruit can give idea of the seed maturation stage

Procedure

Two hundred seeds of *Tetrapleura tetraptera* and *Annona muricata* were divided into four groups consisting of 50 seeds each. Each batch were stored at four temperatures viz: cold room (-20°C), refrigerator (5°C), air conditioner (22°C) and laboratory/room temperature (2°C) as control. 20 seeds each of *Tetrapleura tetraptera* and *Annona muricata* from the four batches, stored at different temperatures were collected and planted to test for the germinability on monthly basis for six months. Planting was done in polythene nylons filled with top-soil and watered daily. The experiments were replicated four times. The number of seeds that germinate in each treatment was recorded and the percentage germination was calculated.

Commented [MO9]: Which concentration, 1% 0.1%????

Commented [MO10]: 20 years ago????? It is important to describe your own procedure. For the goodness of your paper, tell us what you did with the TZ test. It is better than referring to a 20 yr old paper

Commented [MO11]: What were the germination conditions with respect to temperature and light???

Commented [MO12]: What did you consider as your germinated seed? Radicle protrusion or emergence of plumule above the soil medium

RESULTS

Effect of Storage Conditions and Duration of Storage on Seed Germination of *T. tetraptera*

Table 1 shows that various storage condition and duration had different effect on seed germination of *T. tetraptera*. Results obtained revealed that seeds stored in cold room and those stored in the laboratory did not germinate throughout the period of experiment. Seeds stored in the refrigerator had mean germination of 5.20% and 47.00% for the first and second months experiment respectively and shows no further germination at 3rd, 4th, 5th and 6th months of the experiment. Highest seed germination (50%) was recorded in seeds under air condition at the first month of the experiment, mean germination of 39%, 30.50% and 10.70% were observed at 2nd, 3rd and 4th month of the experiment respectively with no further germination at 5th and 6th month in this same (air condition) treatment.

Table 1: Percentage Germination of *T. tetraptera* seeds stored under different storage conditions

Months	Percentage Germination (%)			
	Cold room	Refrigerator	Air conditioned	Room Laboratory
1 st	0	5.20	50.00	0
2 nd	0	47.00	39.00	0
3 rd	0	0	30.50	0
4 th	0	0	10.70	0
5 th	0	0	0	0
6 th	0	0	0	0

Commented [MO13]: Normally, seed germination is higher during the first month of storage and decreases during the storage period.

Commented [MO14]: This is a good result. I expect all your results to follow a similar trend for all storage conditions, but the germination will vary during the storage periods under different conditions.

Commented [MO15]: A graph would present it better

Commented [MO16]: Provide germination data at 0 month of storage. This result is not expected for a cold room.

Commented [MO17]: Why this trend. What was the germination at 0 month of storage

Effects of Storage Conditions and Duration of Storage on Seed Germination of *A. muricata*

Table 2 shows that seeds of *Annona muricata* stored in the cold room responded to germination with unappreciable value of 2.05% only at third month of the experiment, while no emergence was observed at first two months and the last three months of the experiment in this treatment. Only 1.6% of the seeds stored in the laboratory emerged at 5th month of experimental time, seeds stored in the refrigerator did not respond for the first two months, last fifth and sixth month of the experiment but had 15.53% and 35.00% plumule emergence at third and fourth month of the experiment respectively.

Highest percentage germination (58.00%) was observed in seeds stored in the room air conditioned at one month of the experiment, this was followed by the seeds in this same temperature at two month of the experiment with mean plumule emergence value of 39.00%, a decrease in the percentage plumule emergence in seeds under this same temperature was observed at 3rd and 4th months of the experiment with 10.50% and 9.82% respectively. While the seeds under this temperature (air condition) did not respond to germination at 5th and 6th months of the experiment.

Table 2: Percentage Germination of *A. muricata* seeds stored under different storage conditions

Months	Percentage Plumule Emergence (%)			
	Cold room	Refrigerator	Air conditioned	Room Laboratory
1 st	0	0	58.00	0
2 nd	0	0	39.00	0
3 rd	2.05	15.53	10.50	0
4 th	0	35.00	9.82	0
5 th	0	0	0	1.6
6 th	0	0	0	0

Commented [MO18]: Something isn't true with this data.

Commented [MO19]: Good trend. The rest of the conditions must have a decreasing trend, with germination highest at the beginning months

DISCUSSION

Results from this study emphasized the fact that appropriate storage conditions and duration of storage are of great necessity in preserving the viability of seeds from the time of collection to the time when they are needed for sowing. In fact, Genes and Nasreen (1995) observed that the use of seed is one of the propagation methods that can be used to multiply plants but the seeds need to be capable of germinating in the first place., hence the need for knowledge of its methods of germination. Since most plants produce seeds only once a year, knowledge of methods of seeds preservation for next season's generation was deemed important and so was the determination of how soon the seeds would lose its viability.

The results obtained in this study revealed that seeds of *T. tetraptera* and *A. muricata* stored in air conditioned room for short period had the best percentage germination when compared with seeds in other storage conditions. Reason for this could be that temperature in other storage conditions might be too low or high to promote seed germination in this plant or that they were too high to enhance proper germination of this species. This finding agreed with previous observation of Lars (2000) that seeds germinate faster with increasing air temperatures up to a point. Extreme heat will slow germination and also increase moisture loss. The temperatures for optimal germination vary with the type of plant, some annual flowers and vegetables are extremely sensitive to cold, extremely hot or cold temperatures can hamper plant growth, as well as affect seed germination Lars (2000). Gamaene *et al.*, (1999) ascertained that under no storage condition would recalcitrant seed be kept for long as this tends to reduce its viability.

CONCLUSION

Storage conditions with extremely hot or cold temperatures does not encourage seed germination of this plant species. Therefore, storage of seeds of *T. tetraptera* in air conditioned room with temperature ranges between 16 and 26°C for short period to keep its viability intact is recommended

References

- Aldhous, J. R. (1972) Nursery practice. Forestry Comm. Bull. No. 43, London
- Balešević, S., Tatic, M., Miladinovic, J. and Pucarevic, M. (2007). Changes of fatty acids content and vigor of sunflower seed during natural aging, *Helia*, 30 (47): 6168.

Commented [MO20]: This is applicable in controlled germination chambers to determine the best temperature for seed germination. This is not for storage.

Commented [MO21]: The purpose of storage at low temperature is to maintain seed viability for long periods. This is what is done in the gene banks. However, your study shows that the seed did not germinate under cold storage. Were the seeds viable before storage? I don't see the results of TZ test

Commented [MO22]: Purpose of storage is to keep seed alive for long periods

Commented [MO23]: Your experiment is not looking at optimum temperature for seed germination. The focus of your study is on optimum temperature for seed storage. Which storage temperature allows the seed to maintain high viability for long period

Commented [MO24]: This conclusion and recommendation is not true. Lower storage temperature is recommended for seed storage. Seeds stored at 26°C will deteriorate faster than those stored at lower temperature. If you are in charge of the germplasm conservation unit, would you recommend seed storage at higher temperature (26) or lower temperature (20°C)

- Bewley, J. D. and Black, M. (1985). Seed physiology of development and germination. Plenum Press, New York.
- Butalo, J. S. and Badola, H. K. (2004). Effect of pre-sowing treatment on seed germination and seedling vigour in *Angelica glauca*, a threatened medicinal herb. *Cur. Sci.* 87 (6):796-799.
- Chen, S. Y., Kuo, S. R. and Chien, C. T. (2007). Storage behaviour of seeds of *Cinnamomum osmophloem* and *Neolisea aciculate varivariabiliana* (Lauraceae). *Seed Sci. Technol.* 35 (1): 237-243.
- Ellis, R. H., Hong, T. D. and Roberts, E. H. (1992). The low-moisture-content limit to the negative logarithmic relation between seed longevity and moisture content in three sub species of rice. *Annal. Bot.* 69(1):53-58.
- Gamaene, S. H. W., Pritchard, C. and Harris, A. (1999). *Vitellaria paradoxa*. Handling and storage of recalcitrant and intermediate tropical forest tree seeds, Newsletter 5: 15-16. Danida Forest Seed Centre, Humlebaek, Denmark.
- Genes, F. and Nyomora, A. M. S. (2018). Effect of storage time and temperature on germination ability of *Escoecaria bussei*. *Tanzania journal of science* 44 (1); 123-133.
- Hezewijk, M. J., Van, B. A. P., Verleij, J. A. C. and Pieterse, A. H. (1993). Germination of *Orobanche crenata* seeds as influenced by conditioning temperature and period. *Can. J. of Bot.* 71(6):786-792.
- Jamwal, K. S., Sharma, M. L., Chandhoke, N. R. and Chatak, B. J. (1972). Pharmacological action of 6,7- Dimethoxycoumarin (Scoparone) isolated from *Artemisia scoparia* Waldst & Kit. *Indian J. Med. Res.* 60(5):763.
- Lars, S. (2000). Guide to handling of tropical and subtropical forest seed. Centre, Humlebaek, Denmark.
- Martinazzo, A. P. (2006). Secagem, armazenamento e qualidade de folhas de *Cymbopogon citratus* (D.C.) Stapf. 2006. 156 f. Tese (Doutorado em Engenharia Agrícola) – Departamento de Engenharia Agrícola, Universidade Federal de Vicosa, Minas Gerais.
- Martinazzo, A. P., Correa, P. C., Melo, E. C. and Barbosa, F. F. (2007). Difusividade efetiva em folhas de *Cymbopogon citratus* (DC.) Stapf submetidas a secagem com diferentes comprimentos de corte e temperaturas do ar. *Revista Brasileira de Plantas Mediciniais* 9(1): 68-72.

- Masand, S., Madan, S. and Balian, S. K. (2014). Modern concept of storage and packaging of raw herbs used in Ayurveda. *International Journal of Research in Ayurveda and Pharmacy* 5(2): 242 -245.
- Mayuoni-Kirshinbaum, L., Daus, A. and Porat, R. (2013). Changes in sensory quality and aroma volatile composition during prolonged storage of 'Wonderful' pomegranate fruit. *International Journal of Food Science and Technology* 48: 1569-1578.
- Mrđa, J., Crnobarac, J., Dušanić, N., Jocić, S. and Miklič, V. (2011). Germination energy as a parameter of seed quality in different sunflower genotypes - *Genetika* 43(3): 427-436.
- Muller, E., Cooper, E. J. and Alsos, I. G. (2011). Germinability of arctic plants is high in perceived optimal conditions but low in the field. *Botany* 89 (5):337-348.
- Nasreen, S. (1995). Effects of storage period and temperature on seed viability of Wheat. *Park J. Bio Sci.* 2: 1492 – 1493.
- Nasreen, S., Khan, B. R. and Mohmad, A. S. (2000). The effect of storage temperature, storage period and moisture content on seed viability of soyabean. *Pakistan J. of Biol. Sci.* 12/2000; 3:12
- Natural resources Conservation Service (2008). Plants profile *Annona squamosa* L. United States Department of Agriculture, 4 – 17.
- Ojewole, J. A. and Adewunmi, C. O. (2004). Anti-inflammatory and hypoglycemic effects of *Tetrapleura tetraptera* (Taub) (Fabaceae) fruit aqueous extract in rats. *J. Ethnopharmacol* 95(23). 177-182.
- Okwu, D. E. (2003). The potentials of *Ocimum gratissimum*, *Pergularia extensa* and *Tetrapleura tetraptera* fruit as spice and flavouring agents. *Nig. Agric J.* 35: 143-148.
- Onyekwelu, J. C. and Fayose, O. J. (2007) Effect of storage methods on the germination and proximate composition of *Treulia africana* seeds. In: Proceedings of the International Conference on Agricultural Research for Development; October 2007; University of KasselWitzenhausen and University of Göttingen, Tropentag
- Peter, K. V. (2006). Handbook of Herbs and Spices. 3. New York, USA.
- Pradhan, B. K. and Badola, H. K. (2008). Seed germination response of populations of *Swertia chirayita* following periodical storage. *Seed Technol.* 30(1):63–69.
- Schmidt, L. (2002). Guide to handling of Tropical and Subtropical forest seeds Denmark: DANIDA Forest Seed Centre, Borch Tyrk A/S.

Vertucci, C. W. and Roos, E. E. (1990). Theoretical basis of protocols for seed storage. *Plant Physiol.* 94(3):1019–1023.

Zamora, F. J. H. (2013). Effect of moisture on lentil seed germination and growth. <http://en.wikipedia.org/wiki/Germination>.

UNDER PEER REVIEW