

Original Research Article

Assess Nurse's Knowledge Toward Venous Thrombosis At Al-Sharqat General Hospital

ABSTRACT

Background and aims: A blood clot that is stuck in a deep vein is known as deep vein thrombosis, while a blood clot that has broken loose and migrated to the lungs is known as a pulmonary embolus. Thrombotic disorders are the world's most common cause of high mortality and morbidity among preventable diseases.

was to assess nurse's knowledge toward venous thrombosis at AL-Sharqat General Hospital and relationship between knowledge and their demographic data.

Study design: A descriptive study.

Place and Duration of Study: Sample: This study was conducted at Al-Sharqat General Hospital in critical care unit and surgical department, between November 2023 and May 2024

Methodology: A purposive sample (non-probability) consisting of (25) nurses was selected from nurses who work in the critical care units and surgical department.

Results: The results showed that over half of the participants were female nurses, making up the majority of the group. These nurses had graduated from nursing institutes and had between one and five years of experience, without having attended any training courses. Results show that nurses knowledge related to venous thrombosis was poor to moderate and there were no significant relationship between nurse's knowledge and their demographic data at P-value >0.05

Conclusion: Nurses working in the surgical department, emergency units, and critical care units had poor knowledge regarding venous thrombosis because their education level was low and they had not attended training courses

Keywords: Nurse, knowledge, venous thrombosis.

1. INTRODUCTION

A blood clot that is stuck in a deep vein is known as deep vein thrombosis, while a blood clot that has broken loose and migrated to the lungs is known as a pulmonary embolus (1).

Thrombotic disorders are the world's most common cause of high mortality and morbidity among preventable diseases. Deep vein thrombosis, which is regarded as the third main vascular diagnostic after heart attack and stroke, is the most serious and dangerous vascular condition among patients(2) .

DVT has an annual impact of up to 900,000 people (3) ,according to current estimates, DVT causes 60,000 to 100,000 deaths in Americans each year. The fact that 10 to 30% of people who develop DVT die within a month of their

diagnosis is even more concerning (3) .

The majority of DVT cases go undetected and are regarded as silent killers. It is a deadly disorder that has negative effects on patients. Therefore, strategies to enhance nurses' utilization of DVT prevention procedures are crucial(4).

Despite having several risk factors, DVTs are preventable. Understanding risk factors and preventive actions is essential for DVT prevention, claim(5).

Understanding preventative steps can not only lower the incidence of DVT problems but also shorten hospital stays, free up beds, and even save lives. Nurses can significantly contribute to the prevention of Deep Venous Thrombosis, and training will enhance their skills and practices(6).

Nurses lack the necessary skills and knowledge to avoid deep vein thrombosis (DVT) (7). A recent study in India shows that a high percentage of nurses had bad practices and knowledge regarding the prevention of venous thrombosis.(8).

Aims of the study

A. The primary objective

- 1- Was to assess nurse's knowledge toward venous thrombosis.
- 2- Relationship between nurse's knowledge and their demographic data.

B. The secondary objective

1. Identifying the relationships between the Nurses' knowledge with Nurses' staff socio-demographic characteristics variables, such as (age, gender, level of education, years of experience and received training courses).
2. To identify socio-demographic characteristics of nurse.

2. METHODOLOGY

A descriptive study is conducted for the periods of 11th November 2023 to 3ed May 2024. Based on the approval of the council of Nursing College/University of Mosul the study is conducted, the researcher then submitted an outline of the study, including its objectives, to the Salah-Elden Health Directorate (Planning Department, Section of Health Research). This study was conducted at Al-Sharqat General Hospital in critical care unit and surgical department . Whether they agreed to participate in the study or declined, individuals in study groups were given the opportunity to provide informed consent, which was provided to them. A purposive sample (non-probability) consisting of A total of (30) Nurses who are working in surgical department , critical care units and emergency units at the AL-Sharqat General Hospital in AL-Sharqat City. (5) nurses refuse to participate. To make the instrument more valid, it was presented to a panel of (10) experts in the different fields, and Reliability Coefficient for the Pilot study through using Al-Naqeeb Formula[Ref]:

$$\text{Reliability value} = \left(1 - \frac{\text{no. of non coincidences items}}{\text{no. of all items} * \text{sample size of pilot study}} \right) * 100\%$$

. The item was rated and score according to the following designs:

1) The evaluation of the respondent for (Evaluation Effectiveness of an Educational Program on the nurse's knowledge toward a venous thrombosis each question was scored with (2) True answer and (1) False answer.

2) The higher grade of scoring of the questionnaire (MS) means higher knowledge. Data of the study were ordered according to three levels of scale (High, Moderate, and Low) according to the following:

□ low = L(0.00-33.33)

□ Moderate =M (33.34-66.66)

□ High= H (66.67-100)

includes a summary statistics, such that (Mean of Score-MS, Standard Deviation-SD, Relative Sufficiency-RS%, Assessing scored over the three intervals due to RS% by the following assessments: [Low: (0.00 _ 33.33); Moderate: (33.34 _ 66.66); , and High: (66.67 _ 100)].

The self-report and questionnaire are used in the data gathering procedure, and descriptive and inferential statistics are used for analysis.

Statistical Data Analysis approaches were used in order to analyze and assess the results of the study under application of the statistical package (SPSS) ver. (22.0). **Descriptive data analysis:** Summary Statistics tables including: Observed Frequencies, Percents, Mean of score (MS), Standard Deviation (SD), Relative Sufficiency (RS%), Percentile Grand/and Global Mean of Score (PGMS), and Percentile Pooled Standard Deviation (PPSD).

A questionnaire consists of the following parts :

- 1- part one :socio-demographic data
- 2- part two : : This section, which was composed of one domain toward venous thrombosis. includes (20) closed ended (multiple choice) questions to measure nurses' knowledge about venous thrombosis, including: definition, causes, risk factors, signs & symptoms, diagnosis, treatment and complication.

3. RESULTS AND DISCUSSION

3.1. Discussion of Nurses' the Socio-demographic characteristics in the Study Groups.

Twenty five nurses has been selected for each group, total number of male in the study group 11(44%), mean age (29.40 ± 5.97), whereas 44(49%) are accounted in controlled, with mean age (28.92 ± 5.35), as well as no significant differences at P>0.05 are accounted between studied groups with respect to gender and age variables.

Most of educational levels of the studied groups has assigned institute graduation, since 15(60%), and 16(64%) are accounted in the study and controlled groups respectively, with no significant different between studied groups at P>0.05, as well as surgical halls are accounted 10(40%) in each group, and the leftover were distributed similarly among emergency hall, and internal resuscitation unit.

Table (1): Distribution of the studied groups according to (SDCv.) with comparisons significant

SDCv.	Groups	Groups				C.S. (*)
	Classes	Study		Control		P-value
Gender	Male	11	44	12	48	C.C.=0.040
	Female	14	56	13	52	P=0.777 (NS)

Age Groups	< 30 yrs.	6	24	7	28	C.C.=0.207 P=0.525 (NS)
	30 _ 34	10	40	9	36	
	35 _ 39	4	16	7	28	
	> 39 yrs.	5	20	2	8	
	Mean ± SD	29.40 ± 5.97		28.92 ± 5.35		
Education levels	Nursing preparatory school graduate	1	4	0	0.00	C.C.=0.210 P=0.509 (NS)
	Institute graduate	15	60	16	64	
	College graduate	6	24	8	32	
	Postgraduate	3	12	1	4	
Occupation	Emergency hall	7	28	8	32	C.C.=0.052 P=0.936 (NS)
	Internal resuscitation unit	8	32	7	28	
	Surgical halls	10	40	10	40	

(*) **NS: Non Sig. at P>0.05; Testing based on a contingency coefficient (C.C.) test.**

According to analyses of the nurses' demographic data, however there are notable distinctions. The study's quasi-experimental approach, which included both a pre- and post-test, accepts these findings.

The study's objective was to assess nurses' knowledge regarding venous thrombosis .

According to Table 1. With respect to gender, the current study's findings on nurse gender showed that most nurses were female. This might be because there are often fewer male nurses than female nurses in the nursing field.

These results corroborate those of (9) , who performed descriptive research in Egypt to " assess attitude, subjective norms, perceived behavioral control, and intention of nurses towards prevention of deep vein thrombosis among critically ill patients in intensive care units". The research sample mainly consisted of female nurses more than half.

Regarding years of experience, a high percentage of them (more than a third) had been working in nursing for 1–5 years. Most of them (more than half) have experience in the intensive care unit, surgical units, and emergency department for 1–5 years.

These results agreed with study conducted by (10) "*Critical Care Nurses' Knowledge about Pulmonary embolism in Respiratory Care Unit in Baghdad Teaching Hospitals*" A high percentage of them was more than a third For (1-5) years they worked in nursing, and more than half of them had (1-5) years of experience At RCU.

Concerning educational level, The majority of the nurses in the research sample worked in the critical care unit and held a diploma in nursing.

These results agree with study conducted by (11)" Effectiveness of Nursing Education Program on Nurses Practices Toward Arrhythmia in Kirkuk" stated that the nursing institute constituted the majority of his study sample.

This outcome might be explained by the fact that technical nurses work as bedside nurses in government hospitals, but the majority of bachelor nurses serve as supervisors or head nurses.

As regards age, the results of this study indicate that the majority of the study and control samples were in the 30- to 34-year-old age range, with mean ages of (29.40 ± 5.97) and (28.92 ± 5.35) for the study and control groups, respectively. In the researcher's opinion, this outcome might be the result of the demanding nature of work in critical regions, where young adult nurses are needed to combine their severe work load with their energy and strength.

This results disagreed by (12)" *Effectiveness of an Educational Program upon nurses ' knowledge toward The Continuous Positive Airway Pressure (CPAP) Machine in Neonatal Intensive Care Unit at Al-Diwanyia City Hospitals*", he stated that the majority of his study samples ranged in age from 25 to 30

3.2: Discussion of Nurses' Knowledge Toward Venous Thrombosis At Al-Sharqat General Hospital.

Table (2) includes a summary statistics, such that (Mean of Score-MS, Standard Deviation-SD, Relative Sufficiency-RS%, Assessing scored over the three intervals due to RS% by the following assessments: [Low: (0.00 _ 33.33); Moderate: (33.34 _ 66.66); , and High: (66.67 _ 100)]. In addition to that, testing of significant are obtained for each period in light of pre to post periods, as well as testing of significant are presented for testing observed frequencies between the studied groups in each period independently.

Results of testing significant with reference of studied items, as well as scoring scales assessments concerning effectiveness of applying the proposed program reported highly significant differences at P<0.01 through raising grades of studied respondents at the post period, and that could be enable to confirms importance and successfulness of applying the proposed program. In addition to that, and rather than testing significant are too sensitive to improvements that might be occurred for repeated measurements statistically in study group, but all of studied items illustrated too highly and meaningful changeability with a high levels of assessed along pre to post periods.

The results of the controlled group has recorded completely immovable responses over the studied periods with a moderate level of assessed generally.

Table (2): Descriptive Statistics of the studied groups according to (Knowledge related to Venous Thrombosis) Domain along studied periods with comparisons significant

Knowledge related to Venous Thrombosis	Period	N o.	Study					Control					C.S. (*)
			MS	SD	RS %	Ass. (*)	C.S. (*)	MS	SD	RS %	Ass. (*)	C.S. (*)	
Is it considered one of the symptoms of deep vein thrombosis?	Pre	25	0.72	0.46	72	H	1.00	0.76	0.44	76	H	0.019	1.000
	Post	25	0.72	0.46	72	H	0	0.32	0.48	32	L	0.010	0.010
What are the signs and symptoms of PE that a patient may exhibit?	Pre	25	0.72	0.46	72	H	1.00	0.56	0.51	56	M	0.180	0.377
	Post	25	0.72	0.46	72	H	0	0.32	0.48	32	L	0.010	0.010
What is the difference between venous thrombosis (DVT) and arterial thrombosis?	Pre	25	0.80	0.41	80	H	1.00	0.96	0.20	96	H	0.000	0.189
	Post	25	0.84	0.37	84	H	0	0.48	0.51	48	M	0.016	0.016
What is the relationship of Restless Leg Syndrome to DVT?	Pre	25	0.80	0.41	80	H	1.00	0.32	0.48	32	L	0.227	0.001
	Post	25	0.80	0.41	80	H	0	0.52	0.51	52	M	0.072	0.072
What is the difference between deep phlebitis (DVT) and superficial phlebitis?	Pre	25	0.48	0.51	48	M	0.125	0.12	0.33	12	L	0.227	0.012
	Post	25	0.64	0.49	64	M	5	0.32	0.48	32	M	0.046	0.046
What is the relationship	Pre	25	0.52	0.5	52	M	0.03	0.3	0.4	36	M	1.00	0.393

between cancer and the risk of developing DVT?				1			1	6	9			0	
	Post	25	0.76	0.4 4	76	H		0.4 0	0.5 0	40	M		0.021
The common diagnostic technique used to detect Pulmonary Embolism is?	Pre	25	0.52	0.5 1	52	M	0.12 5	0.6 4	0.4 9	64	M	0.42 4	0.567
	Post	25	0.68	0.4 8	68	H		0.4 8	0.5 1	48	M		0.252
What medical tests can be used to determine the effectiveness of anticoagulant treatment?	Pre	25	0.68	0.4 8	68	H	0.06 3	0.5 2	0.5 1	52	M	0.75 4	0.387
	Post	25	0.88	0.3 3	88	H		0.4 4	0.5 1	44	M		0.002
What risk factors may increase the likelihood of blood clots forming?	Pre	25	0.72	0.4 6	72	H	0.03 1	0.3 2	0.4 8	32	L	0.01 3	0.010
	Post	25	0.96	0.2 0	96	H		0.7 2	0.4 6	72	H		0.049
What age group is most susceptible to DVT?	Pre	25	0.72	0.4 6	72	H	0.50 0	0.6 8	0.4 8	68	H	0.45 4	1.000
	Post	25	0.80	0.4 1	80	H		0.5 2	0.5 1	52	M		0.072
What are the main factors that may increase the risk of venous thrombosis?	Pre	25	0.52	0.5 1	52	M	0.03 1	0.5 6	0.5 1	56	M	0.06 5	1.000
	Post	25	0.76	0.4 4	76	H		0.2 8	0.4 6	28	L		0.002
The main factor contributing to the formation of venous thrombosis is?	Pre	25	0.76	0.4 4	76	H	1.00 0	0.2 8	0.4 6	28	L	0.11 8	0.002
	Post	25	0.80	0.4 1	80	H		0.5 6	0.5 1	56	M		0.128
What is the treatment method for DVT?	Pre	25	0.84	0.3 7	84	H	0.50 0	0.4 4	0.5 1	44	M	0.30 2	0.007
	Post	25	0.92	0.2 8	92	H		0.6 4	0.4 9	64	M		0.037

Continue ...

Knowledge related to Venous Thrombosis	Period	N o.	Study					Control					C.S. (*)
			MS	SD	RS %	Ass. (*)	C.S. (*)	MS	SD	RS %	Ass. (*)	C.S. (*)	
What is the first step in managing a PE emergency?	Pre	25	0.68	0.4 8	68	H	0.06 3	0.1 2	0.3 3	12	L	0.00 1	0.000
	Post	25	0.88	0.3 3	88	H		0.6 4	0.4 9	64	M		0.095
What is the main role of surgery in treating DVT?	Pre	25	0.52	0.5 1	52	M	0.01 6	0.4 4	0.5 1	44	M	0.77 4	0.778
	Post	25	0.80	0.4 1	80	H		0.5 2	0.5 1	52	M		0.072
What emergency treatment can be used in severe and serious PE?	Pre	25	0.52	0.5 1	52	M	0.00 4	0.3 2	0.4 8	32	L	0.50 8	0.252
	Post	25	0.88	0.3 3	88	H		0.4 4	0.5 1	44	M		0.002
One of the common ways to prevent deep vein thrombosis is?	Pre	25	0.80	0.4 1	80	H	0.50 0	0.2 8	0.4 6	28	L	0.00 7	0.001
	Post	25	0.88	0.3 3	88	H		0.7 2	0.4 6	72	H		0.289
How can proper nutrition play a role in preventing DVT?	Pre	25	0.56	0.5 1	56	M	0.00 8	0.4 4	0.5 1	44	M	0.75 4	0.572
	Post	25	0.88	0.3 3	88	H		0.3 6	0.4 9	36	M		0.000
One of the factors that may contribute to the	Pre	25	0.84	0.3 7	84	H	1.00 0	0.5 2	0.5 1	52	M	1.00 0	0.032

development of DVT complications is?	Post	25	0.88	0.33	88	H		0.48	0.51	48	M		0.005
What is one of the possible complications of PE?	Pre	25	0.52	0.51	52	M	0.004	0.79	0.41	79	H	0.057	0.072
	Post	25	0.88	0.33	88	H		0.44	0.51	44	M		0.002

(*) Assessments Intervals Scoring Scales of Relative Sufficiency Coefficient (RS%): [L: Low (0.00– 33.33)]; [M: Moderate (33.34 – 66.66)]; [H: High (66.67 – 100)].

Testing are based on McNemar test for testing Pre to Post periods in each group, and Fisher Exact Probability test for testing 2X2 association category dichotomous binary nominal scales of two independent random variables.




Figure (1): Cluster Bar Charts for distribution Grand Mean of Score of studied Main and Sub Main Domains along studied pre to post periods for the studied groups

The study group's pre- and post-test results showed notable variations, but overall, they demonstrated a good degree of understanding in the majority of the topics related to venous thrombosis, indicating that the training program had an impact on the nurses' knowledge. In addition to that, testing of significant are obtained for each period in light of pre to post periods, as well as testing of significant are presented for testing observed frequencies between the studied groups in each period independently.

Results of testing significant with reference of studied items, as well as scoring scales assessments concerning effectiveness of applying the proposed program reported highly significant differences at $P < 0.01$ through raising grades of studied respondents at the post period, and that could be enable to confirms importance and successfulness of applying the proposed program.

In addition to that, and rather than testing significant are too sensitive to improvements that might be occurred for repeated measurements statistically in study group, but all of studied items illustrated too highly and meaningful changeability with a high levels of assessed along pre to post periods. The results of the controlled group has recorded completely immovable responses over the studied periods with a moderate level of assessed generally. Table (3)

The researcher thinks a majority of them have nursing diplomas, all nursing texts are written in English, and they are studying Arabic, . Additionally, the majority of them did not receive previous education on venous thrombosis treatment, which is another reason for their lack of expertise.

These results supported by (13) " *Effect of An Educational Program on Critical Care Nurses Performance regarding Emergency Care for Patients with Pulmonary Embolism*". the study demonstrated the nurses' knowledge prior to the program's deployment, generally poor.

Also these results agree with (14) showed that following an organized education program, there was a significant rise in the post-test knowledge scores. One of the best ways to increase nurses' knowledge regarding deep vein thrombosis prevention is through an organized teaching program.

The study is in agreement with the study done by (15) who claimed that the program improves nurses' knowledge in the intensive care unit and showed a difference between the program's pretest and posttest.

3.3. Discussion of the association between the knowledge with Demographic Variables of the Study

Results shows that weak relationships are proved with (SDCv. and SRv.), since no significant relationships are accounted at $P > 0.05$, and according to that.

Table(3): Relationships (Analysis of Covariance-ANCOVA) for Compliance Regarding to Life Style Modification in the study group and SDCv.

Group	Source of Variations S.O.V.	Type III Sum of Squares	d.f.	Mean Square	F Statistic	Sig. Levels	C.S. (*)
Study Group	Intercept	46506.0	1	46506	1422.9	0.000	HS
	Gender	63.857	1	63.857	1.954	0.190	NS
	Age Groups	53.011	3	17.67	0.541	0.664	NS
	Education Level	12.963	3	4.321	0.132	0.939	NS
	Occupation	163.02	2	81.51	2.494	0.128	NS
	Services yrs.	22.512	2	11.256	0.344	0.716	NS
	Services yrs. at the specific felid	9.641	2	4.82	0.147	0.865	NS
	Error	359.5	11	32.683	R-Square=0.491		
Total	156063.1	25					

(*) HS: Highly Sig. at $P < 0.01$; Non Sig. at $P > 0.05$; Statistical hypothesis based on Analysis of Covariance (ANCOVA).

A The current study found no statistically significant correlation between a nurse's knowledge with venous thrombosis and demographic variables such age, gender, years of experience as a nurse, level of education attained, and years working in surgical or critical care units.

These results agree with (10) " *Critical Care Nurses' Knowledge about Pulmonary embolism in Respiratory Care Unit in Baghdad Teaching Hospitals*". which, in contrast to the current study, showed that there is no significant relationship between age, gender, years of experience, and nurses' knowledge, with the exception of the degree of education, where there is.

Also, these results agree with (Najlala et al.,2021) " *Effectiveness of Education Program on Nurses' Knowledge and Practice toward Preventive Measures of Pulmonary Embolism in AL Nasiriyah City*" show that the knowledge of nurses on preventive measures of pulmonary embolism and demographic characteristics (age, gender, years of service in nursing, educational level, and years of duty in critical or surgical units) did not vary statistically significantly.

Results showed that the overall main domains recorded high significant differences ($P < .001$) toward the effectiveness of a proposed program through raising the information grades of studied respondents at the post-period, and that could be used to confirm the success of applying a proposed program.

In addition to that, and rather than testing significant improvements that occurred for repeated measurement statistics in the study group, all the main domains studied were illustrated as highly and meaningfully changeable with high levels of assessments along the pre- and post-period.

4. CONCLUSION

Nurses working in the surgical department, emergency units, and critical care units had poor to moderate knowledge (low = L(0.00-33.33), (Moderate =M (33.34-66.66) , knowledge regarding venous thrombosis because their education level was low and they had not attended training courses.

CONSENT

All authors declare that 'written informed consent was obtained from the nurses for publication of this case report and accompanying images. A copy of the written consent is available for review by the Editorial office/Chief Editor/Editorial Board members of this journal."

ETHICAL APPROVAL

All authors hereby declare that all experiments have been examined and approved by the appropriate ethics committee and have therefore been performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.

REFERENCES

1. Silva JS da, Lee JA, Grisante DL, Lopes J de L, Lopes CT. Nurses' knowledge, risk assessment, and self-efficacy regarding venous thromboembolism. *Acta Paul Enferm.* 2020;33:eAPE20190125.
2. Parveen B, Kousar Parveen MH, Afzal M. NURSES KNOWLEDGE AND PRACTICES REGARDING RISK FACTORS AND PREVENTION OF PATIENTS DIAGNOSED WITH DEEP VEIN THROMBOSIS. *J Glob Biosci* Vol. 2021;10(6):8780–800.
3. Yohannes et al. (2022). Centers for Disease Control and Prevention (CDC). Obtenido Centers Dis Control Prev CDC <https://www.cdc.gov/dpdx/strongyloidiasis/index.html>. 2020;
4. Hebeshy MI, Hansen DM, Broome B, Abou Abdou S, Murrock C, Bernert D. Development of the Nurses' Intention to Use Deep Vein Thrombosis Preventive Measures Questionnaire. *J Nurs Meas.* 2020;
5. Grabo DJ, Seery JM, Bradley M, Zakaluzny S, Kearns MJ, Fernandez N, et al. Prevention of deep venous thromboembolism. *Mil Med.* 2018;183(suppl_2):133–6.
6. Al-Mugheed KA, Bayraktar N. Knowledge and practices of nurses on deep vein thrombosis risks and prophylaxis: A descriptive cross sectional study. *J Vasc Nurs.* 2018;36(2):71–80.
7. Luo Y, Zhou C, Deng Q, Xu W, Zhang X. Prevention of deep venous thrombosis in stroke: a best practice implementation project. *JBIC Evid Implement.* 2021;19(3):279–87.
8. Alyousef HA, Badawi SEA, Elghoneimy YA, Alameri RA, Almutairi AM. Nurses' Knowledge and Practices Regarding the Prevention of Deep Vein Thrombosis in Saudi Arabia: Descriptive Cross-Sectional Study. *Med Arch.* 2022;76(6):447.
9. Hebeshy MI. Attitude, subjective norms, perceived behavioral control, and intention of Egyptian nurses towards prevention of deep vein thrombosis among critically ill patients in intensive care units. Kent State University; 2018.
10. Najm MA, Jassim AH, Mohammed TR. Critical care nurses' knowledge about pulmonary embolism in respiratory care unit in Baghdad teaching hospitals. *Indian J Forensic Med Toxicol.* 2020;14(3):859–65.
11. M S Hassan S, S Hassan H. Effectiveness of Nursing Education Program on Nurses Practices Toward Arrhythmia in Kirkuk's Teaching Hospitals. *kufa J Nurs Sci.* 2013;3(1):220–30.
12. Abdul-hamza M, Aziz A. Effectiveness of an Educational Program upon Nurses' Knowledge toward the Continuous Positive Airway Pressure Machine in Neonatal Intensive Care Unit at AL-Hussein and the Pediatric Teaching

- Hospital. *Int J Sci Res.* 2017;6(10):10–6.
13. Haza'a A, Al-Qubati FA, Mohammed MA, Abdel-Aziz MA, Mehany MM. Effect of An Educational Program on Critical Care Nurses Performance regarding Emergency Care for Patients with Pulmonary Embolism. *Assiut Sci Nurs J.* 2020;8(20):31–44.
 14. Bhavani A, Sara B. A Study to Assess the Effectiveness of Structured Teaching Programme on Knowledge and Practice Regarding Prevention of Selected Complications of Immobilization among the Patients with Orthopedic Disorders in RMMCH, Chidambaram. *Int J Adv Nurs Manag.* 2017;5(1):51–4.
 15. Hammod HJ, Mohammed S. Effectiveness of an Educational Program on Nurses Knowledge Concerning Complications Prevention of Mechanical Ventilation at Intensive Care Unit in Al-Hussain Teaching Hospital at Nassiryah City. *Kufa J Nurs Sci.* 2016;6(2).

DOI: <https://doi.org/10.52845/CMRO/2024/7-4-28>

DEFINITIONS, ACRONYMS, ABBREVIATIONS

List of Abbreviations	
Symbols	Meaning
ANCOVA	Analysis of Covariance
VTE	Venous thromboembolism
DVT	Deep Vein Thrombosis
PE	pulmonary embolism
CCU	Coronary Care Unit
CDC	Centers for Disease Control and Prevention
ICU	Intensive Care Unit
K-S	Kolmogorov-Smirnov
M.S	Mean of Score
MCQ	Multiple Choice Questions
PGMS	Percentile Grand/or Global Mean of Score
PPSD	Percentile Pooled Standard Deviation
QI	Quality Improvement
RCU	Respiratory Care Unit
RS	Relative Sufficiency
SD	Standard Deviation
SDCv.	Socio-Demographical Characteristics Variables

SPSS	Statistical Package for the Social Sciences
e.g.,	Exempli Gratia (For Example)
etc.	Et Cetera (And So More)
C.S.	Comparison Significant
Sig. or S	Significance
G	Good
H.S	High Significant
M	Moderate
N. S	Non-Significant
P	Poor
df.	Degree Of Freedom
n	Number Of Samples
f.	Frequency
χ^2	Chi-square
\leq	Less than or equal
%	Percentage
<	Less than
\geq	More than or equal

APINDEX: Questionnaire

SECTION A: Socio-Demographic Data of The Participants in The Study.

Instruction: Please fill in the provided space or put an X in the area provided.

All participants are assured that your answers to this test will be treated with strictest confidentiality and you will not be identified in any way.

Thank you very much for your participation.

Socio-Demographic characteristics:

1. Age: /Years

Male	
------	--

Female	
--------	--

2. Gender:

3. Education level:	Secondary Nursing School	
	Nursing institute	
	Nursing college	
	Postgraduate	

4. Years of service:...../Years

5. Work place

6. Years of experience: /Years	CCU	
	Emergency Department	
	Surgical Units	

7. Received Training regarding Venous thrombosis:

Yes	
No	

PRE-TEST/POST-TEST

SECTION B: Participants knowledge regarding venous thrombosis.

Instruction: Please read each question carefully and circle the best answer. Please choose **only ONE** answer.

1. Is it considered one of the symptoms of deep vein thrombosis?

- A. Chest pain
- B. Shortness of breath
- C. Swelling and pain in the leg
- D. Severe headache

2. What are the signs and symptoms of PE that a patient may exhibit?

- A. Persistent cough
- B. Fever and extreme fatigue
- C. Eye pain
- D. Anorexia

3. What is the difference between venous thrombosis (DVT) and arterial thrombosis?

- A. DVT (occurs in the heart and arterial thrombosis in the venous vessels
- B. DVT occurs in the venous vessels and arterial blood clotting in the heart
- C. There is no difference between them
- D. DVT occurs in the deep venous vessels and arterial thrombosis in the arteries

4. What is the relationship of Restless Leg Syndrome to DVT?

- A. There is no relationship between them
- B. It could be an indicator of the presence of DVT
- C. Helps prevent DVT
- D. Causes DVT directly

5. What is the difference between deep phlebitis (DVT) and superficial phlebitis?

- A. Site of inflammation only
- B. Intensity and sharpness
- C. Quality of pain
- D. Time required to heal

6. What is the relationship between cancer and the risk of developing DVT?

- A. Cancer reduces the risk of DVT
- B. Cancer increases the risk of DVT
- C. There is no relationship between them
- D. Cancer only increases the risk of PE

7. The common diagnostic technique used to detect Pulmonary Embolism is?

- A. MRI
- B. X-ray
- C. CT scan
- D. Routine blood test

8. What medical tests can be used to determine the effectiveness of anticoagulant treatment?

- A. Urinalysis
- B. Blood test to check cholesterol level
- C. International Normalized Ratio (INR) analysis
- D. Lung MRI

9. What risk factors may increase the likelihood of blood clots forming?

- A. Long sleep
- B. Frequent exposure to the sun
- C. Obesity, smoking, and family history of clot formation
- D. High ambient temperatures

10. What age group is most susceptible to DVT?

- A. Less than 10 years
- B. Between 20 and 40 years old
- C. Older than 60 years
- D. Between 10 and 20 years

11. What are the main factors that may increase the risk of venous thrombosis?

- A. Intense physical activity
- B. Smoking
- C. Surgery and serious injuries
- D. Taking excess vitamins

12. The main factor contributing to the formation of venous thrombosis is?

- A. Low blood pressure
- B. Hyperperfusion
- C. Low cholesterol
- D. Blood clotting disorders

13. What is the treatment method for DVT?

- A. By anti-inflammatory medications
- B. Using anticoagulant medications

- C. Perform immediate surgery
- D. Massage and physical therapy

14. What is the first step in managing a PE emergency?

- A. Artificial cardiac massage
- B. Giving the patient oxygen
- C. Giving the patient a high dose of anti-inflammatory medication
- D. Perform immediate surgery

15. What is the main role of surgery in treating DVT?

- A. Direct removal of blood clots
- B. Vasodilatation
- C. Repair damaged blood vessels
- D. Improve heart function

16. What emergency treatment can be used in severe and serious PE?

- A. Oxygen therapy only
- B. Artificial cardiac massage
- C. Immediate administration of anticoagulant medications
- D. Surgery to remove a pulmonary clot

17. One of the common ways to prevent deep vein thrombosis is?

- A. Sitting for long periods without moving
- B. Exercise regularly
- C. Eat heavy meals before bed
- D. Refrain from drinking fluids

18. How can proper nutrition play a role in preventing DVT?

- A. Increase your intake of sugars
- B. Reduce protein intake
- C. Eat adequate amounts of vitamin K
- D. Maintain a proper balance of fluid and fiber intake

19. One of the factors that may contribute to the development of DVT complications is?

- A. High ambient temperatures
- B. Not following a high-fat diet
- C. Suddenly stopping taking anticoagulant medications
- D. Low blood cholesterol level

20. What is one of the possible complications of PE?

- A. High blood pressure
- B. coma
- C. Low blood sugar
- D. Lung collapse