

Effect of Sub-Chronic Consumption of Palm Oil-Tainted Feed on Hepatorenal Parameters in Wistar Rats

Abstract

The present study evaluated the effects of sub-chronic consumption of palm oil-tainted feed (POTF) on hepatorenal parameters in male Wistar rats. Twenty-five (25) male Wistar rats with an average weight of 150 ± 15 g were used for the study. They were randomly selected into 5 groups of 5 rats each. Group 1 served as control and accessed only normal feed and water *ad libitum*. Groups 2 and 3 were fed with 10% and 20% POTF and water *ad libitum* respectively. Groups 4 and 5 were fed with 10% and 20% POTF + 0.02mg/ml and 0.04mg/ml of simvastatin respectively, and water *ad libitum* respectively. After 60 days of exposure/consumption of the POTF; blood samples were harvested from the study animals via cardiac puncture after proper sedation. After laboratory and statistical analyses of the quantitative data obtained from the study, the serum level of a liver enzyme—alanine transaminase (ALT) had no significant ($P > 0.05$) change in all the test groups (groups 2, 3, 4, and 5) when compared to the control group. The level of aspartate transaminase (AST) was found to be significantly ($p < 0.05$) raised in test group 3 when compared to that of group 1 (control). The alkaline phosphatase (ALP) level was significantly ($p < 0.05$) reduced in test group 3 but markedly ($p > 0.05$) elevated in test group 4 when compared to group 2. The serum levels of total protein, albumin, and total bilirubin indicated no significant ($P > 0.05$) difference except for conjugated bilirubin which was significantly ($p < 0.05$) reduced in test group 2. In conclusion, except for higher doses, that may elicit some possible unbeneficial effects, mild sub-chronic consumption (10%) of POTF in the study animals did not result in any adverse alterations in both liver enzymes and some renal indices in the study models.

Keywords: *sub-chronic consumption; palm oil tainted feed; hepatorenal parameters; moderate use of safe and natural palm oil.*

Introduction

The pattern of diet across different populations has been linked to the trend of disease prevalence [1, 2]. More so, a healthy dietary pattern is related to reducing the risk of chronic metabolic disease; thus, there is a great need for a better understanding of such benefits [3]. An effective approach to achieving this target is putting up efficient checks on the dietary patterns and their relationship with multiple chronic metabolic disorders; as this can be of great benefit to the populace [4].

Considering *Elaeis guineensis*, a species of palm commonly called oil palm (also known as African oil palm or macaw-fat), it is an age-long trendy food for many West African populations. Again, it is one of the most popular and easy to get, cooking oils used for many delicacies in virtually every home in our population [5, 6]. The plant has several local names like “epo pupa”

in Yoruba, the Hausa of Northern Nigeria called it “*jar mai or man ja*” and the Igbo called it “*Nkwu or nmanunri*” etc.

The plant is known to be rich in beta-carotene and contains saturated (and even unsaturated fats) and some other nutrients like vitamin E and beta-carotene [6, 7].

Even though palm oil and palm kernel oil, are recorded to contain high levels of saturated fatty acids—roughly 50% and 80%, respectively and a source of dietary fat; consuming it as part of a healthy diet does not increase the risk of coronary artery disease [8].

There have been inconsistent reports about the effects of diets rich in palm oil on the increase in risk of cardiovascular diseases (CVDs)/coronary heart diseases (CHDs) in humans [9]. For instance, saturated fats, reported to be contained in palm oil, are known to possess the potential to raise total cholesterol, which includes low-density lipoprotein cholesterol (LDL-C), which is more dangerous [10, 11]. Elevations in LDL-C level are also associated with plaque formation in arteries which is largely contributory to CVD/CHD conditions [12, 13].

It therefore calls for the need for further evaluations to unravel the actual relationship between continuous and increasing use of palm oil and CVDs. The current study therefore investigated the possible effect of sub-chronic consumption of palm oil on hepatorenal parameters in Wistar Rats.

Materials and methods

About (25) male Wistar rats with a weight of 150 ± 15 grams were procured from the Animal House of the Department of Pharmacology, Faculty of Basic Medical Sciences, University of Port Harcourt. The animals were separated into 5 groups of 5 rats and then housed in well-aerated standard top wire-gauzed plastic cages with sawdust beddings. They were maintained at room temperature under the 12-hour light/dark cycle. The animals were allowed two weeks of acclimatization to their new handling and housing and they accessed their differently formulated feeds and water both in this period and through the treatment interval.

Experimental Protocol

Group 1 served as the control group and the animals were allowed access to 100 grams of normal rat feed and water.

Group 2 were fed with 100 grams of rat feed tainted with 10% palm oil and water.

Group 3 was fed with 100 grams of rat feed tainted with 20% palm oil and water.

Group 4 was fed with 100 grams of rat feed tainted with 10% palm oil, water, and 0.02mg/ml of simvastatin.

Group 5 was fed with 100 grams of rat feed tainted with 20% palm oil, water, and 0.04mg/ml of simvastatin.

It is noteworthy that, 10% and 20% gram weight of the palm oil were mixed with 90% and 80% gram weights of the normal rat feed respectively to constitute the 10% and 20% of palm oil tainted feed (POTF).

Harvest of Samples from the study Models

After sixty (60) days of exposure to the different levels of POTF, blood samples were obtained from the study animals through cardiac puncture after proper sedation using 80% trichloromethane (chloroform). The sample was immediately moved into properly labeled lithium heparin bottles and presented for the designated laboratory screening.

Determination of liver enzymes and other biochemical parameters levels

The Randox assay guides and spectrophotometric detection technique (i.e. automated clinical chemistry analyzer) were used to determine the serum concentrations of ALT, AST, ALP, bilirubin, total protein, and albumin [14, 15, 16].

Method of Data Analysis

The numerical data from the present study were subjected to statistical analyses using analyses of variance and Post Hoc tools of the IBM Statistical Product and Service Solutions (SPSS) 21.0V software. The data were presented as Mean \pm Standard error of the mean. Variances between means were determined using Analysis of variance (ANOVA) and post-test using LSD multiple comparison tool and Dunnett at 95% probability.

Ethical Approval

Ethical authorization was sought and obtained from the Ethical Committee of the Department of Human Physiology, Faculty of Basic Medical Sciences, University of Port Harcourt. The study models were handled in line with the recommendations of the US National Institute of Health (NIH) guidelines for the care and use of laboratory animals in experimental research [17].

Result

The data in Table 1 displays the effect of sub-chronic consumption of palm oil on some hepatorenal parameters in male Wistar rats.

The variation in the level of alanine transaminase (ALT) did not show any significant ($p>0.05$) changes when all test groups were compared to the control group and among themselves. For the effect of the sub-chronic consumption of palm oil on aspartate transaminase (AST), only the animals that consumed 20% (group 3) indicated a raised level when compared to that of the animals in the control group. Considering the changes in alkaline phosphatase (ALP) levels following the sub-chronic consumption of palm oil, there appears to be a general reduction in ALP levels when compared to that of control. The aforementioned reductions were seen to be significant in group 2 (treated with 10% of palm oil). On the other hand, the 10% palm oil + simvastatin treated animals (group 4) revealed a significantly elevated ALP level when compared to that of 10% of palm oil treated animals (group 2).

Table 1: Effect of sub-chronic consumption of palm oil tainted feed (POTF) on hepatorenal parameters in male Wistar rats

Groups and Treatment	ALT ($i\mu/L$)	AST ($i\mu/L$)	ALP ($i\mu/L$)	Total Protein (g/L)	Albumin (g/L)	Total Bilirubin $\mu mo/L$	Conjugated Bilirubin $\mu mo/L$
Group 1: Control	5.60 ± 0.98	7.60 ± 0.60	32.20 ± 2.52	64.20 ± 13.68	41.40 ± 2.77	18.84 ± 12.54	4.90 ± 0.65
Group 2: 10% PALM OIL SC	4.80 ± 0.80	11.20 ± 2.24	21.60 ± 3.35 ^a	66.60 ± 7.01	44.00 ± 4.06	5.34 ± 0.98	2.38 ± 0.60 ^a
Group 3: 20% PALM OIL SC	4.80 ± 0.80	15.0 ± 2.75 ^a	29.40 ± 3.93	72.20 ± 5.60	42.60 ± 0.87	5.40 ± 0.90	4.28 ± 0.85
Group 4: 10% PALM OIL SC + SIMV	6.40 ± 0.98	10.80 ± 1.65	32.20 ± 1.71 ^b	43.20 ± 11.12	42.60 ± 1.43	5.54 ± 1.24	3.50 ± 0.76
Group 5: 20% PALM OIL SC + SIMV	5.60 ± 0.98	12.00 ± 2.96	23.80 ± 4.21	38.00 ± 13.80	42.60 ± 2.38	6.16 ± 0.99	4.58 ± 0.72 ^b

Values represent mean ± SEM, n=5; ^a Significant at p<0.05 when compared to Group 1; ^b Significant at p<0.05 when compared to group 2; ^c Significant at p<0.05

The result on total protein changes indicated that the sub-chronic consumption of palm oil in the study animals did not elicit any significant ($p>0.05$) effect on them. Similarly, the changes in albumin and total bilirubin levels as recorded in the present study revealed no—significant ($p>0.05$) change for all test groups when compared to group 1 and among themselves. Meanwhile, for the conjugated bilirubin changes, there were significant ($p<0.05$) reductions in group 2 (10% palm oil-treated animals) and elevation in group 5 (20% palm oil + simvastatin-treated animals).

Discussion

While some schools of thought think that the continuous consumption of palm oil may not possess any adverse effects on the hepato-renal system, heart and blood vessels, or ischemia injuries and plaque buildup [18, 19], others hold that palm oil consumption may relatively increase LDL cholesterol, and other adverse biological alterations [20, 21]. This controversy prompted the focus of the present study, which has made relevant findings that are discussed in the following paragraphs.

A major outcome of the present study indicated a stable or normal level of ALT but a marked increase in AST level of the 20% POTF fed group. Of course, the evaluation of ALT level is considered a very reliable enzyme marker with high specificity in diagnosing liver toxicity [22]. This finding, thus, indicates that the consumption of the POTF in the study models may not have adversely affected the normal function of the hepatocytes. The raised level of AST in the 20% POTF group could be a pointer consumption amount of POTF that may result in damage to extra-hepatic tissues or hemolysis; this outcome validates the earlier submissions of Gowda et al., [23] and Kalas et al., [24], as AST level can be marked influenced by extra-hepatic toxicities; especially, seeing here that the ALT level was unaffected.

Another important serum analyte is alkaline phosphatase (ALP) which has a correlation with the presence of bone, liver, and other diseases when it is elevated in the serum [25]. The depression in ALP level, which is rare, could indicate a lack of zinc, malnutrition, pernicious anemia, thyroid disease, etc., and can affect bones and teeth [26, 27]. Thus the marked depression in ALP level in the POTF group of animals in the present study is a pointer to a possible adverse effect of increasing POTF consumption. Therefore, while further in-depth study may be required to establish this interesting finding of the present study on the possible ALP depressive effects of the consumption of POTF in the study models, there is a need for caution in increasing the use of palm oil as indicated in the study models.

Plasma proteins are known to be majorly synthesized in the liver [28, 29, 30], thus, the marginal effects of consumption of POTFs on the study models' total protein and albumin levels, buttresses the earlier result on ALT in the present study. Therefore the moderate consumption of POTFs in the study animals may not have elicited any severe effects on the hepatorenal system.

The present study also found generally reduced levels of conjugated bilirubin in virtually all rats that fed on the POTFs; this was relatively significant when group 2 (that fed on 10% Palm oil tainted group) values were compared to that of the control group.

Due to bilirubin's hydrophilicity and large molecular size, it is easily reabsorbed via the intestinal walls, but the process of its conjugation prevents the intestinal mucosa from passively reabsorbing it. The conjugation, therefore, aids in removing the potentially harmful metabolic waste products [31, 32]. Now, because the heme oxygenase in Kupffer cells is responsible for the synthesis of bilirubin [33]; it is suggestive to state that the consumption of increasing doses of POTF may have altered the function of the Kupffer cells. This finding also indicates that further study to evaluate the possible difference in the effects of palm consumption on hepatocytes and Kupffer cells of the liver may be necessary.

Conclusion

The present study found that except for higher doses, that may elicit some possible unbeneficial effects, mild sub-chronic consumption of POTF in the study animals may not result in any adverse alterations in both liver enzymes and some renal indices in the study models. It is therefore suggestive to state that, continuous use of palm oil should consider moderate use of safe and natural palm oil to prevent any negative health effects.

References

1. Popkin BM. Global nutrition dynamics: the world is shifting rapidly toward a diet linked with noncommunicable diseases. *The American journal of clinical nutrition*. 2006 Aug 1;84(2):289-98.
2. Jia Q, Xia Y, Zhang Q, Wu H, Du H, Liu L, Wang C, Shi H, Guo X, Liu X, Li C. Dietary patterns are associated with prevalence of fatty liver disease in adults. *European journal of clinical nutrition*. 2015 Aug;69(8):914-21.
3. Li T, Guan L, Wang X, Li X, Zhou C, Wang X, Liang W, Xiao R, Xi Y. Relationship between dietary patterns and chronic diseases in rural population: management plays an important role in the link. *Frontiers in Nutrition*. 2022 Apr 13;9:866400.
4. Panagiotakos DB, Pitsavos C, Skoumas Y, Stefanadis C. The association between food patterns and the metabolic syndrome using principal components analysis: The ATTICA Study. *Journal of the American Dietetic Association*. 2007 Jun 1;107(6):979-87.
5. Achaw OW, Danso-Boateng E. Manufacture of crude palm oil and refined palm oil. *In* *Chemical and Process Industries: With Examples of Industries in Ghana* 2021 Aug 10 (pp. 195-211). Cham: Springer International Publishing.
6. Raji RO, Inengite AK, Godwin J, Ajibesin KK. Assessment of proximate and physicochemical properties of crude palm oil from south-west and South-south Nigeria. *African Journal of Pure and Applied Chemistry*. 2022 Aug 31;16(2):40-56.

7. Edo GI, Makinde MG, Nwosu LC, Ozgor E, Akhayere E. Physicochemical and pharmacological properties of palm oil: an approach for quality, safety, and nutrition evaluation of palm oil. *Food Analytical Methods*. 2022 Aug;15(8):2290-305.
8. Imoisi OB, Ilori GE, Agho I, Ekhatior JO. Palm oil, its nutritional and health implications. *Journal of Applied Sciences and Environmental Management*. 2015;19(1):127-33.
9. Mancini A, Imperlini E, Nigro E, Montagnese C, Daniele A, Orrù S, Buono P. Biological and nutritional properties of palm oil and palmitic acid: effects on health. *Molecules*. 2015 Sep 18;20(9):17339-61.
10. Siri-Tarino PW, Sun Q, Hu FB, Krauss RM. Saturated fatty acids and risk of coronary heart disease: modulation by replacement nutrients. *Current atherosclerosis reports*. 2010 Nov;12:384-90.
11. DiNicolantonio JJ, Lucan SC, O'Keefe JH. The evidence for saturated fat and sugar related to coronary heart disease. *Progress in cardiovascular diseases*. 2016 Mar 1;58(5):464-72.
12. Reiner Ž. Hypertriglyceridaemia and risk of coronary artery disease. *Nature Reviews Cardiology*. 2017 Jul;14(7):401-11.
13. Mortensen MB, Caínzos-Achirica M, Steffensen FH, Bøtker HE, Jensen JM, Sand NP, Maeng M, Bruun JM, Blaha MJ, Sørensen HT, Pareek M. Association of coronary plaque with low-density lipoprotein cholesterol levels and rates of cardiovascular disease events among symptomatic
14. Hsueh CJ, Wang JH, Dai L, Liu CC. Determination of alanine aminotransferase with an electrochemical nano Ir-C biosensor for the screening of liver diseases. *Biosensors*. 2011 Jul 12;1(3):107-17.
15. Zhang P, Wang CY, Li YX, Pan Y, Niu JQ, He SM. Determination of the upper cut-off values of serum alanine aminotransferase and aspartate aminotransferase in Chinese. *World Journal of Gastroenterology: WJG*. 2015 Feb 2;21(8):2419.
16. Ezejindu DN, Chinweife KC, Ihentuge CJ. The effects of moringa extract on liver enzymes of carbon tetrachloride-induced hepatotoxicity in adult wister rats. *Int J Eng Sci*. 2013;2(7):54-9.
17. National Research Council, (2011). Guidelines of the National Institute of Health (NIH) for care and use of laboratory/research animals. [Retrieved in November, 2019 from: <https://www.ncbi.nlm.nih.gov/books/NBK54050/>].adults. *JAMA network open*. 2022 Feb 1;5(2):e2148139.
18. Odia OJ, Ofori S, Maduka O. Palm oil and the heart: a review. *World journal of cardiology*. 2015 Mar 3;7(3):144.

19. Ibrahim, N.I., Fairus, S. and Naina Mohamed, I.,. The effects and potential mechanism of oil palm phenolics in cardiovascular health: a review on current evidence. 2020; *Nutrients*, 12(7), p.2055.
20. Sun Y, Neelakantan N, Wu Y, Lote-Oke R, Pan A, van Dam RM. Palm oil consumption increases LDL cholesterol compared with vegetable oils low in saturated fat in a meta-analysis of clinical trials. *The Journal of nutrition*. 2015 Jul 1;145(7):1549-58.
21. Abdulwaliyu I, Okoduwa SI, Sangodare R, Arekemase SO, Muhammad A. Review of Studies on Palm-Oil Consumption in Relation to Risk of Cardiovascular Diseases. *Journal of Nutrition and Food Security*. 2022.
22. Ozer J, Ratner M, Shaw M, Bailey W, Schomaker S. The current state of serum biomarkers of hepatotoxicity. *Toxicology*. 2008 Mar 20;245(3):194-205.
23. Gowda S, Desai PB, Hull VV, Avinash AK, Vernekar SN, Kulkarni SS. A review on laboratory liver function tests. *The Pan african medical journal*. 2009;3.
24. Kalas MA, Chavez L, Leon M, Taweeseedt PT, Surani S. Abnormal liver enzymes: A review for clinicians. *World journal of hepatology*. 2021 Nov 11;13(11):1688.
25. Sharma U, Pal D, Prasad R. Alkaline phosphatase: an overview. *Indian J Clin Biochem*. 2014 Jul;29(3):269-78. doi: 10.1007/s12291-013-0408-y. Epub 2013 Nov 26. PMID: 24966474; PMCID: PMC4062654.
26. Verma P. Alkaline Phosphatase: A Review Article. *International Journal of Biochemistry and Biomolecules*. 2022 Jul 22;8(1):23-8p.
27. Riancho JA. Diagnostic approach to patients with low serum alkaline phosphatase. *Calcified Tissue International*. 2023 Mar;112(3):289-96.
28. Miller LL, Bly CG, Watson ML, Bale WF. The dominant role of the liver in plasma protein synthesis: A direct study of the isolated perfused rat liver with the aid of lysine- ϵ -C14. *The Journal of experimental medicine*. 1951 Oct 10;94(5):431.
29. Reeve EB, Pearson JR, Martz DC. Plasma protein synthesis in the liver: method for measurement of albumin formation in vivo. *Science*. 1963 Mar 8;139(3558):914-6.
30. De Feo P, Lucidi P. Liver protein synthesis in physiology and in disease states. *Current Opinion in Clinical Nutrition & Metabolic Care*. 2002 Jan 1;5(1):47-50.
31. Wang X, Chowdhury JR, Chowdhury NR. Bilirubin metabolism: applied physiology. *Current Paediatrics*. 2006 Feb 1;16(1):70-4.

32. Sticova E, Jirsa M. New insights in bilirubin metabolism and their clinical implications. World Journal of Gastroenterology: WJG. 2013 Oct 10;19(38):6398.
33. Naito M, Hasegawa G, Ebe Y, Yamamoto T. Differentiation and function of Kupffer cells. Medical Electron Microscopy. 2004 Mar;37:16-28.

UNDER PEER REVIEW