

1  
2  
3  
4  
5  
6  
7  
8  
9  
10

# Impact of Heavy Metal Contamination on Soil and Crop Ecosystem with Advanced Techniques to Mitigate Them

---

## ABSTRACT

Heavy metal pollution in soils has become widespread globally due to both natural processes and human activities. Geologic and anthropogenic activities, such as mining, industrial processes, and agricultural practices, significantly contribute to heavy metal accumulation in soil, posing threats to both plant and animal health. These metals endure in the environment and build up in organisms because they resist breakdown or metabolism. They enter the food chain via primary producers and move up as higher trophic levels consume them. This contamination adversely affects the overall human health, plant growth, performance, and yield and also impacts the soil properties. To minimize this effect of heavy metal toxicity, bioremediation emerges as a highly effective method for treating such polluted soils, often conducted in situ, making it suitable for crop establishment or reestablishment. Bioremediation encompasses various techniques like bioventing, biostimulation, air sparging, natural attenuation, phytoremediation, vermistabilization, rhizofiltration, biosorption etc. Both microorganisms and plants play crucial roles in bioremediation, with the combined approach ensuring more efficient cleanup. However, the success of this method depends significantly on the specific species involved. Traditional physical and chemical remediation methods are often costly and fail to restore soil suitability for plant growth. In contrast, bioremediation offers a more environmentally friendly and economically viable solution, leveraging natural processes to encourage plant growth on contaminated soils.

*Keywords: Bioremediation, biostimulation, environment, heavy metals, phytoremediation,*

## 1. INTRODUCTION

“Heavy metals are metallic elements characterized by high atomic weights and densities and often accumulate in the environment, posing risks to ecosystems and human health. A considerable number of heavy metals pose toxicity, contributing to environmental pollution. Due to their non-biodegradable and water-soluble properties, they easily permeate plants and human bodies, leading to negative consequences”[1]. “The term “heavy metal” denotes a metal or metalloid possessing an atomic density at least five times higher than that of water”[2,3]. “Common heavy metals present in the environment comprise cadmium (Cd), lead (Pb), nickel (Ni), silver (Ag), zinc (Zn), cobalt (Co), chromium (Cr), iron (Fe), arsenic (As), and mercury (Hg). The high concentration of heavy metals in water, air, and soil poses a threat to all living organisms”[4]. “While heavy metals naturally occur in soil, human activities, both geological and anthropogenic, escalate their concentrations to levels detrimental to plant and animal health. These activities comprise mining and metal smelting, fossil fuel combustion, agricultural use of fertilizers and pesticides, industrial production of batteries and metal goods, sewage sludge disposal, and municipal waste management. Recorded instances show that plants growing on soils contaminated with heavy metals experience growth reduction due to alterations in their physiological and biochemical processes” [5,6,7]. “Heavy metals predominantly exist in dispersed form within rock formations. The human impact of heavy metal presence in the biosphere has surged due to industrialization and urbanization. These metals are most

abundant in soil and aquatic ecosystems, with a smaller presence in the atmosphere as particles or vapors. The toxicity of heavy metals to plants depends on various factors such as plant species, the particular metal involved, its concentration, chemical form, and soil properties including composition and pH. Interestingly, several heavy metals are essential for plant growth, and similarly, many serve as essential trace nutrients for animals and the human body”[8]. “Similarly, numerous heavy metals play essential roles for plants by serving as cofactors, triggering enzyme reactions, exhibiting ductility and conductivity, and ensuring cation stability”[9]. “Heavy metals enter the human body through different routes, including contaminated food, water, skin contact, and inhalation. Those absorbed through the intestine are typically water-soluble and travel to various organs via the circulatory system. Even at low concentrations, heavy metals impact the respiratory tract and various cell types such as endothelial and epithelial cells”[10].

These heavy metals persist in the environment and accumulate in living organisms, as they are not easily broken down or metabolized. They enter the ecological food chain starting from primary producers and then move up through consumption at higher trophic levels. Plants, being stationary, primarily interact with heavy metal ions through their roots, while in aquatic systems, the entire plant body is exposed to these ions. Additionally, heavy metals can directly adhere to leaves through particles deposited on foliar surfaces. Human exposure to heavy metals occurs mainly through inhalation or ingestion, with ingestion being the primary route for the general population. However, urbanization, traffic, industrial and agricultural activities, waste incineration, and mining significantly contribute to heavy metal inhalation exposure. Metals like Cadmium (Cd), Nickel (Ni), Arsenic (As), and Chromium (Cr) present various hazards to human health and are potent carcinogens. Cadmium intake can lead to Itai-Itai disease, while mercury intake can result in Minamata disease. Arsenic and other heavy metals can cause poisoning through contamination of drinking water.

“Numerous methods are available for remediating metal-polluted soils, encompassing a spectrum from physical and chemical approaches to biological ones. However, many physical and chemical methods, including encapsulation, solidification, stabilization, electrokinetics, vitrification, vapor extraction, soil washing, and flushing, tend to be costly and may not render the soil conducive to plant growth”[11]. The biological approach, known as bioremediation, promotes the growth or re-growth of plants in polluted soils. It stands out for its environmental friendliness, as it relies on natural processes. Additionally, bioremediation proves to be a cost-effective technique compared to other remediation methods. This paper explores the characteristics of soils contaminated with heavy metals, evaluates plant growth and productivity in such soils, and emphasizes the utilization of biological methods for remediating heavy metal pollution.

## 2. SOURCES OF HEAVY METAL POLLUTION

### 2.1 NATURAL SOURCES

“Heavy metals originating from rock materials constitute the "lithogenic" aspect. The concentration and composition of heavy metals formed in soil are primarily determined by the type of parent rock”[12]. “The main heavy metal pollutants contributed by parent rocks comprise Cobalt (Co), Chromium (Cr), Iron (Fe), Manganese (Mn), Nickel (Ni), and Zinc (Zn). The weathering process of igneous rocks, such as Augite, Olivine, and Hornblende, results in substantial amounts of heavy elements, whereas sedimentary rocks contribute only a minor portion”[13].

“In addition to rocks, natural sources of heavy metals include volcanoes, wind-blown dust and storms, wildfires, sea sprays, and aerosols (especially in coastal regions)”[14]. “Geothermal sources, along with volcanic eruptions, have also contributed significantly to atmospheric pollutants and toxic waste”[15].

### 2.2 ANTHROPOGENIC SOURCES

2.2.1 **Agricultural sources:** Inorganic and organic fertilizers, manure, lime, pesticides, and other agricultural inputs contain varying levels of Chromium (Cr), Cadmium (Cd), Nickel (Ni), Zinc (Zn), Lead (Pb), and other heavy metals. Similarly, many widely used chemical pesticides such as Bordeaux mixture and lead arsenate contain Copper (Cu), Mercury (Hg), Manganese (Mn), Lead (Pb), or Zinc (Zn). Additionally, the utilization of municipal and industrial wastewater for irrigation serves as a significant source of heavy metal contamination in soil.

2.2.2 **Industrial Sources:** “Industrial activities such as mining, smelting, and metal processing primarily produce Chromium (Cr) and Nickel (Ni), while Vanadium (V), Titanium (Ti), and Manganese (Mn) are predominantly derived from oil and coal-related operations”[16]. “Coal mining releases significant amounts of Arsenic (As), Cadmium (Cd), and Iron (Fe), while gold mining elevates Mercury (Hg) levels in the environment”[17]. “These heavy metals are typically emitted in particulate and vapor forms, which, upon interaction with atmospheric moisture, form aerosols. These aerosols can be dispersed by wind (dry deposition) or deposited through rainfall (wet deposition), leading to soil and water contamination”[13]. “Moreover, combustion processes involving coal,

92 petroleum, and nuclear power stations release heavy metals such as Selenium (Se), Cadmium (Cd), Boron (B),  
93 Copper (Cu), Cesium (Cs), Zinc (Zn), and Nickel (Ni) into the atmosphere”[18]. “Additionally, activities like plastics  
94 processing, microelectronics manufacturing, wood preservation, textiles production, and paper processing also  
95 contribute to heavy metal pollution in the environment”[3].

96 2.2.3 **Domestic Effluents:**“The primary source of heavy metal pollution in water bodies is domestic effluents. These  
97 effluents encompass untreated wastewater, as well as substances that have been processed through biological  
98 treatment plants and discharged through sewage outfalls. Additionally, many commonly used enzymatic  
99 detergents contain small amounts of elements such as Iron (Fe), Chromium (Cr), Manganese (Mn), Zinc (Zn),  
100 Cobalt (Co), Strontium (Sr), and Boron (B), thereby adding to the burden of heavy metal pollution”[19].

### 101 3. EFFECT OF HEAVY METAL CONTAMINATION IN SOIL

102  
103 Heavy metals possess metallic characteristics like ductility, malleability, conductivity, cation stability, and ligand specificity.  
104 They are characterized by their high density and atomic weight, typically having an atomic number greater than 20. While  
105 some heavy metals like Cobalt (Co), Copper (Cu), Iron (Fe), Manganese (Mn), Molybdenum (Mo), Nickel (Ni), Vanadium  
106 (V), and Zinc (Zn) are essential in small quantities for organisms, excessive amounts of these elements can be  
107 detrimental. On the other hand, heavy metals such as Lead (Pb), Cadmium (Cd), Mercury (Hg), and Arsenic (As)  
108 (considered a metalloid but often grouped with heavy metals) have no beneficial effects on organisms and are considered  
109 significant threats due to their harmful effects on both plants and animals. Metals can be found in soil either independently  
110 or in association with other soil constituents. These constituents encompass exchangeable ions adhered to the surfaces  
111 of inorganic solids, nonexchangeable ions, and insoluble inorganic metal compounds like carbonates and phosphates.  
112 Additionally, metals may exist as soluble compounds or free ions in the soil solution, as complexes with organic materials,  
113 or attached to silicate minerals [11]. Metals attached to silicate minerals constitute the inherent soil metal levels and  
114 typically do not pose contamination or pollution concerns, unlike metals found independently or in high concentrations  
115 within the other four components [20].  
116  
117

118 Various soil properties have distinct impacts on the availability of metals. Soil pH emerges as a crucial determinant of  
119 metal availability [21]. For example, the presence of Cadmium (Cd) and Zinc (Zn) for *Thlaspi caerulescens* roots  
120 diminishes with higher soil pH levels. Moreover, organic matter and hydrous ferric oxide have demonstrated the ability to  
121 reduce heavy metal availability by immobilizing these metals [22]. Additionally, notable positive associations have been  
122 observed between heavy metals and certain soil physical attributes such as moisture content and water holding capacity  
123 [23].

124 Metal availability in soil is influenced by several factors, including the density and charge type of soil colloids, the extent of  
125 complexation with ligands, and the relative surface area of the soil [11,24]. Soil colloids, with their extensive interface and  
126 specific surface areas, play a crucial role in regulating heavy metal concentrations in natural soils. Additionally, in polluted  
127 soils, soluble metal concentrations may be decreased by soil particles possessing high specific surface areas, although  
128 this effect may vary depending on the metal [11]. For example, the addition of amendments containing hydroxides with  
129 high reactive surface areas reduced the solubility of Arsenic (As), Cadmium (Cd), Copper (Cu), Molybdenum (Mo), and  
130 Lead (Pb), while the solubility of Nickel (Ni) and Zinc (Zn) remained unchanged [25]. Furthermore, soil aeration, microbial  
131 activity, and mineral composition have been demonstrated to impact heavy metal availability in soils [26]  
132

133 On the other hand, heavy metals can alter soil properties, particularly biological aspects [27]. Monitoring changes in soil  
134 microbiological and biochemical properties post-contamination serves as a sensitive method for assessing soil pollution  
135 intensity, providing faster results compared to monitoring physical and chemical properties [28]. Heavy metals influence  
136 the abundance, diversity, and functions of soil microorganisms, with their toxicity dependent on various factors including  
137 soil temperature, pH, clay minerals, organic matter, inorganic ions, and chemical forms of the metal.

138 Studies examining the impact of heavy metals on soil biological properties yield varying results, with some showing  
139 negative effects while others find no significant relationship. Discrepancies likely stem from differences in experimental  
140 setups, including the use of artificially contaminated soils in labs versus naturally polluted soils in the field. Regardless of  
141 soil origin, further detailed research is needed to fully understand heavy metals' effects on soil ecosystems. It's advisable  
142 to employ multiple methods, such as assessing microbial biomass and enzymatic activities, to gain a comprehensive  
143 understanding. Additionally, heavy metals can influence each other's availability in soil, affecting plant uptake. Interactions  
144 like antagonism and synergy exist among heavy metals, as observed in various studies. For instance, the inhibitory effect  
145 of manganese on carbon mineralization can be countered by cadmium. Similarly, competition between copper and zinc,  
146 as well as nickel and cadmium, has been noted in plant membrane carriers. However, the relationship between metals  
147 can be complex, as copper has been found to enhance zinc toxicity in certain plants, highlighting the need for further

investigation in this field. Furthermore, different species of the same metal can interact with each other. For instance, the presence of arsenite strongly inhibited the uptake of arsenate by rice plants growing in polluted soil[29].

#### 4. EFFECT OF HEAVY METAL CONTAMINATION IN CROPS

“The heavy metals accessible for plant uptake are either those existing as soluble components in the soil solution or those readily solubilized by root exudates. While plants require specific heavy metals for their growth and maintenance, excessive amounts of these metals can pose toxicity risks. Plants' ability to accumulate essential metals also facilitates their uptake of other nonessential metals. Metals found in elevated levels in the soil influence seed germination, plant growth, production, as well as physiological, biochemical, and genetic aspects of the plant system”[30].

Elevated concentrations of heavy metals can cause various direct toxic effects, such as inhibiting cytoplasmic enzymes and damaging cell structures due to oxidative stress. An example of an indirect toxic effect is the displacement of essential nutrients at cation exchange sites in plants. Furthermore, heavy metals' detrimental impact on the growth and activities of soil microorganisms can indirectly affect plant growth. For instance, a decrease in the population of beneficial soil microorganisms due to high metal concentrations may lead to reduced organic matter decomposition, resulting in soil nutrient depletion. Additionally, heavy metal interference with soil microorganisms' activities can impede enzyme functions crucial for plant metabolism. These toxic effects, whether direct or indirect, ultimately contribute to a decline in plant growth and, in severe cases, may lead to plant death. Some of the common examples of heavy metal contamination on crops are as follows:

##### 1. Arsenic (As) toxicity:

- **Rice:** Reduced seed germination, decreased seedling height, reduced leaf area, and lower dry matter production.
- **Tomato:** Decreased fruit yield and lower leaf fresh weight.
- **Canola:** Stunted growth, chlorosis (yellowing of leaves), and wilting.

##### 2. Cadmium (Cd) toxicity:

- **Wheat:** Reduced seed germination, decreased plant nutrient content, and shorter shoot and root length.
- **Garlic:** Reduced shoot growth and accumulation of Cd in the plant.
- **Maize:** Reduced shoot growth and inhibition of root growth.

##### 3. Cobalt (Co) toxicity:

- **Tomato:** Reduction in plant nutrient content.
- **Mung bean:** Reduced antioxidant enzyme activities, decreased plant sugar, starch, amino acids, and protein content.
- **Radish:** Reduced shoot length, root length, and total leaf area; decreased chlorophyll content; reduced plant nutrient content, antioxidant enzyme activity, plant sugar, amino acid, and protein content.

##### 4. Chromium (Cr) toxicity:

- **Wheat:** Reduced shoot and root growth.
- **Tomato:** Decreased plant nutrient acquisition.
- **Onion:** Inhibition of germination process and reduction of plant biomass.

##### 5. Copper (Cu) toxicity:

- **Bean:** Accumulation of Cu in plant roots, root malformation, and reduction.
- **Black bindweed:** Plant mortality, reduced biomass, and decreased seed production.
- **Rhodes grass:** Reduced root growth.

##### 6. Mercury (Hg) toxicity:

- **Rice:** Decreased plant height, reduced tiller and panicle formation, yield reduction, and bioaccumulation in shoot and root of seedlings.
- **Tomato:** Reduced germination percentage, decreased plant height, reduced flowering and fruit weight, and chlorosis.

##### 7. Manganese (Mn) toxicity:

- **Broad bean:** Mn accumulation in shoot and root, reduced shoot and root length, and chlorosis.
- **Spearmint:** Decreased chlorophyll a and carotenoid content, and accumulation of Mn in plant roots.
- **Pea:** Reduced chlorophylls a and b content, lower relative growth rate, reduced photosynthetic O<sub>2</sub> evolution activity, and decreased photosystem II activity.
- **Tomato:** Slower plant growth and decreased chlorophyll concentration.

##### 8. Nickel (Ni) toxicity:

- **Pigeon pea:** Decreased chlorophyll content, reduced stomatal conductance, and decreased enzyme activity affecting the Calvin cycle and CO<sub>2</sub> fixation.
- **Rye grass:** Reduced plant nutrient acquisition, decreased shoot yield, and chlorosis.

- **Wheat:** Reduced plant nutrient acquisition.
  - **Rice:** Inhibition of root growth.
9. **Lead (Pb) toxicity:**
- **Maize:** Reduced germination percentage, suppressed growth, reduced plant biomass, and decreased plant protein content.
  - **Portia tree:** Reduced number of leaves and leaf area, decreased plant height, and reduced plant biomass.
  - **Oat:** Inhibition of enzyme activity affecting CO<sub>2</sub> fixation.
10. **Zinc (Zn) toxicity:**
- **Cluster bean:** Reduced germination percentage, decreased plant height and biomass, and reduced chlorophyll, carotenoid, sugar, starch, and amino acid content.
  - **Pea:** Reduced chlorophyll content, altered chloroplast structure, decreased photosystem II activity, and reduced plant growth.
  - **Rye grass:** Accumulation of Zn in plant leaves, growth reduction, decreased plant nutrient content, and reduced efficiency of photosynthetic energy conversion.

**Table 1. Range of Toxic heavy metals in plants and crops [31]**

Heavy metals	Range in soils (ppm d. wt.)	Range in crops (ppm d. wt.)
Cadmium	0.01–0.7	0.2–0.8
Nickel	10–1000	1.0
Arsenic	0.1–40	0.01–5
Mercury	0.01–0.2	0.005–0.02
Lead	2–200	0.1–10

**Table 2. Regulatory limits of some toxic heavy metals on humans [32]**

Heavy metals	Max. acceptable conc. (WHO)	Max. acceptable conc. in drinking water (EPA)
Cadmium	0.003 ppm	0.005 ppm
Nickel	–	2–4.3 ppb
Arsenic	0.01 ppm	0.01 ppm
Mercury	0.001 ppm	2 ppb
Lead	0.01 ppm	50 ppb

## 5. APPROACHES TO MITIGATE HEAVY METAL TOXICITY

The biological approach, known as bioremediation, promotes the establishment or reestablishment of plants in polluted soils. It's considered environmentally friendly as it relies on natural processes for its implementation [33]. Bioremediation presents an opportunity to neutralize or eliminate various pollutants through natural biological processes. It relies on cost-

effective, simple techniques that are generally well-received by the public and can often be conducted on-site. However, it may not always be feasible as it has limitations in its effectiveness against certain contaminants, operates on relatively lengthy timeframes, and may not always achieve desired levels of decontamination. While the methods involved are not overly complex, designing and executing a successful bioremediation program may necessitate significant expertise and experience to thoroughly evaluate site suitability and optimize conditions for optimal results.

**Techniques involved in bioremediation are as follows:**

1. **Bioventing:** Bioventing is an in-situ bioremediation technique that involves supplying oxygen to the unsaturated (vadose) zone of the subsurface to stimulate the aerobic biodegradation of contaminants by indigenous microorganisms. Air is delivered through injection or extraction wells, and the technique is primarily used for treating fuels, volatile organic compounds (VOCs), semi-volatile organic compounds (SVOCs), pesticides, and herbicides. It requires the presence of indigenous microbes capable of degrading the target contaminants and necessary nutrients for their growth. Bioventing is relatively inexpensive, low-maintenance, and enjoys public acceptance. However, it is less effective in treating high water table areas, low moisture soils, and sites with high concentrations of inorganic salts, heavy metals, or organic compounds that inhibit microbial growth.

2. **Biostimulation:**Biostimulation involves introducing water-based solutions containing nutrients, electron acceptors (e.g., oxygen, nitrate), or other amendments to stimulate the growth and activity of indigenous microorganisms capable of degrading contaminants in soil and groundwater. It is designed for treating fuels, non-halogenated VOCs, SVOCs, pesticides, and herbicides. While biostimulation can be applied to halogenated organics, it may be less effective. The technique requires the presence of indigenous degraders and bioavailability of contaminants. It is not suitable for sites with high levels of inorganic salts, heavy metals, or organic compounds that inhibit microbial growth. It can be relatively inexpensive compared to other remediation methods but may increase contaminant mobility, necessitating treatment of underlying groundwater.

3. **Air sparging:** Air sparging is an in-situ technique that involves injecting air below the water table to promote aerobic biodegradation of groundwater contaminants by indigenous microorganisms. It is primarily used for treating fuels, non-halogenated VOCs, SVOCs, pesticides, organics, and herbicides. Air sparging has also been demonstrated to be effective for remediating petroleum hydrocarbon-contaminated aquifers. Like other bioremediation techniques, it requires the presence of indigenous degraders, nutrients, and bioavailability of contaminants. Air sparging can be relatively inexpensive and low-maintenance but is not applicable in sites with high levels of inorganic salts, heavy metals, or organic compounds that inhibit microbial growth.

4. **Natural attenuation:** Natural attenuation, also known as passive remediation or intrinsic bioremediation, is a proactive approach that relies on monitoring and verifying natural remediation processes without active intervention. It is suitable for treating fuels, non-halogenated VOCs, SVOCs, pesticides, herbicides, and hydrophobic contaminants like high molecular weight PAHs that tend to sorb tightly to soil particles. Natural attenuation processes can be destructive (destroying the contaminant) or non-destructive (reducing contaminant concentrations). While relatively simple and low-cost, natural attenuation can be time-consuming, and contaminant plumes may migrate if attenuation rates are too slow. It requires long-term monitoring and may face public scepticism due to the perception of inaction.

5. **Landfarming:** Landfarming involves excavating contaminated soil and applying it to the soil surface, where it is periodically tilled to mix and aerate the material, promoting biodegradation, transformation, and immobilization of contaminants through biotic and abiotic reactions. It is primarily used for treating soil contaminated with fuels, PAHs, non-halogenated VOCs, SVOCs, pesticides, and herbicides. While simple and inexpensive, landfarming requires large areas and may sometimes result in contaminant volatilization rather than biodegradation. Liners or other methods may be used to control leachate.

6. **Phytoremediation:** Phytoremediation is the use of plants to remove, degrade, or immobilize contaminants in soil and groundwater. It can be used for remediating toxic heavy metals, radionuclides, organic contaminants like chlorinated solvents, BTEX compounds, non-aromatic petroleum hydrocarbons, nitrotoluene ammunition wastes, and excess nutrients. Plants facilitate remediation through direct uptake and incorporation of contaminants, immobilization (phytostabilization), release of enzymes that act on contaminants, and stimulation of microbially mediated degradation in the rhizosphere. Phytoremediation is a relatively low-cost and environmentally friendly approach but is limited by factors such as depth of contamination, time requirements, and the ability of the site to support plant growth. Types of phytoremediation techniques are as follows:

**A. Phytoextraction:** This technique utilizes plants that can accumulate and concentrate high levels of contaminants, especially heavy metals, in their biomass from the soil or water. The contaminated plant parts can then be harvested and

disposed of safely. It is particularly useful for removing metals like lead, cadmium, zinc, nickel, and arsenic from contaminated sites. Key factors include selecting hyperaccumulator plants, enhancing metal bioavailability through soil amendments, and optimizing plant growth conditions.

**B. Phytostabilization:** In this approach, metal-tolerant plants are used to immobilize and reduce the mobility of contaminants, primarily heavy metals, in the soil. The plants help stabilize the contaminated soil, preventing erosion, leaching, and further spread of the contaminants. This technique is useful for stabilizing and revegetating metalliferous mine wastes, tailings, and other contaminated sites.

**C. Phytodegradation:** This involves the breakdown and degradation of organic contaminants, such as chlorinated solvents, herbicides, and ammunition wastes, by enzymes produced within the plant tissues after uptake. The contaminants are transformed into simpler, less toxic molecules that can be incorporated into the plant biomass or released into the environment.

**D. Phytovolatilization:** Plants and associated soil microbes are used to take up certain contaminants, like selenium, arsenic, and mercury, from the soil, transform them into volatile forms, and release them into the atmosphere. This technique is primarily employed for removing mercury and selenium from contaminated sites.

**7. Composting:** Composting is a controlled biological process in which organic materials are decomposed by microorganisms under aerobic conditions. The process involves several stages, including the mesophilic, thermophilic, and maturation phases. During composting, various factors such as temperature, moisture content, pH, aeration, and the carbon-to-nitrogen ratio are carefully controlled to optimize the degradation of organic contaminants. Successful application of composting for the bioremediation of polycyclic aromatic hydrocarbons (PAHs), petroleum hydrocarbons, phenol derivatives, polychlorinated biphenyls (PCBs), phthalic acid esters, and pesticides is possible. Several studies have demonstrated the ability of composting to reduce the concentrations of these contaminants through biodegradation, volatilization, and other mechanisms.

**8. Vermistabilization:** Vermistabilization, also known as vermicomposting, is a composting process that involves the use of earthworms to break down organic matter and facilitate the biodegradation of contaminants. Earthworms play a crucial role in aerating and bioturbating the soil, improving its nutritional status and fertility, and hindering the binding of organic contaminants to soil particles, thereby increasing their bioavailability. Earthworm species like *Eisenia fetida*, *Lumbricus terrestris*, and *Allobophora chlorotica* have been found to facilitate the removal of heavy metals, PAHs, PCBs, atrazine, and metamitron from contaminated soil during vermicomposting.

**9. Rhizofiltration:** It involves using plant roots to absorb, precipitate, and concentrate contaminants, primarily heavy metals, from contaminated water sources. The water is passed through a network of plant roots that can sorb or uptake the contaminants. The roots can then be harvested, processed to recover the metals, or disposed of safely. This technique is suitable for treating large volumes of water with low concentrations of metal contaminants.

**10. Controlled solid-phase biotreatment:** These processes involve the ex-situ treatment of contaminated soils in controlled environments, such as prepared treatment beds, biotreatment cells, and soil piles or biopiles. The contaminated soil is excavated and mixed with soil amendments, and parameters like moisture, heat, nutrients, oxygen, and pH are carefully controlled to enhance biodegradation. These techniques differ from traditional landfarming in that the treatment processes are often enclosed to control off-gases and leachate. They offer the advantage of containing toxic byproducts or metabolites formed during the biodegradation process.

**11. Slurry-phase bioremediation:** Slurry-phase bioremediation involves the treatment of excavated contaminated soils in a controlled bioreactor environment. The soil is processed to separate stones and rubble, then mixed with water to form a slurry, typically containing 10-40% solids. Electron acceptors, nutrients, and specialized microorganisms (if necessary) are added to the bioreactor, and parameters like pH and temperature are controlled to optimize biodegradation processes. This technique is favored over in-situ biological treatments for heterogeneous soils, low permeability soils, or when faster treatment times are required. It also has the advantage of containing toxic metabolites produced during biodegradation.

**12. Biosorption:** Biosorption is a highly promising bioremediation technique for removing heavy metal ions from aqueous solutions, particularly due to their extreme toxicity and persistence in the environment. This process involves the binding and concentration of contaminants by certain types of inactive or dead microbial biomass. One of its significant advantages is its cost-effectiveness, coupled with high efficiency and the potential for metal recovery. Unlike conventional treatment methods, biosorption minimizes the generation of chemical or biological sludge and allows for the regeneration of biosorbents. Recently, attention has shifted towards utilizing low-cost renewable organic materials, such as waste byproducts from industrial operations and agricultural residues, as effective biosorbents. These materials have shown remarkable abilities to accumulate heavy metals, including lead, cadmium, uranium, copper, zinc, and chromium, with some biosorbents capable of binding multiple metal types simultaneously. The choice of biomass for biosorption experiments depends on factors such as its origin, availability, and specificity towards certain metal ions. Various sorption processes, such as ion exchange, complexation, chelation, microprecipitation, and oxidation/reduction, contribute to the complex phenomenon of biosorption. Understanding the scientific principles, kinetics, thermodynamics, and equilibrium interactions of biosorption is crucial for optimizing this bioremediation technique, which has been extensively explored and documented in the scientific literature. Batch and continuous modes of operation are commonly employed in laboratory-scale biosorption processes, with factors like pH, temperature, ionic strength, biosorbent dosage, and agitation rate

357 influencing bacterial batch biosorption. Overall, biosorption holds significant potential as a sustainable and effective  
358 approach for mitigating heavy metal pollution in water bodies and industrial effluents.  
359

## 360 **6. CONCLUSION**

361 Heavy metal contamination poses significant threats to soil health and crop productivity. The accumulation of  
362 heavy metals in soil can lead to toxic effects on plants, including inhibition of enzyme activity, oxidative stress,  
363 and nutrient displacement. Consequently, this contamination jeopardizes crop quality and yield, posing risks to  
364 human health through food chain transfer. However, bioremediation offers a promising solution to mitigate these  
365 adverse effects. Through the use of microorganisms and plants, bioremediation processes can detoxify  
366 contaminated soil by immobilizing, degrading, or extracting heavy metals. This eco-friendly approach not only  
367 restores soil health but also helps in the restoration of ecosystem balance. By harnessing the natural processes of  
368 microbial and plant interactions, bioremediation presents a sustainable and cost-effective strategy to combat  
369 heavy metal contamination, safeguarding both soil fertility and crop productivity for future generations. Embracing  
370 bioremediation practices can lead us towards a healthier environment and more resilient agricultural systems.  
371  
372

## 373 **7. FUTURE PROSPECTS**

374 Looking ahead, the future prospects for bioremediation in addressing heavy metal contamination are promising.  
375 Continued research and technological advancements hold the potential to enhance the efficiency and scalability of  
376 bioremediation techniques. Furthermore, integrating bioremediation strategies with other sustainable agricultural practices  
377 could offer holistic solutions for managing heavy metal pollution while promoting soil health and crop productivity.  
378 Additionally, increasing awareness and adoption of bioremediation among farmers, policymakers, and industries can  
379 accelerate its implementation on a larger scale. By embracing bioremediation as a key component of environmental  
380 stewardship and sustainable agriculture, we can strive towards a cleaner, healthier, and more resilient future for our soil  
381 and crops.  
382

## 383 **8. COMPETING INTERESTS**

384 The authors assert with utmost clarity that they do not possess any competing interests.  
385

## 386 **9. REFERENCES**

- 387 1. Bharti, R., & Sharma, R. (2022). Effect of heavy metals: An overview. *Materials Today: Proceedings*, 51, 880-885.
- 388 2. Hawkes JS (1997) Heavy metals. *J Chem Educ* 74:1369–1374
- 389 3. Tchounwou PB, Yedjou CG, Patlolla AK, Sutton DJ (2012) Heavy metal toxicity and the environment. In:  
390 *Molecular, clinical and environmental toxicology*. Springer, Basel, pp 133–16
- 391 4. Oliveira A, Pampulha ME (2006) Effects of long-term heavy metal contamination on soil microbial characteristics.  
392 *J BiosciBioeng* 102(3):157–161
- 393 5. Chatterjee, J., & Chatterjee, C. (2000). Phytotoxicity of cobalt, chromium and copper in cauliflower. *Environmental*  
394 *pollution*, 109(1), 69-74.
- 395 6. Öncel, I., Keleş, Y., & Üstün, A. S. (2000). Interactive effects of temperature and heavy metal stress on the growth  
396 and some biochemical compounds in wheat seedlings. *Environmental pollution*, 107(3), 315-320.
- 397 7. Oancea, S., Foca, N., & Airinei, A. (2005). Effects of heavy metals on plant growth and photosynthetic activity.  
398 *AnaleleUniv. "Al. I. Cuza*, 1, 107-110.
- 399 8. Wintz, H., Fox, T., & Vulpe, C. (2002). Responses of plants to iron, zinc and copper deficiencies. *Biochemical*  
400 *Society Transactions*, 30(4), 766-768.
- 401 9. Stohs, S. J., & Bagchi, D. (1995). Oxidative mechanisms in the toxicity of metal ions. *Free radical biology and*  
402 *medicine*, 18(2), 321-336.
- 403 10. EPA, U. (1996). Integrated Risk Information System (IRIS)[online electronic data file]. US Environment Protection  
404 Agency, Office of research and development. National Center for Environment Assessment. Last updated  
405 December, 1.
- 406 11. Marques, A. P., Rangel, A. O., & Castro, P. M. (2009). Remediation of heavy metal contaminated soils:  
407 phytoremediation as a potentially promising clean-up technology. *Critical Reviews in Environmental Science and*  
408 *Technology*, 39(8), 622-654.  
409  
410  
411

- 412 12. Goyal, D., Yadav, A., Prasad, M., Singh, T. B., Shrivastav, P., Ali, A., Dantu, P. K., & Mishra, S. (2020). Effect of  
413 heavy metals on plant growth: An overview. In *Contaminants in Agriculture: Sources, Impacts, and Management*  
414 (pp. 79-101).
- 415 13. Nagajyoti PC, Lee KD, Sreekanth TVM (2010) Heavy metals, occurrence and toxicity for plants: a review. *Environ*  
416 *Chem Lett.* 8:199–216
- 417 14. Seaward MRD, Richardson DHS (1990) Atmospheric sources of metal pollution and effects on vegetation. In:  
418 Shaw AJ (ed) *Heavy metal tolerance in plants evolutionary aspects*. CRC Press, Boca Raton, pp 75–94
- 419 15. Eshleman A, Siegel SM, Siegel BZ (1971) Is mercury from Hawaiian volcanoes a natural source of pollution?  
420 *Nature* 223:471–475
- 421 16. Guan Q, Wang F, Xu C, Pan N, Lin J, Zhao R, Luo H (2018) Source apportionment of heavy metals in agricultural  
422 soil based on PMF: a case study in Hexi Corridor, northwest China. *Chemosphere* 193:189–197
- 423 17. Lacerda LD (1997) Global mercury emissions from gold and silver mining. *Water Air Soil Pollution* 97:209–221
- 424 18. Verkleji JAS (1993) The effects of heavy metals stress on higher plants and their use as bio monitors. In: Markert  
425 B (ed) *Plant as bioindicators: indicators of heavy metals in the terrestrial environment*. VCH, New York, pp 415–  
426 424
- 427 19. Angino EE, Magnuson LM, Waugh TC, Galle OK, Bredfeldt J (1970) Arsenic in detergents possible danger and  
428 pollution hazard. *Science* 168:389–392
- 429 20. Ramos, L., Hernandez, L. M., & Gonzalez, M. J. (1994). Sequential fractionation of copper, lead, cadmium and  
430 zinc in soils from or near Donana National Park (Vol. 23, No. 1, pp. 50-57). American Society of Agronomy, Crop  
431 Science Society of America, and Soil Science Society of America.
- 432 21. Harter, R. D. (1983). Effect of soil pH on adsorption of lead, copper, zinc, and nickel. *Soil Science Society of*  
433 *America Journal*, 47(1), 47-51.
- 434 22. Yi, L., Hong, Y., Wang, D., & Zhu, Y. (2007). Determination of free heavy metal ion concentrations in soils around  
435 a cadmium rich zinc deposit. *Geochemical Journal*, 41(4), 235-240.
- 436 23. Sharma, M. R., & Raju, N. S. (2013). Correlation of heavy metal contamination with soil properties of industrial  
437 areas of Mysore, Karnataka, India by cluster analysis. *International Research Journal of Environment Sciences*,  
438 2(10), 22-27.
- 439 24. Norvell, W. A. (1984). Comparison of chelating agents as extractants for metals in diverse soil materials. *Soil*  
440 *Science Society of America Journal*, 48(6), 1285-1292.
- 441 25. McBride, M. B., & Martínez, C. E. (2000). Copper phytotoxicity in a contaminated soil: remediation tests with  
442 adsorptive materials. *Environmental Science & Technology*, 34(20), 4386-4391.
- 443 26. Magnuson, M. L., Kelty, C. A., & Kelty, K. C. (2001). Trace metal loading on water-borne soil and dust particles  
444 characterized through the use of Split-flow thin-cell fractionation. *Analytical Chemistry*, 73(14), 3492-3496.
- 445 27. Friedlova, M. (2010). The influence of heavy metals on soil biological and chemical properties.
- 446 28. Nannipieri, P., Badalucco, L., Landi, L., & Pietramellara, G. (1997). Measurement in assessing the risk of  
447 chemicals to the soil eco system. In *Ecotoxicology: Responses, Biomarkers and Risk Assessment* (pp. 507-534).  
448 OECD.
- 449 29. Abedin, M. J., Feldmann, J., & Meharg, A. A. (2002). Uptake kinetics of arsenic species in rice plants. *Plant*  
450 *physiology*, 128(3), 1120-1128.
- 451 30. Guidotti, T. L., McNamara, J., & Moses, M. S. (2008). The interpretation of trace element analysis in body  
452 fluids. *Indian Journal of Medical Research*, 128(4), 524-532.
- 453 31. Ksheminska, H., Jaglarz, A., Fedorovych, D., Babyak, L., Yanovych, D., Kaszycki, P., & Koloczec, H. (2003).  
454 Bioremediation of chromium by the yeast *Pichia guilliermondii*: toxicity and accumulation of Cr (III) and Cr (VI) and  
455 the influence of riboflavin on Cr tolerance. *Microbiological research*, 158(1), 59-67.
- 456 32. Martin, S., & Griswold, W. (2009). Human health effects of heavy metals. *Environmental Science and Technology*  
457 *briefs for citizens*, 15(5).
- 458 33. Chibuikwe, G. U., & Obiora, S. C. (2014). Heavy metal polluted soils: effect on plants and bioremediation methods.  
459 *Applied and environmental soil science*, 2014.
- 460
- 461