

## Review Article

# Effectiveness of soil amended biochar for improving the vegetable production, soil health and environmental considerations

### Abstract

Increased emphasis on sustainable growth in agriculture without endangering the environment has assumed greater importance. The introduction of biochar has the potential to increase crop yields and soil fertility in agriculture while reducing greenhouse gas emissions at the same time. Biochar production technology and biomass type has a great influence on its quality and yield which in turn is responsible for altering soil fertility status and enhancing other soil properties. Biochar is obtained by thermochemical conversion from biomasses by the process of pyrolysis and different pyrolysis conditions which include heating rate, temperature or residence time affects the product distribution and yield for biochar produced from same biomass. Biochar is produced sustainably from eco-friendly sources under manageable conditions such as agricultural woody and non-woody wastes or crop residues or animal wastes. Biochar's acknowledgement in removing heavy toxic metals, organic solvents and pesticide residues from soil as well as in waste water management aids in environmental protection. Yield response of biochar produced from various feedstock sources on vegetable crop production is discussed to compare their relative effectiveness. Biochar is a carbonaceous substance produced that sequester carbon and remain in soil for thousands of years and thus offers stability in soil. Biochar as Negative Emissions Technology (NET) in removing CO<sub>2</sub> from atmosphere and other greenhouse gases such as N<sub>2</sub>O and CH<sub>4</sub> is a prominent methodology in mitigating serious concern of climate change. Biochar The objective of this review paper is to extensively justify the role and importance of biochar on agricultural fields in long run to harness the benefits for sustainable growth.

**Keywords:** biochar, biomass, negative emissions technology, carbon sequestration

### Introduction?

The major setback faced by agriculture globally is decline in productivity of agricultural lands owing to the poor soil condition. Intensive farming practices over decades has degraded soils and has reduced the potential of soils to sustain plant health causing decline in yields. To

**Comment [i1]:** Effectiveness of soil-amended biochar in improving crop production, soil health and environmental aspects

**Comment [i2]:** Emphasizing sustainable growth of agriculture without endangering the environment has become even more important. The introduction of biochar can increase yields and soil fertility in agriculture, reducing greenhouse gas emissions

**Comment [i3]:** Biochar is produced by thermochemical conversion of biomass using a pyrolysis process, and different pyrolysis conditions, such as heating, temperature or residence time, affect the product distribution and yield of biochar produced from the same biomass. Biochar is sustainably produced from environmentally friendly sources under controlled conditions, such as wood and non-wood agricultural waste or crop residues or animal waste.

**Comment [i4]:** in the cultivation of vegetable crops

**Comment [i5]:** to remove

**Comment [i6]:** from the atmosphere is an important method to mitigate

**Comment [i7]:** delete

**Comment [i8]:** It is preferable to write every 6-8 sentences in one paragraph. Applies to every article

**Comment [i9]:** the loss of

feed the growing population which is estimated to reach 9.8 billion by 2050 with shrinking land resources and infertile soils amid climate change stress is a challenging goal (Shubham *et al.* 2022). The fact that land under cultivation is limited, there is need to elevate the quality of soils under agriculture for better yield through application of soil remedy techniques. The negative consequences of green revolution on agricultural soils and environment have come to the fore. Soil has lost its healthy physical, chemical and biological properties over a period of time. The direct implications on soil properties have caused soil salinization, soil acidification, organic matter loss, soil compaction, desertification, loss of biodiversity in soil environment and decreased productivity. It is imperative to take essential measures for improvement in agriculture with adaptability to climate change and variations.

**Comment [i10]:** Feeding a

**Comment [i11]:** Since the cultivated area

**Comment [i12]:** it is necessary to improve the quality of the cultivated soil using soil improvement techniques to achieve better yields. The negative effects of the green revolution on agriculture and the environment have emerged.

**Comment [i13]:** its

**Comment [i14]:** loss of organic matter

The increase in concentration of greenhouse gases (GHG's) is responsible for relative increase in global temperatures. The implications of rising CO<sub>2</sub> levels on equilibrium climate sensitivity and related concerns have risen substantially in the past few years. Methane and Nitrous oxide are equally responsible for driving global climate change. As per the report by Climate Watch (2023), Agriculture is the fourth largest sector contributor of GHG's globally in the year 2020 with 5.87 billion tonnes emissions measured in carbon dioxide-equivalents while agriculture being the dominant sector in different countries (Anonymous 2020). Among non-CO<sub>2</sub> greenhouse gases, Nitrous Oxide (N<sub>2</sub>O) and Methane (CH<sub>4</sub>) are the dominant gases responsible for greenhouse effect not in terms of mass since they make up only small fraction of GHG's emissions. For instance, according to a source, nitrous oxide warms the planet 300 times as much as carbon dioxide (Canfield *et al.*, 2010). As for Methane, one tonne of methane would generate 28 times the amount of warming as one tonne of CO<sub>2</sub> (Saunoi *et al.*, 2020). Agriculture sector is the leading cause of Methane and Nitrous oxide emissions at 3.54 and 2.33 billion tonnes respectively measured in carbon dioxide-equivalents in the year 2020 as per the data source of Climate Watch (2023) (Anonymous 2020). Fertilizer application to the soil has been known to decrease the CH<sub>4</sub> sink capacity and increased N<sub>2</sub>O and CH<sub>4</sub> emissions to the atmosphere. On an average, most of the applied fertilizers are lost as runoff or released into the atmosphere as GHG emissions due to multiple factors. Methane emissions are linked to livestock, rice cultivation, biomass burning, waste matter decomposition or fossil fuel production. On the other side, Nitrous Oxide emissions are majorly contributed by the application of nitrogenous fertilizers to soils.

**Comment [i15]:** temperature

**Comment [i16]:** According to a Climate Watch report

**Comment [i17]:** more than

**Comment [i18]:** According to the data source Climate Watch (2023) (Anonymous 2020), the agricultural sector is the largest source of methane emissions, with nitrous oxide emissions measured as carbon dioxide equivalent in 2020 between 3.54 and 2.33 billion tons.

**Comment [i19]:** Delete

**Comment [i20]:** mostly caused

**Comment [i21]:** nitrogen

Among various amendment measures, biochar is a unified approach for improving soil fertility as well as mitigating climate change. Biochar has wider applications possibly required not just to comply with food security demand but also to meet the sustainable goals of development. A wealth of benefits that biochar withhold include increased nutrient availability as well as fertility in soil, improved crop yield and reduction in GHG's emissions. Biochar is prepared from waste biomass and therefore is sustainable and has an economical approach since it can be produced at feasible cost. The range of waste biomass that can be used in biochar preparation is vast and thus aids in waste management. Biochar is a carbonaceous material produced from biomass residue that plays an important role in enriching soil carbon pool. The concept of capturing carbon and

**Comment [i22]:** that may be needed to meet not only

**Comment [i23]:** residues

**Comment [i24]:** supply

storing it in agricultural lands seems to be an innovative approach that can be defined as carbon sequestration. However, it would be resourceful to not only capture carbon emissions from bulk industrial sources but also channelizing the focus to sequester carbon from crop residues rich in biomass. Biochar preparation is the potent way of converting residues into useful resources. Production of biochar offers multitude of advantages ranging from production of renewable fuels to reduction of methane and nitrous oxide emissions from soil. Biochar is an excellent soil conditioner with potent ability to reduce the impact of heavy metal pollution in soil. Therefore, it is crucial to acknowledge the scope of biochar for agricultural lands in long run. The factors responsible for quality of biochar and their relative soil dynamics are further explored. A comprehensible knowledge of influence of biochar on vegetable cropping system is reviewed to allow grower's make robust decisions for enhanced productivity. The utilization of biochar to decrease greenhouse gas (GHG) footprint per unit yield within agricultural ecosystems is studied to not just illustrate potentials but to occupy knowledge gap to maximize the benefits. Biochar's recognition as negative emissions technology has also been discussed to unleash the comparative edge over other options in mitigating climate change.

**Comment [i25]:** Agricultural land carbon capture and storage appears

**Comment [i26]:** containing a lot of

**Comment [i27]:** producing

**Comment [i28]:** reducing

**Comment [i29]:** an effective

**Comment [i30]:** contamination

**Comment [i31]:** of crop in agroecosystems was

**Comment [i32]:** climate change mitigation options.

### Decisive factors influencing biochar yield

The key factors that influence the yield and properties of biochar include feedstock material used in biochar preparation and processing conditions which in turn determine its role as a soil applicator for soil remediation and productivity. Organic matter undergoes thermal transformation in an oxygen-limited environment to produce biochar. The common thermochemical techniques used for biochar production include pyrolysis, hydrothermal carbonization, gasification, flash carbonization and torrefaction (Yaashika *et al.* 2020). Pyrolysis has been considered as the most cost-effective and highly efficient method for biochar production (Cha *et al.* 2016). During Pyrolysis, besides biochar, bio-oil (can be used as energy source) and gaseous by-products which is a mixture of CO, CO<sub>2</sub>, H<sub>2</sub>, CH<sub>4</sub>, H<sub>2</sub>O and other volatile compounds are released. The biochar properties are heavily affected by the extent of pyrolysis (pyrolytic temperature and pressure) and entirely by the size of biomass and kiln (or furnace) residence time (Anonymous 2018). Pyrolysis is divided into slow pyrolysis (300 -700°C), fast pyrolysis (600 -1000°C), intermediate and flash pyrolysis (> 1000 °C). The data in table 1 provides the range of operating parameters and product distribution of different categories of pyrolysis processes (Kazawadi *et al.* 2021). From table 1, it can be concluded that although fast and flash pyrolysis allows for rapid thermal decomposition but main products are gases and bio-oil (thinner). On the contrary, slow and intermediate pyrolysis yields char, gases and bio-oil (tar) as major products. It can be explained by the fact that as the pyrolysis temperature increase, moisture and volatile matter content decrease while fixed-carbon content increase. Also, ash content per unit mass increase with increase in temperature. Figure 1 summarizes different pyrolysis conditions and the effect on product distribution. Therefore, it is evident that each type of pyrolysis differs in temperature, residence time, heating rate and products obtained.

**Comment [i33]:** influencing

**Comment [i34]:** conditions,

**Comment [i35]:** spreader in improving land and increasing

**Comment [i36]:** delete

**Comment [i37]:** was

**Comment [i38]:** are released, which are

**Comment [i39]:** compounds

**Comment [i40]:** biomass size

**Comment [i41]:** It can be concluded from Table 1

**Comment [i42]:** decomposition, the

**Comment [i43]:** increases


**Comment [i44]:** according to

Property	Slow	Intermediate	Fast	Flash
Heating rate (°C/s)	1.1–1	1–10	10–200	>1000
Feed size (mm)	5–50	1–5	<1	<0.5
Reaction temperature (°C)	400–500	400–650	850–1250	>1000
Vapor residence time (s)	300–550	0.5–20	0.5–10	<1
Feed water content (%)	Up to 40	Up to 40	<<10	<<10
Biooil yield (%)	20–50	35–50	60–75	60–75
Biochar yield (%)	25–35	25–40	10–25	10–25
Gas yield (%)	20–50	20–30	10–30	10–30

[Source: Kazawadiet al. 2021]

**Table 1: Classification of pyrolysis processes**

Comment [i45]: write in above table



**Figure 1: Summarizing different pyrolysis conditions and the effect on product distribution.**

Comment [i46]: Summary of

Comment [i47]: effects

Credit: (Created based on Xavier DEGLISE, Emeritus Professor at University Henri Poincaré, France. 2006 BEEMS Module C2, Brian He.)

[Source: [4.1 Biomass Pyrolysis | EGEE 439: Alternative Fuels from Biomass Sources \(psu.edu\)](#)]

Comment [i48]: what is this ?

In order to maximize the biochar yield, the production technique must be compatible with feedstock properties. The feedstock material is classified into woody and non-woody substrates.

Biomass	Temp. (°C)	Yield (%)	Proximate analysis (wt. %)
---------	------------	-----------	----------------------------

Woody biomass is procured from trees and woody forest residues. Non-woody biomass is plant material from crop residues and agricultural lignocellulosic materials; animal and industrial solid wastes. Woody biomass is characterized by low moisture, low ash, high calorific value, high bulk density and less voidage; in contrast, non-woody biomass exhibits the opposite characteristics (Tomczyk *et al.* 2020). The analysis of heterogeneity of different biomasses is crucial for understanding their respective composition. There are several parameters to determine the compositional variance in different biomasses which are broadly categorized into physical and chemical properties of biomasses used in production of biochar. Physical properties include moisture content (%), water holding capacity (g water g<sup>-1</sup> dry), bulk density (kg m<sup>-3</sup>) and porosity (%). Chemical properties include pH, EC (dS m<sup>-1</sup>), Total organic matter (%), Total organic carbon (%), C/N ratio, Total nitrogen (%), Total potassium (%) and others. Animal biochars are mostly constituted of animal protein such as gelatin, collagen, and polysaccharides (cellulose, starch and carbohydrates) (Huang *et al.* 2021). Since majority of plant-based biomass is composed of cellulose, hemicellulose and lignin; these components undergo different reaction conditions to produce biochar (Yaashika *et al.* 2020). The pyrolysis temperature and feedstock type influences the physical and chemical properties of biochar due to stagewise decomposition of structure and chemical bonds (Huang *et al.* 2021). Different types of feedstocks respond heterogeneously to particular pyrolysis conditions due to variations of cellulose (C<sub>5</sub>H<sub>8</sub>O<sub>4</sub>)<sub>m</sub>, hemicelluloses (C<sub>5</sub>H<sub>8</sub>O<sub>4</sub>), lignin [C<sub>9</sub>H<sub>10</sub>O<sub>3</sub>(OCH<sub>3</sub>)<sub>0.9-1.7</sub>]<sub>n</sub> and inorganic mineral contents (Hassan *et al.*, 2020). It is evident that different materials yield varying amounts of biochar at different temperatures, for instance, in a study using different agricultural wastes (straw rice, sawdust, sugarcane and tree leaves), researchers found that biochar yield decreased variably as the pyrolysis temperature increased from 400 to 800-degree Celsius (Khater *et al.* 2024). Biochar yield decreases with increasing pyrolysis temperature as a result of the increased burning rate, because of the variation of lignin and cellulose content of biomass and conversion of organic matter to ash, which reduced the carbon content of the biochar (Khater *et al.* 2024) but it yields more energy and offers more stability of carbon in soil. In table 2, Yield and proximate analysis (volatile matter, ash, carbon fixed) of biochars produced at different pyrolysis temperatures for animal source biomass and plant-based biomass is demonstrated (Domingues *et al.* 2017). It can be analyzed from the data given in table 2 that with increase in temperature,

Comment [i49]: Wood biomass is obtained

Comment [i50]: lignocellulose

Comment [i51]: variety of different biomasses,

Comment [i52]: total

Comment [i53]: proteins

Comment [i54]: minerals

Comment [i55]: However, the biochar production decreases

Comment [i56]: , produces more energy, and ensures greater carbon stability in the soil.

			<b>Volatile Matter</b>	<b>Ash</b>	<b>Carbon Fixed</b>
Chicken manure	350	69.7	36.9 Ab	52.0 Ba	11.1 Cd
	450	63.0	30.6 Ba	55.3 Aa	14.1 Be
	750	55.9	26.5 Ca	56.4 Aa	17.0 Ae
Eucalyptus sawdust	350	42.5	36.9 Ab	0.9 ABe	62.2 Cb
	450	36.0	28.5 Bb	0.7 Be	70.8 Bb
	750	28.2	6.5 Cd	1.1 Ae	92.4 Aa
Coffee husk	350	43.5	34.6 Ac	12.9 Bb	52.5 Cc
	450	37.7	26.2 Bc	12.9 Bb	60.9 Bc
	750	31.6	17.6 Cb	19.6 Ab	62.8 Ad
Sugarcane bagasse	350	37.5	35.0 Ac	1.9 Ad	63.0 Ca
	450	33.2	24.0 Bd	2.1 Ad	73.9 Ba
	750	26.9	7.7 Cc	2.2 Ad	90.1 Ab
Pine bark	350	59.6	38.5 Aa	8.3 Bc	53.2 Cc
	450	49.3	29.3 Ba	7.9 Bc	62.8 Bc

**Comment [157]:** is prefer to write the Duncan latter in small ,

	750	38.9	6.0 Cd	14.5 Ac	79.4 Aa
--	-----	------	--------	---------	---------

biochar yield decreases for any feedstock source. However, it is evident from the figures in table that Chicken manure (animal-based biomass) showed highest yield and ash content at temp 350-degree Celcius (slow pyrolysis) comparative to plant-based biomasses.

Comment [i58]: ?, it's not clear

Comment [i59]: °C

[Source: Domingues *et al.* 2017]

Uppercase letters compare pyrolysis temperatures within the same biomass and lowercase letters compare biomass at the same temperature. The same letter do not differ by the Tukey test at  $p < 0.05$ .

Comment [i60]: what is this ?

**Table 2: Yield and proximate analysis (volatile matter, ash, carbon fixed) of biochars produced at different pyrolysis temperatures.**

Comment [i61]: write above the table

### Quality of biochar and related soil-dynamics

Comment [i62]: Biochar quality and related soil dynamics

The IBI (2015) has formalised various physical parameters linked to biochar quality which include pH, volatile compound content, water holding capacity, ash content, bulk density, pore volume, and specific surface area. The nutrient content and physico-chemical properties of biochar is a product of feedstock material, activators or amendments added and processing conditions. It can be noted that biochar produced from same feedstock material varies in properties due to varying conditions of pyrolysis process. For instance, Biochar produced from straw-based feedstock and low-temperature pyrolysis exhibits numerous surface functional groups, enhancing its surface activity and adsorption capacity, resulting in increased active sites for biochar surface adsorption, reduced energy requirements for reactions, enhanced atomic bonding orbitals, and improved potential for charge transfer between biochar and soil adsorbates (He *et al.* 2024). Further, application of biochar for desired purpose in soil is calibrated by balancing the parameters based on needs. The impact of biochar application in soil is not solely the function of biochar properties but includes analysis of several factors. These factors include type of climate and soil, physico-chemical properties of soil, crop-type, biochar application rate and method, treatments applied, post-application management and so on. Biochar-soil dynamics thus yields conceptual framework for analysing its effectiveness. The following table indicates the effectiveness of application of different types of biochar on yield of vegetable crops, its role in soil nutrient enhancement and remediation as well environmental impact.

Comment [i63]: IBI (2015) formalized

Comment [i64]: related

Comment [i65]: compounds

Comment [i66]: nutritional

Comment [i67]: the properties of biochar produced from the same raw material differ due to the different conditions of the

Comment [i68]: In addition, the

Comment [i69]: soil type

Comment [i70]: , etc.

Comment [i71]: provides a

Comment [i72]: in the cultivation

Biochar type	Yield response on vegetable crops	Impact on soil health	Environmental impact	References

<b>Rice Husk Biochar</b>	<p>In acidic soil, biochar treatment yielded 212% more yield than the control in chili pepper, 210% in tomato, 182% in soybean and 612% in yard long bean. While in neutral soil, biochar treatment yielded 76% more tomato yield than the control; and in alkaline soil, biochar treatment yielded 189% more than the control in chili pepper, 223% in sweet pepper, 139% in Phaseolus bean and 303% in carrot.</p>	<p><b>Alkaline Soil-</b>No significant response in alkaline soil</p> <p><b>Acidic Soil-</b>improvement of soil physical properties such as permeability, water holding capacity and nutrient use efficiency.</p>	<p>1. Biochar has also been extensively used to remove different types of pollutants (e.g. heavy metals, dye, pharmaceuticals, pesticides/herbicides and phenols) from wastewaters.</p> <p>2. The high efficiency of biochar to adsorb heavy metals and organic pollutants is related to its high porosity, high number of functional groups and high pH.</p>	<p>Williams <i>et al.</i>(2023)</p> <p>Asadiet <i>al.</i>(2021)</p>
<b>Coconut Husk Biochar</b>	<p>18% increase in yield of Water spinach</p>	<ul style="list-style-type: none"> <li>○ Coconut husk increases water holding capacity, improves pH, organic carbon, porosity and nutrient availability in soil.</li> <li>○ It contains relatively high concentration of essential minerals</li> </ul>	<p>Soil amendment, Carbon sequestration and Water treatment are the environmental applications for coconut husk biochar.</p>	<p>Zhao <i>et al.</i> (2019)</p> <p>Nagulaet <i>al.</i> (2022)</p> <p>Ajienet <i>al.</i> (2022)</p>

		<p>including P, K, Na, Ca), Mg, Mo, Zn, Mn, Cu, Ni, Fe and Si.</p> <ul style="list-style-type: none"> <li>○ Coconut based biochar is alkaline and thus promotes higher microbial activity, higher organic matter mineralization, higher plant nutrient availability and higher acid neutralization.</li> </ul>		
<b>Wheat straw Biochar</b>	<p>Amending the soil of lower fertility with wheat straw biochar of concentration BC<sub>3</sub>% (Biochar concentration) significantly increased green pepper yield, due to improved soil chemical properties (pH, P, K, SOM, and CEC), and enhanced water and nutrient use.</p>	<ul style="list-style-type: none"> <li>○ Increase the sorption of deltamethrin in soil. The enhanced sorption of deltamethrin in the biochar amended soil could reduce its leachability in the soil.</li> <li>○ Therefore, wheat straw biochar can immobilize deltamethrin in the soil to a certain extent and minimize deep leachability, reduce the risk of pesticide contamination</li> </ul>	<p><b>1.</b> Wheat straw biochar aids in mitigating climate change by decreasing the N<sub>2</sub>O and CO<sub>2</sub> emissions from agricultural soils.</p> <p><b>2.</b> Wheat straw biochar application reduces the sensitivity to temperature changes of the decomposition of the new organic matter added to the soil.</p>	<p>Purkaysthaet <i>al.</i> (2022)</p> <p>Palangiet <i>al.</i> (2021)</p> <p>Duanet <i>al.</i> (2020)</p>
<b>Sugarcane Bagasse biochar (SBB)</b>	<p>Yield contributing characters of Okra treated with 10% SBB</p>	<p>It enhances the water holding capacity of saline soil which has high seepage problem. It also enhances soil</p>	<p>Increasing concentration of nitrate in drinking water can cause</p>	<p>Sikdaret <i>al.</i> (2023)</p> <p>Nieet <i>al.</i> (2018)</p>

	+ CF treatment is increased significantly as compared to no control treatment  PL- 113% (↑) IPW-88% (↑) PD-69.7% (↑) NPP-32.8% (↑) PWP-113% (↑) NSP- 148% (↑)	quality due to its high surface area and surface chemical functional groups	harmful effect on human health, Sugarcane Bagasse biochar is an effective adsorbent for the nitrate adsorption from aqueous solution.	Hafshejaniet al. (2016)
<b>Chicken Manure Biochar</b>	An increase in yield is seen by 23.69% when chicken biochar is used in red chilli.	<ul style="list-style-type: none"> <li>○ It can help in increasing the availability of phosphorus in the soil.</li> <li>○ It contributes complete nutrients needed by plants. It increases soil pH and soil aggregates, increases soil water content.</li> <li>○ Increase the ability of soil to provide Ca, Mg, P and K, increase soil microbial respiration, increase cation exchange capacity.</li> </ul>	Chicken manure-based biochar can be used as an effective adsorbent for removing contaminants from water. For instance, it has been explored for arsenic removal.	Mahendra et al. (2020)  Babaei et al. (2023)

**Abbreviations:** PL- Pod length; IPW- Individual pod weight; PD- Pod diameter; NPP- No. of pods/plant; PWP- Pod weight/ plant; NSP- No. of seeds/ plant; SBB- Sugarcane Bagasse Biochar; CF- Chemical Fertilizer treatment

**Table 3: Yield response of different biochar types on vegetable crops, their environmental impact and impact on soil health**

**Comment [173]:** Write above the table

## Biochar in mitigating climate change

**Comment [i74]:** Biochar in Climate Change Mitigation

Crop residue accounts for global concern as how to manage it effectively by not polluting the surroundings further. For safe disposal of agricultural residues, biochar is the convenient approach which not only allows residue management but enhances soil health, reduces GHG emissions as well as provides the most stable carbon source that takes thousands of years to degrade. Biochar sequester carbon from environment and has the potential to remove 6% of global emissions which is significant (Anonymous 2023). The carbon that would otherwise be released into environment from degradation of residues is being stored by converting the biomass to biochar. Biochar application has been known to reduce nitrous oxide (N<sub>2</sub>O) emissions from soil (Woolf *et al.* 2018). The biochar has a strong adsorption capacity due to its high porosity, large specific surface area, and functional group properties on the surface and therefore, biochar can reduce N<sub>2</sub>O emissions by absorbing NH<sub>4</sub><sup>+</sup>-N and NO<sub>3</sub><sup>-</sup>-N which are N<sub>2</sub>O precursors, as well as directly capturing N<sub>2</sub>O (Liang *et al.* 2023). Methane emissions from soil is attributed majorly to microbial activities in soil. Global methane fluxes are net positive as rice cultivation is a much larger source of CH<sub>4</sub> than the sink contribution of upland soils (Jeffery *et al.* 2016). Biochar has more porous structure and high surface area which allows for adsorption of methane molecules and it also provides space for methane oxidizing microbes. Long term application of biochar in soils can help mitigate global temperature incline by reducing the GHG emissions. Apart from this biochar's role as negative emissions technology for removing CO<sub>2</sub> from atmosphere is established by the 2018 report of Intergovernmental Panel on Climate Change (IPCC). Biochar is an effective NET since it is produced from sequestered atmospheric carbon and is highly stable, hence, carbon is stored in soil for longer period that would otherwise be released into atmosphere.

**Comment [i75]:** emissions,

**Comment [i76]:** delete

**Comment [i77]:** The use of biochar is known to reduce nitrogen

**Comment [i78]:** Due to its high porosity, high specific surface area and surface functional group properties, biochar has a strong adsorption capacity, and therefore

**Comment [i79]:** positive because

**Comment [i80]:** a more

**Comment [i81]:** a high

## Conclusion

Biochar is a single promising solution for multiple environmental remediations including soil health improvement, improving crop yields, agricultural waste management, climate change mitigation, waste water treatment and water conservation. Physico-chemical properties exhibited by biochar is a function of availability of wide range of biomasses including both plant and animal sources and different operational parameters of biochar production. Bio-oil and gaseous by-products are yielded during biochar production that can be harvested using high-end technology. Biochar dynamics in soil depends on several factors from climate to soil. Biochar aids in increasing yield of different crops. Biochar is a definite technology for managing global temperatures in desired range by reducing impact of GHG emissions.

**Comment [i82]:** one

**Comment [i83]:** many

**Comment [i84]:** factors,

**Comment [i85]:** Biochar helps to increase the yield of various crops.

## References?

1. Ajien, A., Idris, J., Sofwan, N.M., Husen, R. & Seli, H. Coconut shell and husk biochar: A review of production and activation technology, economic, financial aspect and application. *Waste Management & Research*. (2023);41(1):37-51.  
DOI: <https://doi.org/10.1177/0734242X221127167>
2. Anonymous (2018). (1) (PDF) Biochar and its Application (researchgate.net)
3. Anonymous (2020). Emissions by sector: where do greenhouse gases come from? - Our World in Data
4. Anonymous (2023). What is the future of biochar as a climate change mitigation tool? | World Economic Forum (weforum.org)
5. Anonymous. 4.1 Biomass Pyrolysis | EGEE 439: Alternative Fuels from Biomass Sources (psu.edu)
6. Asadi, H., Ghorbani, M., Rashti, M.R., Abrishamkesh, S., Amirahmadi, E., Chengrong, C. & Gorji, M. (2021). Application of Rice Husk Biochar for Achieving Sustainable Agriculture and Environment. *Rice Science*, Volume 28, Issue 4, Pages 325-343, ISSN 1672-6308, DOI: <https://doi.org/10.1016/j.rsci.2021.05.004>.
7. Babaei, K., Salimi, S., Mahmoudabadi, Z.S. & Tavasoli, A. (2023). Synthesis and application of chicken manure biochar as an effective nanoporous adsorbent for removal of arsenic from wastewater: a cost-effective approach. *J IRAN CHEM SOC* **20**, 541–550.  
DOI: <https://doi.org/10.1007/s13738-022-02686-6>
8. Canfield, D.E., Glazer, A.N. & Falkowski, P.G. The Evolution and Future of Earth's Nitrogen Cycle. *Science* **330**, 192-196 (2010). DOI: [10.1126/science.1186120](https://doi.org/10.1126/science.1186120)
9. Cha, J.S., Park, S.H., Jung, S.C., Ryu, C., Jeon, J.K., Shin, M.C. & Park, Y.K. (2016). Production and utilization of biochar: A review. *Journal of Industrial and Engineering Chemistry*, Volume 40, Pages 1-15, ISSN 1226-086X.  
DOI: <https://doi.org/10.1016/j.jiec.2016.06.002>
10. Domingues, R.R., Trugilho, P.F., Silva, C.A., Melo, I.C., Melo, L.C., Magriotis, Z.M., & Sánchez-Monedero, M.A. (2017). Properties of biochar derived from wood and high-nutrient biomasses with the aim of agronomic and environmental benefits. *PLoS ONE*, *12*. DOI: [10.1371/journal.pone.0176884](https://doi.org/10.1371/journal.pone.0176884)
11. Duan, M., Wu, F., Jia, Z., Wang, S., Cai, Y. & Chang, S.X. Wheat straw and its biochar differently affect soil properties and field-based greenhouse gas emission in a Chernozemic soil. *Biol Fertil Soils* **56**, 1023–1036 (2020). DOI: <https://doi.org/10.1007/s00374-020-01479-4>
12. Hafshejani, L.D., Hooshmand, A., Naseri, A.A., Mohammadi, A.S., Abbasi, F. & Bhatnagar, A. (2016). Removal of nitrate from aqueous solution by modified sugarcane bagasse

**Comment [i86]:** Review the sources. Some sources include the year of publication at the end, the source, and others after the names of the researchers. Therefore, it is preferable to unify the sources according to the magazine's instructions

**Comment [i87]:** Rewrite the references in right form

- biochar. *Ecological Engineering*, Volume 95, Pages 101-111, ISSN 0925-8574, DOI: <https://doi.org/10.1016/j.ecoleng.2016.06.035>.
13. Hassan, M., Liu, Y., Naidu, R., Parikh, S. J., Du, J., Qi, F., & Willett, I. R. (2020). Influences of feedstock sources and pyrolysis temperature on the properties of biochar and functionality as adsorbents: A meta-analysis. *The Science of the total environment*, 744, 140714. DOI: <https://doi.org/10.1016/j.scitotenv.2020.140714>
  14. He, D., Luo, Y., & Zhu, B. (2024). Feedstock and pyrolysis temperature influence biochar properties and its interactions with soil substances: Insights from a DFT calculation. *The Science of the total environment*, 922, 171259. DOI: <https://doi.org/10.1016/j.scitotenv.2024.171259>
  15. Huang, H., Reddy, N.G., Huang, X., Chen, P., Wang, P., Zhang, Y., Huang, Y., Lin, P. & Garg, A. Effects of pyrolysis temperature, feedstock type and compaction on water retention of biochar amended soil. *Sci Rep* 11, 7419 (2021). DOI: <https://doi.org/10.1038/s41598-021-86701-5>
  16. Jeffery, S., Verheijen, F.G.A., Kammann, C. & Abalos, D. (2016). Biochar effects on methane emissions from soils: A meta-analysis. *Soil Biology and Biochemistry*, Volume 101, Pages 251-258, ISSN 0038-0717, DOI: <https://doi.org/10.1016/j.soilbio.2016.07.021>.
  17. Kazawadi, D., Ntalikwa, J. & Kombe, G. A Review of Intermediate Pyrolysis as a Technology of Biomass Conversion for Coproduction of Biooil and Adsorption Biochar", *Journal of Renewable Energy*, vol. 2021, Article ID 5533780, 10 pages, 2021. DOI: <https://doi.org/10.1155/2021/5533780>
  18. Khater, E. S., Bahnasawy, A., Hamouda, R., Sabahy, A., Abbas, W., & Morsy, O. M. (2024). Biochar production under different pyrolysis temperatures with different types of agricultural wastes. *Scientific reports*, 14(1), 2625. DOI: <https://doi.org/10.1038/s41598-024-52336-5>
  19. Liang, X., Zhou, W., Yang, R., Zhang, D., Wang, H., Li, Q., Qi, Z., Li, Y. & Lin, W. (2023). Microbial mechanism of biochar addition to reduce N<sub>2</sub>O emissions from soilless substrate systems. *Journal of Environmental Management*, Volume 348, 119326, ISSN 0301-4797. DOI: <https://doi.org/10.1016/j.jenvman.2023.119326>.
  20. Mahendra, K.A., Situmeang, Y.P. & Suarta, M.W. (2020). Effect of Biochar and Compost from Chicken, Goat, and Cow Manure on Cultivation of Red Chili (*Capsicum annum* L). (2020). *SEAS (Sustainable Environment Agricultural Science)*, Volume 04, Number 02, pages: 95~101, ISSN 2614-0934, DOI: <http://ejournal.warmadewa.ac.id/index.php/seas>
  21. Nagula, S., Usha, P.B., & Thampatti, M. & Joseph, B. (2022). Tender Coconut Husk Derived Biochar Impact on Soil Properties, Yield and Fruit Quality of Banana. *Journal of the Indian Society of Soil Science*. 69. 334-338. DOI: [10.5958/0974-0228.2021.00050.5](https://doi.org/10.5958/0974-0228.2021.00050.5)
  22. Nie, C., Yang, X., Niazi, N.K., Xu, X., Wen, Y., Rinklebe, J., Ok, Y.S., Xu, S. & Wang, H. (2018). Impact of sugarcane bagasse-derived biochar on heavy metal availability and

- microbial activity: A field study. *Chemosphere*, Volume 200, Pages 274-282, ISSN 0045-6535, DOI: <https://doi.org/10.1016/j.chemosphere.2018.02.134>.
23. Palangi, S., Bahmani, O. & Atlassi-pak, V. (2021). Effects of wheat straw biochar amendments to soil on the fate of deltamethrin and soil properties. *Environmental Technology & Innovation*, Volume 23, 101681, ISSN 2352-1864, DOI: <https://doi.org/10.1016/j.eti.2021.101681>.
  24. Purkaystha, J., Prasher, S., Afzal, M.T., Nzediegwu, C. & Dhiman, J. Wheat straw biochar amendment significantly reduces nutrient leaching and increases green pepper yield in a less fertile soil. (2022). *Environmental Technology & Innovation*, Volume 28, 102655, ISSN 2352-1864, DOI: <https://doi.org/10.1016/j.eti.2022.102655>.
  25. Saunio, M., Stavert, A. R., Poulter, B., Bousquet, P., Canadell, J. G., Jackson, R. B., ... & Ciais, P. (2020). The global methane budget 2000–2017. *Earth System Science Data*, 12(3), 1561-1623. DOI: <https://doi.org/10.5194/essd-12-1561-2020>
  26. Shubham, Sharma, R., Kaushal, Y. & Sharma, P. (2022). Effect of Forest Fires on Soil Carbon Dynamics in Different Land Uses under NW Himalayas. *Indian Journal of Ecology* 49(6):2322-2329. DOI: [10.55362/IJE/2022/3828](https://doi.org/10.55362/IJE/2022/3828)
  27. Sikdar, A., Zaman, M.N., Dev, S., Chakma, N. & Sharif, M.O. (2023). Effect of Co-application of Different Biochars and Inorganic Fertilizers on the Growth and Yield of Okra (*Abelmoschus esculentus* L.). *Journal of Agroforestry and Environment*. 16(2):20-27. DOI: [10.55706/jae1623](https://doi.org/10.55706/jae1623)
  28. Tomczyk, A., Sokołowska, Z. & Boguta, P. Biochar physicochemical properties: pyrolysis temperature and feedstock kind effects. *Rev Environ Sci Biotechnol* 19, 191–215 (2020). DOI: <https://doi.org/10.1007/s11157-020-09523-3>
  29. Williams, R., Belo, J.B., Lidia, J., Soares, S., Ribeiro, D., Moreira, C.L., Almeida, L., Barton, L. & Erskine, W. Productivity gains in vegetables from rice husk biochar application in nutrient-poor soils in Timor-Leste. *Sci Rep* 13, 10858 (2023). DOI: <https://doi.org/10.1038/s41598-023-38072-2>
  30. Woolf, D., Lehmann, J., Cowie, A., Cayuela, M. L., Whitman, T. & Sohi, S. (2018). Biochar for climate change mitigation: navigating from science to evidence-based policy. In R. Lal, & B. A. Stewart (Eds.), *Soil and Climate CRC Press*. DOI: <https://www.taylorfrancis.com/books/e/9780429487262>
  31. Yaashikaa, P.R., Kumar, P.S., Varjani, S. & Saravanam, A. (2020). A critical review on the biochar production techniques, characterization, stability and applications for circular bioeconomy. *Biotechnology reports (Amsterdam, Netherlands)*, 28, e00570. DOI: [10.1016/j.btre.2020.e00570](https://doi.org/10.1016/j.btre.2020.e00570)
  32. Zhao, F., Zou, G., Shan, Y., Ding, Z., Dai, M. & He, Z. Coconut shell derived biochar to enhance water spinach (*Ipomoea aquatica* Forsk) growth and decrease nitrogen loss under tropical conditions. *Sci Rep* 9, 20291 (2019). DOI: <https://doi.org/10.1038/s41598-019-56663-w>

UNDER PEER REVIEW

