

## Original Research Article

### Structure and Floristic Composition of Existing Agroforestry Systems in Allahabad District of Uttar Pradesh, India

Comment [A1]: remove

#### Abstract

The Structure and Floristic Composition of Existing Agroforestry Systems was carried out in the farmers field of village Kashari (site-I) and Korihar (site-II) at Allahabad District of Uttar Pradesh, India during 2008-2010. The quantitative characteristics of vegetation were determined by the quadrat method. The floristic diversity study was made by adopting random sampling approach, under which, 20 quadrates of 10 m x 10 m were laid down. The density, frequency, abundance, IVI, Species –diversity, concentration of dominance, species-richness, equitability and beta diversity and other useful parameters for comparison of different types of existing agroforestry systems were analyzed by the simple arithmetic calculation. 20 quadrates of 10 x 10 m size were laid out in each site for the observation of trees and shrubs. The vegetative structure of trees and shrub in each four sites was estimated for frequency; density and abundance by using expressions given by Curtis and Mc Intosh, 1950. Floristic diversity studies of the herbaceous species were not attempted.

Comment [A2]: remove

Comment [A3]: the quadrats

Comment [A4]: remove and put the conclusion from the study

#### **Introduction:**

India has a long existing of agroforestry (AF) systems and several indigenous agroforestry systems, based on people's needs and site-specific characteristics have been developed over the years. The existing agroforestry systems in India include trees on farms, community forestry and a variety of local forest management and ethno forestry practices (Gairola et al 2008). In India, the practice of growing scattered trees on farmlands is quite old and has not changed much over centuries; these trees are multipurpose, used for shade, fodder, fuel wood, fruit, vegetables and medicinal uses (Bijalwan, et al., 2011, Dey et al, 2022). The landuse pattern is one of the important aspects for analyzing the structure, composition and phyto diversity for understanding the vegetation dynamics of the existing sites (Umrao et al, 2017). The farmers have adopted the trend of growing trees around agricultural land due to significant economics as well as social benefits. But it will be worthwhile to work out an integrated approach with the help of agricultural and forestry scientists, depending on the suitability of crops and locations (Ranjan et al., 2016, Chaurasia et al, 2022). A new land-use options that increase livelihood security and reduce vulnerability to climate and environmental change are necessary (Pathak et al., 2016, Mishra et al, 2022). Community structure, composition and vegetative function are the most important ecological attributes of forests, which show variations in response to environmental as well as anthropogenic variables (Gairola et al., 2008; Timilsina et al., 2007). Species diversity is an important concept and one of the major attributes of a natural community (Chaurasia et al, 2020). These variations in vegetation structure, richness, diversity and distribution are directly correlated with the intensity of variables like geographical location, agricultural practices and their extent (Criddle et al., 2003). In Central and Eastern Uttar Pradesh, the agroforestry systems are well established. As multiple cropping is a kind of natural resource management based on environmental, social and economic criteria which is done by combination of trees, pasture

and cropland, in order to increase social, economic and environmental benefits (Rawat et al, 2009). Planting and harvesting of wood products, fruits, roots, leaves, fuel, and fodder along with agricultural crops on the same piece of land has been practiced since the old days. However, under present conditions, the agroforestry techniques applied by the farmers appear to be poorly developed and exploitative. In most cases, the trees are neither protected nor properly managed. Hence, there is immense potential for improvement of traditional agroforestry systems in order to realize the real production potential of existing agroforestry systems. Greater agrobiodiversity also may ensure longer term stability of C storage in fluctuating environments (Henry *et al.*, 2009), apart from augmenting biomass production potential (Kumar, 2006). Main aimed of this paper are to describe the structural attributes of density, frequency, diversity, equitability and species richness on the existing agroforestry systems in Fatehpur and Allahabad districts of Uttar Pradesh. These studies helps in determination of predominating communities of timber trees, horticultural trees and shrubs species and further identify the localities having protection and promotion of these plants.

Comment [A5]: was

#### MATERIAL AND METHODS:

The study was carried out in the two selected village sites in Allahabad in eastern part of UP. The two sites namely Kashari (site-I) and Korihar (site-II) were selected in Allahabad district, these sites are situated on the right side of the Ganga river at a distance of about 20 km and 35 km away from Allahabad city (81°50' E longitude and 25° 27' N latitude).



Map.1 - Map of study areas

Comment [A6]: the

**Floristic -Diversity:** The floristic diversity study was made by adopting random sampling approach, under which, 20 quadrates of 10 m x 10 m and 5 m x 5 m size were laid out in each sites for the observation of trees and shrubs respectively. The status of density, frequency, Abundance,IVI, Species –diversity, concentration of dominance, species-richness, Equitability and Beta diversity and other useful parameters for comparison of different types of existing agroforestry systems were analyzed by the simple arithmetic calculation. The quantitative characteristics of vegetation were determined by the quadrat method. The vegetative structure of trees and shrub in each two sites was estimated for frequency; density

and abundance by using expressions given by Curtis and Mc Intosh, 1950. The floristic diversity study was made by adopting random sampling approach, under which, 20 quadrates of 10 x 10 m size were laid out in each sites for the observation of trees and shrubs. The vegetative structure of trees and shrub in each four sites was estimated for frequency; density and abundance by using expressions given by Curtis and Mc Intosh, 1950. Floristic diversity studies of the herbaceous species were not attempted. Basal area of trees was calculated as cross sectional area of stem at DBH (1.37 m), while basal area of shrubs was calculated as cross sectional area of main stem at 15 cm above from the ground level. Basal Area =  $\pi (d^2/4)$ , Where d is the diameter of tree

Comment [A7]: remove

Comment [A8]: remove

Number of sampling units in which species occurred

$$\text{Frequency} = \frac{\text{Number of sampling units in which species occurred}}{\text{Total number of sampling unit studies}} \times 100$$

Total number of sampling unit studies

Total number of individual of species

$$\text{Density} = \frac{\text{Total number of individual of species}}{\text{Total number of quadrat studied}}$$

Total number of quadrat studied

Total number of individual of the species in all sampling units

$$\text{Abundance} = \frac{\text{Number of sampling unit in which the species occurred}}{\text{Total number of individual of the species in all sampling units}}$$

Number of sampling unit in which the species occurred

Total basal area of trees

$$\text{Basal area per tree} = \frac{\text{Total basal area of trees}}{\text{Number of trees}}$$

Number of trees

**Relative Basal Area:** The relative density, relative frequency, relative basal areas were calculated using following formula.

Density of individual of species

$$\text{Relative density} = \frac{\text{Density of individual of species}}{\text{Total density of all species}} \times 100$$

Total density of all species

Frequency of the individual species

$$\text{Relative frequency} = \frac{\text{Frequency of the individual species}}{\text{Total frequency of all the species}} \times 100$$

Total frequency of all the species

Basal area of the individual of species

$$\text{Relative basal area} = \frac{\text{Basal area of the individual of species}}{\text{Total basal area of all the species}} \times 100$$

**Importance Value Index:** The importance value index (IVI), which is an integrated measure of the relative frequency, relative density and relative basal area/dominance, was calculated for each tree species given by Curtis, 1959.

$$\text{Importance Value Index (IVI)} = \text{RD} + \text{RF} + \text{RBA}$$

The number of trees falling in the sample unit was counted and classified as per their diameter and height characteristics. The Species diversity (Shannon index), concentration of dominance (Simpson index) and other useful parameters for comparison of different types of existing agroforestry systems were calculated (Shannon and Weaver, 1963 and Simpson, 1949).

**Tree diversity analysis:**

Tree diversity in all four sites of agroforestry systems (trees and shrubs) were calculated by the following diversity indices.

- (a) **Species Diversity Index.** It was calculated by the formula given by Margalef, 1958.

$$H = -\sum [(ni/N) \log (ni/N)]$$

Where  $n_i$  was the total number of individuals of species  $N$  was the total number of individuals of all the species on that site.

- (b) **Concentration of dominance** was measured by the formula of Simpson Index developed by Simpson, 1949.

$$Cd = -\sum \left[ \left( \frac{n_1}{N} \right)^2 + \left( \frac{n_2}{N} \right)^2 + \dots + \left( \frac{n_n}{N} \right)^2 \right]$$

Where  $N$  was the total number of individuals of species  $n_i$  was the total number of individuals of all the species on that site.

- (c) **Equitability (e)** was calculated as suggested by Pielou (1975) as  $e = H / \ln s$

Where  $H$  is the Shannon – Wiener Index and  $s$  = total number of species

- (d) **Species richness** was calculated by the following equation of Margalef (1958)  $d = s - 1 / \ln N$

Where  $s$  = number of species, and  $N$  = number of individuals of all species

(e) **Beta diversity** was calculated as outlined by Whittaker, 1977

$$bd = S_c / s$$

Where  $S_c$  = total number of species in all sites and average species per site.

## RESULTS AND DISCUSSION

### Floristic-diversity analysis and distribution patterns of tree species:

The distribution patterns and species composition of existing agroforestry systems viz. agrisilvicultural and agrihorticultural system commonly practiced in Allahabad was studied.

**Floristic diversity at Site –I (Kashari) of Allahabad district:** The results of floristic-diversity at site-I are presented in table 1 and graphically illustrated in figure 1. It is evident from the data that among tree species available in site –I timber and fuelwood, horticultural and shrubs species were 18, 8 and 4, respectively. The dominant and co-dominant species were *Acacia nilotica* and *Azadirachta indica* showed IVI values of 24.54 and 24.09, respectively. The highest (70 trees ha<sup>-1</sup>) value of density was recorded for *Acacia nilotica* in timber and fuel wood tree species followed by *Azadirachta indica* (65 trees ha<sup>-1</sup>). The highest tree density was recorded for *Emblica officinalis* (70 trees ha<sup>-1</sup>) among horticultural tree species followed by *Mangifera indica* (35 trees/ha). Total basal cover for tree was recorded higher for *Ficus religiosa* (5.144 m<sup>2</sup>/ha) followed by *Madhucalatifolia* (5.1094 m<sup>2</sup>/ha). Among horticulture and shrubs species, the dominant and co-dominant species were *Emblica officinalis* and *Mangifera indica* with IVI values of 18.89 and 16.207, respectively.

**Table 1: Floristic- diversity of site-I (Kashari)of Allahabad district**

**Timber and fuelwood trees**

Name of the species	Density (100m <sup>2</sup> )	Frequency	Abundance	BA (cm <sup>2</sup> )/100m <sup>2</sup>	Relative Density	Relative Frequency	Relative Basal Area(RBA)	IVI	B.A. M <sup>2</sup> /ha.
<i>Acacia nilotica</i> L.Willd.ex del.	0.70	55	1.27	393.56	8.04598	8.02920	8.46728	24.54246	3.9356
<i>Aegle marmelos</i> (L.)Corr.	0.25	20	1.25	151.81	2.87356	2.91971	3.26609	9.05936	1.5181
<i>Artocarpusheterophyllum</i> L.	0.25	15	1.67	146.61	2.87356	2.18978	3.15428	8.21762	1.4661
<i>Azadirachta indica</i> L.	0.65	60	1.08	365.45	7.47126	8.75912	7.86248	24.09287	3.6545
<i>Dalbergiasissoo Roxb.</i>	0.20	15	1.33	138.80	2.29885	2.18978	2.98625	7.47488	1.388
<i>Eucalyptus tereticornis</i> Sm.	0.30	25	1.20	143.23	3.44828	3.64964	3.08144	10.17935	1.4323
<i>Ficus benghalensis</i> L.	0.25	20	1.25	373.92	2.87356	2.91971	8.04468	13.83795	3.7392
<i>Ficus glomerata</i> L.	0.30	25	1.20	179.69	3.44828	3.64964	3.86592	10.96383	1.7969
<i>Ficus religiosa</i> L.	0.35	20	1.75	514.40	4.02299	2.91971	11.06723	18.00992	5.144
<i>Limonia acidissima</i> L.	0.25	25	1.00	162.47	2.87356	3.64964	3.49556	10.01875	1.6247
<i>Madhucalatifolia</i> Macb.	0.40	30	1.33	510.94	4.59770	4.37956	10.99258	19.96984	5.1094
<i>Pongamia pinnata</i> L.	0.20	10	2.00	187.04	2.29885	1.45985	4.02408	7.78279	1.8704
<i>Populus deltoides</i> Bartr.ex.Marsh	0.20	15	1.33	87.226	2.29885	2.18978	1.87663	6.36526	0.8723
<i>Prosopis juliflora</i> (sw) DC.	0.25	15	1.67	34.851	2.87356	2.18978	0.74980	5.81315	0.3485
<i>Syzygium cumini</i> (L.)Skeel	0.20	10	2.00	187.26	2.29885	1.45985	4.02875	7.78745	1.8726
<i>Tamarindus indica</i> L.	0.20	15	1.33	224.53	2.29885	2.18978	4.83072	9.31935	2.2453

<i>Tectonagrandis</i> L inn.f.	0.60	50	1.20	194.27	6.89655	7.29927	4.17967	18.37549	1.9427
<i>Zizyphusmaritima</i> Lam	0.25	25	1.00	129.84	2.87356	3.64964	2.79346	9.31666	1.2984
<b>Horticultural trees</b>									
<i>Carica papaya</i> L.	0.20	10	2.00	12.031	2.29885	1.45985	0.25885	4.01755	0.1203
<i>Carissa corandus</i> L.	0.30	25	1.20	9.9589	3.44828	3.64964	0.21426	7.31217	0.0996
<i>Citrus lemon</i> (L)Burm.f	0.30	30	1.00	10.424	3.44828	4.37956	0.22427	8.05210	0.1042
<i>Cordia myxa</i> Roxb.	0.15	10	1.50	3.5652	1.72414	1.45985	0.07670	3.26070	0.0357
<i>Emblica officinalis</i> Gaertn	0.70	60	1.17	96.851	8.04598	8.75912	2.08372	18.88882	0.9685
<i>Mangifera</i> indica L.	0.35	30	1.17	362.76	4.02299	4.37956	7.80474	16.20729	3.6276
<i>Musa paradesiaca</i> L.	0.15	10	1.50	2.3339	1.72414	1.45985	0.05021	3.23420	0.0233
<i>Psidium</i> guajavaL .	0.15	10	1.50	21.959	1.72414	1.45985	0.47245	3.65644	0.2196
<b>Shrubs species</b>									
<i>Bougainvillaglab</i> raL.	0.20	15	1.33	0.8458	2.29885	2.18978	0.01820	4.50683	0.0085
<i>Jatropha</i> curcas L.	0.15	10	1.50	0.6453	1.72414	1.45985	0.01388	3.19788	0.0065
<i>Ricinus</i> communis L.	0.15	15	1.00	0.7074	1.72414	2.18978	0.01522	3.92914	0.0071
<i>Ziziphus</i> zizyphus L.	0.10	10	1.00	0.0332	1.14943	1.45985	0.00071	2.60999	0.0003
<b>Total</b>	<b>8.70</b>	<b>685</b>	<b>40.74</b>	<b>4648</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>300.0</b>	<b>46.48</b>




Fig. 1: Floristic diversity of site-I (Kashari) of Allahabad district


## 2. Floristic- diversity analysis at site –II (Korihar) of Allahabad district:

Data on Floristic diversity findings of tree layer at site-IV are presented in table 2 and graphically illustrated in figure 2 revealed that total number of timber and fuel wood, horticultural and shrubs species were 18, 6 and 4, respectively. The dominant and co-dominant species were *Tectonagrandis* and *Acacia nilotica* and showed IVI values of 23.24 and 23.0, respectively. The highest density value was recorded for *Tectonagrandis* (65 trees ha<sup>-1</sup>) followed by *Acacia nilotica* (60 trees ha<sup>-1</sup>). In horticulture and shrub species, the dominant and co-dominant species were *Mangifera indica* and *Emblia officinalis* with their IVI values of 18.05 and 18.01, respectively. The highest value of tree density was recorded for *Emblia officinalis* (60 trees ha<sup>-1</sup>) followed by *Madhuca latifolia* (30 trees ha<sup>-1</sup>). Total tree basal cover was observed comparatively higher in *Madhuca latifolia* (4.53 m<sup>2</sup> ha<sup>-1</sup>) followed by *Azadirachta indica* (3.63 m<sup>2</sup> ha<sup>-1</sup>). In horticultural and shrubs species, basal cover was comparatively greater in *Mangifera indica* (4.27 m<sup>2</sup> ha<sup>-1</sup>) and *Emblia officinalis* (0.830 m<sup>2</sup> ha<sup>-1</sup>). The results also conform to the earlier findings of Bijalwan et al. 2011.

**Table 2 Floristic -diversity of site-II (Korihar) of Allahabad district**

Name of the species	Density (100m <sup>2</sup> )	Frequency	Abundance	BA (cm <sup>2</sup> )/ 100m <sup>2</sup>	Relative Density	Relative Frequency	Relative Basal Area(RB A)	IVI	B.A. M <sup>2</sup> /ha.
<b>Timber and fuelwood trees</b>									
<i>Acacia nilotica</i> L. Willd.ex del.	0.60	50	1.20	281.36	7.69231	8.19672	7.19095	23.0799	2.8136
<i>Aegle marmelos</i> (L.)Corr.	0.25	20	1.25	137.94	3.20513	3.27869	3.52549	10.0093	1.3794
<i>Albizia procera</i> L.	0.15	10	1.50	62.999	1.92308	1.63934	1.61011	5.17253	0.63
<i>Artocarpus heterophyllus</i> L.	0.25	20	1.25	127.23	3.20513	3.27869	3.25165	9.73547	1.2723
<i>Azadirachta indica</i> L.	0.50	45	1.11	363.06	6.41026	7.37705	9.27909	23.0663	3.6306
<i>Dalbergia sissoo</i> Roxb.	0.25	15	1.67	219.78	3.20513	2.45902	5.61721	11.2813	2.1978
<i>Eucalyptus tereticornis</i> Sm.	0.25	20	1.25	123.95	3.20513	3.27869	3.16787	9.65169	1.2395
<i>Ficus benghalensis</i> L.	0.20	20	1.00	281.84	2.56410	3.27869	7.20319	13.0459	2.8184
<i>Ficus glomerata</i> L.	0.30	25	1.20	189.05	3.84615	4.09836	4.83165	12.7761	1.8905
<i>Ficus religiosa</i> L.	0.20	20	1.00	268.56	2.56410	3.27869	6.86371	12.7065	2.6856
<i>Madhucalatifolia</i> Macb.	0.40	30	1.33	453.11	5.12821	4.91803	11.58055	21.6267	4.5311
<i>Pithecellobium dulce</i> (Roxb.)Benth.	0.20	15	1.33	152.19	2.56410	2.45902	3.88953	8.91265	1.5219
<i>Pongamia pinnata</i> L.	0.30	25	1.20	129.73	3.84615	4.09836	3.31562	11.2601	1.2973
<i>Populus deltoids</i> <i>Bartrx.ex.Marsh</i>	0.20	10	2.00	41.854	2.56410	1.63934	1.06969	5.27314	0.4185
<i>Prosopis juliflora</i> (sw) DC.	0.30	15	2.00	41.821	3.84615	2.45902	1.06885	7.37402	0.4182
<i>Syzygium cumini</i> (L.)Skeel	0.20	10	2.00	143.61	2.56410	1.63934	3.67024	7.87369	1.4361
<i>Tectona grandis</i> Linn.f.	0.65	55	1.18	230.62	8.33333	9.01639	5.89412	23.2438	2.3062
<i>Zizyphus maritima</i> Lam	0.35	25	1.40	102.33	4.48718	4.09836	2.61529	11.200	1.0233
<b>Horticultural trees</b>									
<i>Carica papaya</i> L.	0.20	10	2.00	12.031	2.56410	1.63934	0.30749	4.51094	0.1203
<i>Carissa corandus</i> L.	0.30	25	1.20	9.9589	3.84615	4.09836	0.25453	8.19904	0.0996

<i>Emblca officinalis</i> Gaertn	0.60	50	1.20	83.015	7.69231	8.19672	2.12169	18.0107	0.8302
<i>Mangifera indica</i> L.	0.30	20	1.50	427.76	3.84615	3.27869	10.93273	18.0575	4.2776
<i>Musa paradisiaca</i> L.	0.15	10	1.50	2.3339	1.92308	1.63934	0.05965	3.62207	0.0233
<i>Psidium guajava</i> L.	0.15	10	1.50	21.321	1.92308	1.63934	0.54491	4.10733	0.2132
<b>Shrubs species</b>									
<i>Bougainvillea glabra</i> L.	0.20	15	1.33	0.8458	2.56410	2.45902	0.02162	5.04474	0.0085
<i>Jatropha curcas</i> L.	0.15	15	1.00	3.7623	1.92308	2.45902	0.09616	4.47825	0.0376
<i>Ricinus communis</i> L.	0.15	15	1.00	0.7074	1.92308	2.45902	0.01808	4.40017	0.0071
<i>Ziziphus zizyphus</i> L.	0.05	10	0.50	0.0166	0.64103	1.63934	0.00042	2.28079	0.0002
<b>Total</b>	<b>7.80</b>	<b>610</b>	<b>37.6</b>	<b>3912.8</b>	<b>100.0</b>	<b>100.0</b>	<b>100.0</b>	<b>300.0</b>	<b>39.128</b>



**Fig.2 Floristic diversity of site-II(Korihar)of Allahabad district**

**3. Diversity index analysis:** Diversity index analysis of vegetation of both sites of the district Allahabad are presented in Table 3. The value of index of dominance (Simpson index) was higher in agrisilviculture system at site –II (0.038) followed by site-I (0.035). The species diversity (sd) was highest at site –II (0.999) In agrihorticultural system highest species of diversity was recorded at site-I (0.330) and lowest at site-II (0.308). Equitability (e) of trees were found maximum (0.055) in agrisilviculture system at site-I, whereas minimum was recorded at site-II (0.038) in agrihorticulture system. Highest Species richness was found in agrisilviculture system in site –II (0.153) followed by site –I (0.152) in agrihorticulture system where as in lowest found in agrisilviculture system in site -I(0.146). Beta diversity were found highest in agrihorticulture system in site –II (5.167) and lowest at site –I (3.875) in agrihorticulture system. The results of diversity index are also supported by the findings of Knight, 1975. In case of modified Simpson's Index Value almost similar observation was reported by Jose *et.al.*, (1994). Basha (1987) has also reported Simpson's index of diversity of 0.94 for evergreen forests of Silent Valley. The Shannon-Wiener Index Value (H') showed that the diversity was almost similar to that in a tropical forest (5.45). H' value in tropical rain forest generally varies between 5.06 in a young stand to 5.4 in an old stand (Knight, 1975).

**Table 3: Diversity Index of study sites-I and II of Allahabad districts:**

AF system	Species – Diversity Index (Shannon Index)		Simpson Index (Concentration of Dominance)		Species Richness		Equitability		Beta Diversity	
	Trees	Shrubs	Trees	Shrubs	Trees	Shrubs	Trees	Shrubs	Trees	Shrubs
AS/S <sub>I</sub>	0.970	0.060	0.035	0.0002	0.146	0.250	0.053	0.020	3.944	4.667
AS/S <sub>II</sub>	0.999	0.064	0.038	0.0002	0.153	0.272	0.055	0.021	3.944	4.667
AH/S <sub>I</sub>	0.308	0.027	0.007	0.0002	0.152	0000	0.038	0.027	3.875	6.000
AH/S <sub>II</sub>	0.262	0.028	0.008	0.0002	0.147	0000	0.043	0.028	5.167	6.000

**Comment [A9]:** AS MEANS.....  
AH MEANS.....

### CONCLUSIONS:

In view of the above findings, it is concluded that the IVI was recorded maximum in *Acacia nilotica* in both the research sites. As far as concerned to maximum value of Concentration of Dominance (Simpson Index), Equitability, Beta diversity and Species diversity, Site-I (Kashari) is superior while species richness was found at site –II (Korihar).

**Comment [A10]:** RE Frame your conclusion pls

### REFERENCES

- Bhandari, B.S. and Tiwari, S.C.** (1997). Dominance and diversity along an altitudinal gradient in a montane forest of Garhwal Himalaya. Proc. Indian National Science Academy 64:437-446.
- Bijalwan, A., Sharma, C.M. and Kediya, V.K.** (2011). Socioeconomic status and livelihood support through traditional agroforestry systems in hill and mountain agroecosystems of Garhwal Himalaya, India. *Indian Forester*, 137(12) 1423-1431.
- Bijalwan, A., Swami, S.L., Sharma, C.M., Umrao, R. and Paliwal, H.B.** (2011). Structure, Composition and diversity of tree vegetation in Sal mixed dry tropical forest in Chhatisgarh plains of India. *Indian Forester*, 137(4):453-462.
- Buvaneshwaran, C.; George, M.; Manivachakam, P.; Subramanian, V., (2001).** Comparative studies on the performance and productivity of teak in farmland and in forest plantation. *Rangemanagement and Agroforestry*, 22 (1):113-117.
- Chandra, S. Negi and Nautiyal S. (2005).** Phyto-sociological studies of a traditional reserve forest-ThalkeDharPithoragarh, Central Himalayas (India).
- Chaurasia, S. and Mishra, K. (2022).** Phytosociological Study of Kushmi Forest Division of Gorakhpur Uttar Pradesh, Int. J. Agriworld, 3 [2]: 20-33.
- Chaurasia, S., Hemant Kumar, Kumud Dubey and Y.K. Agarwal (2020).** Assessment of growing stock, above ground biomass and sequestered carbon in sal dominated kushmi forest Gorakhpur, *Indian Forester*, 146(11): 1041-1045.
- Dwivedi, A.P. (1993).** Babul (*Acacia nilotica*): A multipurpose tree of dry areas. Scientific Publisher, Jodhpur, India. Pp 69.

- Dey, A., Hemant Kumar and Sangjukta Das** (2022).Significance of Non-Timber Forest Products (NTFPs) to the Tribal Wealth of Ranchi District, Jharkhand, *Eco.Env.& Cons.* 28 (4) :2014-2020
- Gairola, S., Rawal, R.S., and Todaria, N.P.** (2008).Forest vegetation patterns along an altitudinal gradient in sub-alpine zone of west Himalaya, India.*African Journal of Plant Science*, 2(6):42-48.
- Hussain,M.S.,Sultana,A.,Khan,J.A.andKhan ,A.(2008).**Species composition and community structure of forest stands in KumaonHimalaya,Uttarakhand,India.*Tropical Ecology*,49(2):167-181.
- Jose,S.Sreepathy, A., Kumar,B. and Venugopal, V.K.**(1994).Structural, Floristic and edaphic attributes of the grassland-Shola forests of Eravikulam in Penunsular India.**For.Eco.Management**.65(2-3):279-291.
- Kharkwal,G.andRawat,Y.S.**(2010).Structure and composition of vegetation in subtropical forest of KumaunHimalaya.*African Journal of Plant Science*,4(4):116-121.
- Knight,D.H.**(1975).A phytosociological analysis of species rich tropical forests on Barro Colorado Island,Panama.*Ecol.Monogr*,45:259-289.
- Kumar and Bhatt.**(2006).Plant Biodiversity and Conservation of Forests in Foot Hills of GarhwalHimalaya.*Lyonia a Journal of Ecology and Application*.Des.1-19.
- Kumar, B.M; Suman Jacob George and Chiimani, S.** (1994).Diversity, structure and standing stock of wood in the homegardens of Kerala in peninsular India.*Agroforestry Systems*, 25:243-262.
- Malik, N.Z.,Arshad.M.andMirza,S. N.**(2007).Phytosociological attributes of different plant communities of PirChinasi Hills of Azad Jammu and Kashmir. *International Journal of Agriculture and Biology*, 9(4):569-574.
- Marglef, D.R.**(1958).*Information Theory in Ecology*. Year book of the society for *General Systems Research*, 3:36-71.
- Menon, A. R.R. and Suraj, M.A.** (2009).Phytosociological Analysis of woody vegetation along an altitudinal gradient in Ponnudi Hill, Thrissur District, Kerala, South India.*Indian Forester*.vol.135(6):799-806.
- Minj,A.V.andQuli ,S.M.S.**(2000).Impact of Agroforestry on socio-economic status of respondents. *Indian Forester*, 126(7):788-791.
- Mishra, R.** (1968). *Ecology work book*. Oxford and IBH publishing co, Calcutta.p.244.
- Mishra, K., Kumar, H., N.N Mishra and BhagyashreeKesharwani** (2022). Tree species composition, biomass and carbon stocks in Urban Green Spaces of Prayagraj,*Eco.Env. & Cons.* 28 (4) :1912-1919
- Mughal,A.H.Ara,T. and Bhattacharya,P.** (2000).Socio-Economic aspects of Agroforestry in rural Srinagar of Kashmir valley. *Indian Forester*, 126 (3) 234-240.
- Negi, B.S., Chauhan, D.S. and Todaria, N.P.** (2008) .Comparative plant diversity between panchayat and adjoining reserve forests in GarhwalHimalaya.*Indian Journal of Forestry*, 31(4):585-593.
- Negi, C.S.** (2003).Role of Traditional Knowledge and Beliefs in Conservation-case Studies from Central Himalaya, India.*Man in India*, 83(3&4):371-391.
- Pandey, D. N.** (2007).Multifunctional agroforestry systems in India.*Current Science*, Vol.92 (4).
- Pandey,U. and Singh, J.S.** (1984).Energetics of Hill agro-ecosystem: A case study from Central Himalayas. *Agricultural Systems*, 13: 83-85.
- Pane, V.G. and Sukhatme, P.V.**(1978). *Statistical Methods for Agricultural Workers*. ICAR

Publication, New Delhi.

- Pielou, E.A.** (1975). *Ecological Diversity*. John Wiley and Sons. New York, pp. 165.
- Rajvanshi, R. V. Kumar, W. Bachpai, Rajgopal and Raj, S.F.H.** (1987). Herbaceous under growth in some forest habitat in Nilgiris. *Indian Forester*, 113(9):599-608.
- Ralhan, P.K., A.K. Saxena and Singh, J.J.** (1982). Analysis of forest vegetation at and around Nainital in Kumaon Himalaya. *Proc. Indian National Science Academy*, 48:121-137.
- Ranganathan, C.R.** (1938). Studies in the ecology of the Shola grassland vegetation of the Nilgiri plateau. *Indian Forester*, 64:523-541.
- Rawat, V.S. and Chandhok, A.** (2009). Phytosociological analysis and distribution patterns of tree species: A case study from Govind Pashu Vihar, National Park, Uttarakhand. *New York Science Journal*, 2(4):58-63.
- Saravanan, S., Ravichandran, K., Balsubramanian, A. and Paneerselvam, K.** (2013). Structure and floristic composition of tree diversity in Andaman tropical evergreen forest of middle Andaman, India. *Indian Journal of Forestry*, Vol. 36 (2): 167- 171.
- Saxena, A.K. and Singh J.S.** (1982). A phytosociological analysis of woody species in forest communities of a part of Kumaon Himalaya. *Vegitatio.*, 50:3-32.
- Semwal, R.L., Nautiyal, S., Rao, K.S., Maikhuri, R.K. and Bhandari, B.S.** (1999). Structure of forests under community conservation: A preliminary study of Jardhar village initiative in Garhwal Himalaya. *ENVIS bulletin-Himalayan Ecology and Development*, 7(2):16-27.
- Shannon, C. and Weaver, W.** (1963). *The mathematical theory of communication*. University of Illinois press, Urbana, U.S.A.. pp. 117.
- Singh, B., Sharma, K.N. and Kumar, D.** (2011). Influence of tree species on improvement of soil fertility status in central zone of Punjab. *Indian Journal of Agriforestry*. 13(1):35-39.
- Singh, and Beniwal, S.K.** (1995). Socio-economic development through community Forest. Natl. Seminar on com. Forestry: Biodiversity of Forest species, May 2-3, Abs. & Souvenir, ISTS, Solan and M.S. Swaminathan Research Foundation, pp. 88.
- Umrao, R., B. Mehera, N. Khare and Hemant Kumar** (2017) Structure and floristic composition of existing agroforestry systems in Fatehpur district of U.P., *Current World Environment*, 12(1), 124-131.
- Verma, R.K., Totey, N.G. and Gupta, B.N.** (1997). Analysis of forest vegetation in the permanent preservation plot of Tamna in Orissa. *Indian Forester*, 123(11):1007-1016.