

Impact of Nano DAP and Zn EDTA on cowpea growth and yield

Comment [h1]: The first letter of each word should be capital in title.

ABSTRACT

Pulses are gaining more important position in Indian agriculture. But the average productivity of pulses ~~is pulses~~ in India (764 kg ha^{-1}) is much less productive than the global average (848 kg ha^{-1}). ~~An increased yield of cowpeas can be achieved by Basally applied nutrients are lost over time, thus~~ applying important fertilizer to ~~cowpeas~~ during their critical growth ~~as with the passage of time basally applied nutrients are lost, is a effective strategy to enhance their yield.~~ Especially smart fertilizers like Nano DAP and Zn EDTA are very helpful in that. Therefore, the field experiment was done in the dairy Farm, School of Agriculture and Animal Science, Gandhigram Rural Institute, Gandhigram, Tamil Nadu during January 2024 to March 2024 to assess the impact of Nano DAP and Zn EDTA on growth and yield of cowpea. It was laid out in Randomized Block Design (RBD) with seven treatments ~~with under~~ three replications. Among the different treatments, **RDF 50% P, 100% NK + Seed treatment** and two foliar sprays of 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T_7) ~~was achieved a~~ supremely improved growth parameters like plant height, no. of branches per plant, leaf area index, dry matter production, and yield attributes viz., pod length, no. of pods per plant, no. of seeds per pod and test weight. Further it recorded a higher seed yield, haulm yield, ~~harvest and harvest~~ index. The ~~minimum value was recorded under~~ control treatment (T_1) ~~was recorded the minimal value.~~ Experimental results clearly revealed that RDF 50% P, 100% NK + Seed treatment and two foliar sprays of 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T_7) has recorded the highest grain yield of 1720 kg ha^{-1} .

Comment [h2]: Starting of this sentence lacks the flow and link to the previous sentence.

Comment [h3]: At first the full words should be written along with abbreviations in bracket, later on the abbreviation can be used.

Comment [h4]: Full forms would be encouraged.

Key words: Nano DAP, Zn EDTA, Cowpea, Yield.

Introduction

Cowpea is well-known for its ability to withstand dryness. It is also known as black-eyed pea or southern pea. The shadowing effect of its broad, drooping leaves preserves soil moisture. It can be used as a vegetable, food, feed, fodder, forage and green manuring. **In the semi-arid tropical regions of Asia, Africa, Southern Europe, Central America, and South America, it is a**

vital food legume. Worldwide, the use of cowpeas as food, fodder, and a raw resource for numerous industries is increasing. Cowpea, also referred to as vegetable meat, is a valuable source of vitamins, minerals, and protein for a healthy diet. 4.3 g protein, 2.0 fibre, 8.0 g carbohydrate, 74 mg phosphorus, 2.5 mg iron, 13.0 mg vitamin C, and 0.9 mg minerals, etc. (~~Damoar et al., 2020~~) are all provided presently in 100 g of green tender pods (~~Damoar et al., 2020~~). India has low pulse production and productivity as pulses are cultivated mainly on marginal and submarginal soils with inadequate management techniques. India is the world's biggest importer and user of pulses. With 2,745.28 tonnes production, Uttar Pradesh is the top cowpea producer (NHB, 2021-2022). India's productivity of pulses (764 kg ha⁻¹) is significantly lower than the global average (848 kg ha⁻¹). Overall area cultivated for pulses remains almost unchanged at 22 to 24 million hectares, with a production that has remained relatively constant at 12 to 14 million tonnes. In contrast to the FAO/WHO recommendation of 80 g day⁻¹, the current per capita availability of pulses has reduced from 51.1 g day⁻¹ in 1970-71 to less than 41.9 g day⁻¹ (2017). It results in an alarming shortage of pulses in India, worsening the issue of malnourishment among a significant percentage of the vegetarian population.

Thus, there is an urgent need to increase the pulses production in order to fulfill the demand by adopting the appropriate production technologies. To improve its production and productivity, nutrient management strategies like foliar application of important nutrients is a feasible one. The cowpea crop responds well to the fertilizer. Today, foliar application of fertilizer is gaining momentum because photosynthetic efficiency is increased through foliar application of fertilizer by delaying the leaf senescence. Legumes are crops that are fond of phosphorus, and ~~is~~ are necessary for the development of seeds, nodules, and flower initiation (Haruna and Aliyu, 2011). (~~Ndakemi~~ and Dakora, 2007). Zinc is a major micronutrient that is crucial to the growth and development of plants. In developing countries, Zn deficiency occurs due to inadequate supply of micronutrient. Several research findings also reported that, considerable increment in the yield of the crop is due to the application of Zn fertilizers. Smart fertilisers, such as Zn EDTA and Nano DAP, will be key in boosting agricultural productivity in sustainable manner by ~~enhance~~ enhancing the nutrient use ~~efficienece~~ efficiency. Keep the above points in view that, ~~the~~ this study was carried out to analyse the effect of Nano DAP and Zn EDTA on growth and yield of Cowpea.

Materials and methods

Comment [h5]: Reference should be present.

Comment [h6]: Reference required.

Comment [h7]: The most recent report's reference should be added.

Comment [h8]: After full stop first letter of first word should be capital.

Comment [h9]: Reference should be added.

Comment [h10]: Reference?

Comment [h11]: Write correct spelling of this name

Comment [h12]: Reference should be added.

Comment [h13]: Seed sowing detail should be mentioned, such as the spacing of seeds? The source from where the seeds were taken should be mentioned. This section can be improved more.

Comment [h14]: The recorded parameters must be mentioned, their data recording methodology can be mentioned. Details of location can be added like temperature.

The field experiment was conducted at Gandhigram Rural Institute, Gandhigram during 2024 (January- April) to find out the effect of Nano DAP and Zn EDTA on growth and yield attributes of Cowpea. The treatment consists of seven treatments ~~which are in~~ including the combination ~~with of~~ Zn EDTA and Nano DAP. Plot size ~~were was~~ 5×4 m. The details of the treatment ~~were are~~, T₁ – Control, T₂ – RDF 100% NPK (N- 12.5kg/ha, P₂O₅- 30Kg/ha, K₂O - 16.7Kg/ha), T₃ – RDF 100%NPK+ Foliar spray of 3% Zn EDTA, T₄ – RDF 25%P, 100% NK + Seed treatment and two foliar spray of 0.2% Nano DAP, T₅ - RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP, T₆ – RDF 25% P, 100% NK+ Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA, T₇ – RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA. ~~It was foliar sprayed~~ The foliar spray was done at the 45 DAS and 60 DAS. It was laid out in the randomized block design with three replication ~~with having~~ an objective to identify the most efficient fertilizer schedule to enhance the productivity and also to reduce the usage of phosphorus fertilizer. The soil of the experimental field was sandy clay in texture with a pH of 8.41. The soil was low in carbon, low in available nitrogen, low in available phosphorus, high in available potassium and medium in zinc. Biometric observations were taken from ~~the~~ randomly chosen five plants from the each plot and the mean value ~~arrived obtained were was~~ analysed statistically.

Results and discussion

Plant height

Among the treatments, RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇) ~~was~~ recorded the maximum plant height of 55.32 cm. It is followed by the results of RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The least plant height of 40.21cm was observed at the harvesting stage in the control treatment (T₁).

The combination of basal application of conventional fertilizers and foliar spraying of nano fertilizers could have developed a significant increase in plant height by stimulating the plant's auxin metabolism and enzyme activity. This, in turn, might have prompted the cells to elongate and enlarge, ultimately leading to taller plants. This is in conformity with the works of

Yasser *et al.*, (2020) and Gupta *et al.*, (2022).Hagaget *et al.*, (2018) ascribed that nanofertilizers promote the uptake of water and nutrients, which is reflected in plant growth. Moreover, nanofertilizers have a huge surface compared to conventional fertilizers, and this increases the plant's metabolic efficiency.

The application of micronutrients through foliar fertilization resulted in a considerable increase in plant height. This increase in plant height might have been triggered by the micronutrient supply being adequate, which in turn accelerated the enzymatic activity and auxin metabolism in plants (Sudha and Stalin, 2015). An increase in plant height might be due to zinc application and its interrelationship with auxin production, an important growth parameter regulating the stem elongation and cell enlargement (Khanda *et al.*, 1997; Sreenivasa Rao, 2003).

Leaf area index

The maximum leaf area index of 3.56 was noted in RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇) at harvesting stage. This treatment was followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The least leaf area index of 1.93 at harvesting stage was recorded in the control treatment (T₁).

Nano fertiliser easily enters leaves through stomata and other apertures, thus promoting the growth and elongation of the leaf by regulating the rate of cell division or size (Venkatesh *et al.*, 2022). Phosphorus influences plant growth from the cellular to the whole plant level, by increasing leaf area through the processes of cell division and enlargement (Assuero *et al.*, 2004). The findings aligned with the findings of Shawa *et al.*, (2022) for philodendron and Juthery *et al.*, (2020) with wheat. Phosphorus increases leaf area and number along with acceleration of cell division by accumulating at meristematic areas. (Sruthy *et al.*, 2023).

Foliar application of nano DAP (0.4%) at 20 DAS and 40 DAS might have concurred with the nutrient demand of the crop through faster absorption and assimilation as suggested by Manikandan and Subramanian (2016) in maize. Further, supplementing soil application of P with foliar application of nano P might also have resulted in quicker absorption of P in the nano form as it might have easily penetrated into the leaves resulting in more leaf area and LAI. Similar study was conducted by Rashmi and Prakash

Comment [h15]: In reference chapter, the citation starts with different name, this should be corrected here.

Comment [h16]: This is missing in reference chapter.

Comment [h17]: The name in text should always be same as the initial name mentioned in the reference chapter.

(2023).Foliar application of water-soluble fertilizers resulted in enhanced Plant growth, increased translocation of photosynthesis from source to sink, and reduce flower and pod shedding .

Number of branches per plant

Among the treatments, RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇) ~~was~~ recorded the maximum number of branches of 7.92. It is followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅) .The least branches ~~of 5.3~~ was ~~were~~ observed at the harvesting stage in the control treatment (T₁).

The tiny size of nano DAP facilitates the direct adsorption of nutrients into the leaves, where they are more quickly and effectively absorbed. Sufficient nutrients can promote growth and elongation of the cells. More branches per plant result in increased nutrient uptake in plant. Similar observations were recorded by Vagharet *al.*, (2020) and Manjunath Gondi (2018).

By increasing nutrient absorption, the application of nano -P promotes better metabolic processes like photosynthesis, which raises the accumulation of photosynthates and their translocation to the economically valuable parts of the plant, enhancing crop growth, development, and yield. A comparable outcome was observed in broad beans, where the application of nano- fertilizer improved nutrient accumulation and boosted growth activity due to smart delivery system of the fertilizers (Poudelet *al.*, 2023).

Dry matter production

RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇) ~~was registered~~ resulted in the highest dry matter production of 4800 kg ha⁻¹ at the harvesting stage. It is followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The control treatment (T₁) registered the least dry matter production of 3042 kg ha⁻¹.

According to Arifet *al.*, (2016), increasing availability of nutrients to crops through foliar application is a practical method to increase crop yield. Based on their research, they concluded that applying nano-DAP and foliar spraying nutrient solutions at different growth stages

improved wheat yield, yield components, biomass, and pronounced P content – even at 75% lower input than commercial DAP.

Meena *et al.*, (2021) reported after the application of NFs, wheat grain and straw yield increased by 44.6 and 13.1%, respectively. This increase in output could be attributed to higher growth hormone, enhanced metabolic processes, and photosynthetic activities.

Comment [h18]: In reference chapter, the year for this reference is 2020.

In cowpea, foliar application of nano P was positively correlated with DMP (Kumari *et al.*, 2017). A sufficient supply of RDF and foliar application of P in nano form may be the cause of enhanced nutrient intake and efficient photosynthetic translocation, which in turn promotes DMP.

Barua and Sakia (2018) also reported that ZnSO₄ application @ 25kg/ha as basal + 0.5 % foliar application resulted in the plant's maximum straw yield. The reason for this is that zinc has positive influence on root growth, which increase plant nutrient intake from the soil that feeds the aerial portions of the plants and promotes vegetative growth. Likewise, similar conclusion was obtained by Beebout *et al.*, 2011.

Number of pods per plant

In addition, the simultaneous application of traditional and nano DAP ensured optimum and balanced nutrient availability during the crop season, particularly in the crucial stages of the crop. This is because of the nanoparticles' smaller size and greater effective surface area, which allow them to enter the plant more readily and improve nitrogen and phosphorus uptake. Plant components grow to their maximum potential as a result of the increased absorption. Similar results were reported by Manjili *et al.*, 2014 and Abdelghany *et al.*, 2022.

The maximum no. of pods of 19.10 was noted in RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇) at harvesting stage. This treatment was followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The least No. of pods per plant of 13.67 at harvesting stage was recorded in the control treatment (T₁).

Applying Zinc to crops improved the biological activity, growth parameters, and metabolism of nutrients; therefore, applying Zinc increased the activity of enzymes, which in

turn promoted the growth of vegetative branches and pods plant⁻¹(Michailet *al.*, 2004). The seed yield was enhanced by the combined action of growth parameters, yield attributes, and increased cowpea nutrient uptake (Channabasappa *et al.*, 2004).

Under different environmental conditions, the effectiveness of applying soil fertilizer is less than that of foliar fertilization because foliar fertilization provides the necessary nutrients directly to the leaves, absorbs more quickly doesn't depend on root activity, and increase plant growth parameters due to soil water availability (Poudelet *al.*, 2023).

No.of seeds per pod

RDF 50% P, 100% NK + Seed treatment and two foliar spray 0. 2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇) was registered the highest seeds per pod of 13.6 at the harvesting stage. It is followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The control treatment (T₁) registered the least seeds per pod of 9.13.

Then use of Nano-DAP results in a significant increase in the number of seeds produced, particularly when paired with traditional fertilizer. This is because the nutrients are stored in plant cells and released slowly to mitigate biotic and abiotic stress and enhance no of seeds. Increased plant development and metabolic activities, such photosynthesis, are encouraged by nano fertilizers, which increases the concentration and transport of photosynthate to essential plant components. Similar outcomes were reported by El-Azizy *et al.*, (2021) demonstrating the efficacy of nano phosphorus fertilizer in increasing seed yield.

Test weight

However, RDF 50% P, 100% NK + Seed treatment and two foliar spray 0. 2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇) was registered the maximum test weight of 13.90 g. This treatment was followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The least test weight of 13.78 g was noted in the control treatment (T₁).

Legumes are phosphorus- loving plants; they need it for seed development, growth, and most importantly, the energy-producing process of nitrogen fixation. Cowpea yield depends on phosphorus because it is known to promote growth, stimulate nodule development and influence

how effectively of the rhizobium-legume symbiosis functions. (Haruna and Aliyu, 2011). In immature cells with high metabolism and rapid cell division including shoot and root tips, it is essential in enormous amounts. It also promotes the growth of seeds, fruits and flowers (Ndakemi and Dakora, 2007).

Grain yield

Increased yield components may be responsible for higher grain yield. The optimum and balanced nutrient availability was ensured throughout the crop period, particularly during the critical stages of the crop, by applying conventional and nano fertilizers (nano urea and DAP) in combination. This is because nanoparticles have a greater surface area and are smaller size, allowing them to more readily enter plants and improve nitrogen and phosphorus uptake. Increased uptake leads to optimal growth of plant parts and metabolic processes, such as photosynthesis, which increases the accumulation of photosynthates and their translocation to the economically productive parts of the plant. This process increase biomass, yield- attributing characters, and yield by enhancing the translocation of assimilates to seeds – all of which contribute to increased yield. Similar results were reported by Bhargavi and Sundari (2023) and Rajesh (2023).

Comment [h19]: This reference is missing.

The maximum seed yield of 1720 kg ha⁻¹ was achieved by the RDF 50% P, 100% NK + Seed treatment and two foliar spray 0. 2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇). It is followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The control treatment (T₁) resulted in the lowest seed yield of 710 kg ha⁻¹.

Zn foliar treatment significantly increased the production of cowpea grains and straw, which may be associated to its function in photosynthesis, cell division, protein synthesis, membrane structure retention and disease resistance. Since zinc is essential for the proper growth and development of plants, its role in the synthesis if lipids,carbohydrates, and nucleic acid was another factor contributing to the higher yield. Moreover, increased crop performance and yield were naturally the consequence of pollen, fertilization, and germination processes functioning properly as a result of Zn foliar application (Cakmak *et al.*,2008,Aravind,P.,and Prasad 2003, Pandey 2006).

Comment [h20]: Et al. should be removed as citation shows there is only one author.

Comment [h21]: Reference is missing in Reference chapter.

Haulm yield

The treatments, RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇) at harvesting stage recorded the supremely highest haulm yield of 3670 kg ha⁻¹. It is followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The least haulm yield of 2105 kg ha⁻¹ was observed in the control treatment (T₁).

The rapid uptake and ease of translocation of nano fertilizers by plants may have contributed to the increase in haulm production following foliar spraying. This increased rate of photosynthesis and accumulation of dry matter resulted in a higher haulm yield. This corresponds to the results of Mallikarjuna (2021) and Rajput *et al.*, (2022).

Harvest index

Harvest index was significantly influenced by the foliar application of Nano DAP and Zn EDTA fertilization. The significantly highest harvest index of 31.91 was noticed in the RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA (T₇). It is followed by the RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP (T₅). The lesser harvest index of 25.22 was noted in the control treatment (T₁).

Plant growth mechanisms are impacted by the enhanced absorption of nanonutrients caused by the small size of nano fertilizers. The accumulation of dry matter in leaves and proper nutrient supply led to an increase in plant metabolic activities such as photosynthetic and chlorophyll synthesis, both of which promote vegetative growth. This prolonged photosynthesis period additionally enables the plant to grow generally in terms of the harvest index. Similar findings were recorded by Sharma *et al.*, (2022) and Maheta *et al.*, (2023).

Table 1. Effect of Nano DAP and Zn EDTA on growth attributes of cowpea

Treatments	Plant height (cm)	LAI	No. of branches per plant	Dry matter production
T ₁ – Control	40.21	1.93	5.3	3042

T ₂ – RDF 100% NPK (N-12.5kg/ha, P ₂ O ₅ - 30Kg/ha, K ₂ O -16.7Kg/ha)	47.22	2.82	6.95	4159
T ₃ – RDF 100%NPK+ Foliar spray of 3% Zn EDTA	49.87	3.08	7.31	4376
T ₄ – RDF 25%P, 100% NK + Seed treatment and two foliar spray of 0.2% Nano DAP	43.49	2.32	6.18	3718
T ₅ - RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP	52.71	3.31	7.63	4581
T ₆ – RDF 25% P, 100% NK+ Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA	46.19	2.57	6.63	3945
T ₇ – RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA	55.32	3.56	7.92	4800
S.Ed	1.26	0.07	0.11	98
CD (P = 0.05)	2.56	0.18	0.25	197

Table 2. Effect of Nano DAP and Zn EDTA on yield attributes of cowpea

Treatments	No.of pods per plant	Pod length (cm)	No.of seeds per pod	Test weight (g)
T ₁ – Control	13.67	12.11	9.13	13.78

Comment [h22]: This parameter is missing in results.

T ₂ – RDF 100% NPK (N- 12.5kg/ha, P ₂ O ₅ - 30Kg/ha , K ₂ O -16.7Kg/ha)	16.61	14.80	11.87	13.85
T ₃ – RDF 100%NPK+ Foliar spray of 3% Zn EDTA	17.40	15.53	12.39	13.85
T ₄ – RDF 25%P, 100% NK + Seed treatment and two foliar spray of 0.2% Nano DAP	14.92	13.35	10.56	13.81
T ₅ - RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP	18.31	16.25	13.02	13.88
T ₆ – RDF 25% P, 100% NK+ Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA	15.77	14.15	11.21	13.83
T ₇ – RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA	19.10	16.93	13.6	13.90
S.Ed	0.35	0.30	0.23	0.29
CD (P = 0.05)	0.76	0.63	0.51	NS

Table 3. Effect of Nano DAP and Zn EDTA on yield of cowpea

Treatments	Seed yield (kg ha ⁻¹)	Haulm yield (kg ha ⁻¹)	Harvest index
------------	-----------------------------------	------------------------------------	---------------

T ₁ – Control	710	2105	25.22
T ₂ – RDF 100% NPK (N-12.5kg/ha, P ₂ O ₅ - 30Kg/ha , K ₂ O-16.7Kg/ha)	1447	3111	31.74
T ₃ – RDF 100%NPK+ Foliar spray of 3% Zn EDTA	1545	3315	31.79
T ₄ – RDF 25%P, 100% NK + Seed treatment and two foliar spray of 0.2% Nano DAP	1279	2783	31.48
T ₅ - RDF 50% P, 100% NK+ Seed treatment and two foliar spray of 0.2% Nano DAP	1628	3508	31.69
T ₆ – RDF 25% P, 100% NK+ Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA	1358	2949	31.53
T ₇ – RDF 50% P, 100% NK + Seed treatment and two foliar spray 0.2% of Nano DAP + Foliar spray of 3% Zn EDTA	1720	3670	31.91
S.Ed	33	74	0.04
CD (P = 0.05)	68	151	0.11

Reference

Abdelghany, A. M., El-Banna, A. A., Salama, E. A., Ali, M. M., Al-Huqail, A. A., Ali, H. M., & Lamlom, S. F. (2022). The individual and combined effect of nanoparticles and

Comment [h24]: This section should be improved carefully as per the criteria of this journal. Moreover references should be in one format.

biofertilizers on growth, yield, and biochemical attributes of peanuts (*Arachis hypogea* L.). **Agronomy**, **12**(2), 398.

Aravind.P., M.N.V. Prasad, Zinc alleviates cadmium-induced oxidative stress in *Ceratophyllum demersum* L.: a free-floating freshwater macrophyte, **Plant Physiol. Biochem.** **41** (4) (2003) 391–397.

Arif, U., Hussain, S., Shah, S. Z. A., Hamid, A., Yaqoob, A., Arif, A. A., & Muneer, N. (2016). Interactive effect of phosphorus and zinc on the growth, yield and nutrient uptake of garlic (*Allium sativum* L.) variety Gulabi. **Asian Journal of Agriculture and Food Sciences**, **4**(5).

Assuero SG, Mollier A, Pellerin S. The decrease in growth of phosphorus- deficient maize leaves is related to a lower cell production. *Plant, Cell and Environment*. 2004;**27**(7):887-895.

Barua, D., & Saikia, M. (2018). Agronomic biofortification in rice varieties through zinc fertilization under aerobic condition. **Indian Journal of Agricultural Research**, **52**(1), 89-92.

Beebout, S. J., Francis, H. C. R., Dennis, S. J. T., & Rane, C. M. (2011, October). Reasons for variation in rice (*Oryza sativa*) grain zinc response to zinc fertilization. In **3rd International Zinc Symposium** (pp. 10-14).

Bhargavi, G., & Sundari, A. (2023). Effect of nano urea on the growth and yield of rice (*Oryza sativa*) under SRI in the Cauvery delta zone of Tamil Nadu. **Crop Research**, **58**(1and2), 12-17.

Cakmak, I. (2008). Enrichment of cereal grains with zinc: agronomic or genetic biofortification. **Plant and soil**, **302**, 1-17.

Channabasappa KS, Madiwalar SL, Manjappa K, Patil SK. Effect of integrated nutrient management on productivity of rice-cowpea cropping system under low land hill region. **Karnataka J. Agric. Sci.**, 2004;**17**(3):623-625.

Damoar, K., Sharma, R. K., & Maida, P. (2020). Response of cowpea (*Vigna unguiculata* L.) varieties to under Malwa region of Madhya Pradesh. **Journal of Pharmacognosy and Phytochemistry**, 9(2), 1749-1753.

El-Azizy, F. A., & Habib, A. A. M. (2021). Effect of nano phosphorus and potassium fertilizers on productivity and mineral content of broad bean in North Sinai. **Journal of Soil Sciences and Agricultural Engineering**, 12(4), 239-246.

El-Shawa GM, Alharbi K, AlKahtani M, AlHusnain L, Attia KA, Abdelaal K. Improving the quality and production of philodendron plants using nanoparticles and humic acid. **Horticulturae**. 2022;8(8):678.

Gupta SP, Mohapatra S, Mishra J, Yadav SK, Verma S, Singh S, et al. Effect of nano nutrient on growth attributes, yield, Zn content, and uptake in wheat (*Triticum aestivum* L.). **International Journal of Environment and Climate Change**. 2022;12(11):2028-2036.

Hagagg, L.F., Mustafa, N.S., Genaidy, E.A.E., and El-Hady, E.S. (2018) Effect of spraying nano-NPK on growth performance and nutrients status for (Kalamat cv.) olive seedling. **Bioscience Research** 15: 1297–1303

Haruna, I. M., & Aliyu, L. (2011). Yield and economic returns of sesame (*Sesamum indicum* L.) as influenced by poultry manure, nitrogen and phosphorus at Samaru, Nigeria. **Elixir Agric**, 39, 4884-4887.

Haruna, I. M., and Aliyu, L., (2011). Yield and economic returns of sesame (*Sesamum indicum* L.) as influenced by poultry manure, nitrogen and phosphorus at Samaru, Nigeria. **Elixir Agric**, 39: 4884-4887.

Kumari MS, Rao PC, Padmaja G, Ramulu V, Saritha JD, Ramakrishna K. Effect of bio and nano phosphorus on yield, yield attributes and oil content of groundnut (*Arachis hypogaea* L.). **Environment Conservation Journal**. 2017;18(3):21-26.

Maheta A, Gaur D, Patel S. Effect of nitrogen and phosphorus nano-fertilizers on growth and yield of maize (*Zea mays* L.). **Pharma Innovation Journal**. 2023;12(3):2965-2969.

Comment [h25]: This reference is mentioned twice.

Mallikarjuna PR. Effect of nano nitrogen and nano zinc nutrition on nutrient uptake, growth and yield of irrigated maize during summer in the southern transition zone of Karnataka. M. Sc. (Agri.) Thesis, KeladiShivappaNayaka University of Agricultural Sciences Shivmogga (India); c2021.

Manikandan, A., & Subramanian, K. S. (2016). Evaluation of zeolite based nitrogen nano-fertilizers on maize growth, yield and quality on inceptisols and alfisols. **Int. J. Plant Soil Sci**, 9(4), 1-9.

Manjili MJ, Bidarigh S, Amiri E. Study on effect of foliar application of nano chelate molybdenum fertilizer on yield and yield components of peanut. **Egyptian Academic Journal of Biological Sciences**. 2014;5(1):67-71.

Manjunath Gondi. Effect of nano nutrients on growth, seed yield and quality in cowpea [*Vignaunguiculata* (L.) Walp.]. M.Sc (Agri) Thesis, Univ. Agric. Sci., GKVK, Bengaluru, 2018.

Meena, R.H.; Jat, G.; Jain, D. Impact of foliar application of different nano-fertilizers on soil microbial properties and yield of wheat. **J. Environ. Biol**. 2020, 42, 302–308.

Comment [h26]: Year is different in the text.

Michail T, Walter T, Astrid W, Walter G, Dieter G, Maria SJ et al. A survey of foliar mineral nutrient concentrations of *Pinuscanariensis* at field plots in Tenerife. **Forest. Ecol. Manag** ,2004 :189:49 -55.

Ndakidemi, P. A., & Dakora, F. D. (2007). Yield components of nodulated cowpea (*Vignaunguiculata*) and maize (*Zea mays*) plants grown with exogenous phosphorus in different cropping systems. **Australian Journal of Experimental Agriculture**, 47(5), 583-589.

Nkaa, F., Nwokeocha, O. W., & Ihuoma, O. (2014). Effect of phosphorus fertilizer on growth and yield of cowpea (*Vignaunguiculata*). **IOSR Journal of Pharmacy and Biological Sciences**, 9(5), 74-82.

Comment [h27]: Not present in text.

Paikaray, R. K., Mishra, K. N., Khanda, C. M., & Garnayak, L. M. (1997). Effect of nitrogen, phosphorus and plant density on yield and nutrient uptake in late sown niger (*Guizotia abyssinica*).

Comment [h28]: Missing in text.

Pandey, N., & Gupta, B. (2012). Improving seed yield of black gram (*Vigna mungo* L. var. DPU-88-31) through foliar fertilization of zinc during the reproductive phase. **Journal of plant nutrition**, 35(11), 1683-1692.

Comment [h29]: Not present in text.

Poudel A, Singh SK, Jiménez-Ballesta R, Jatav SS, Patra A, Pandey A. Effect of nano-phosphorus formulation on growth, yield and nutritional quality of wheat under semi-arid climate. **Agronomy**. 2023;13(3):768.

Rajput JS, Thakur AK, Nag NK, Chandrakar T, Singh DP. Effect of nano fertilizer in relation to growth, yield and economics of little millet (*Panicum sumatrense* Roth) under rainfed conditions. **Pharma Innovation Journal**. 2022;11(7):153-156.

Rashmi, C. M., & Prakash, S. S. (2023). Effect of nano phosphorus fertilizers on growth and yield of maize (*Zea mays* L.) in central dry zone of Karnataka.

Rupa, T. R., Rao, C. S., Rao, A. S., & Singh, M. (2003). Effects of farmyard manure and phosphorus on zinc transformations and phyto-availability in two alfisols of India. **Bioresource Technology**, 87(3), 279-288.

Comment [h30]: Missing in text.

Sharma SK, Sharma PK, Rameshwar LM, Sharma V, Chaudhary R, Pandey R, et al. Effect of foliar application of nano-urea under different nitrogen levels on growth and nutrient content of pearl millet (*Pennisetum glaucum* L.). **International Journal of Plant and Soil Science**. 2022;34(20):149-155.

Sruthy, H., Pillai, P. S., Shimi, G. J., Bindhu, J. S., & Sajeena, A. (2023). Growth and Yield of Grain Cowpea (*Vigna unguiculata* sub sp. *Cylindrica*) in Response to Foliar Nutrition and Graded Levels of Phosphorus and Potassium. **International Journal of Environment and Climate Change**, 13(11), 4001-4014.

Sudha, S., & Stalin, P. (2015). Effect of zinc on yield, quality and grain zinc content of rice genotypes. **International journal of farm sciences**, 5(3), 17-27.

- Vaghar MS, Sayfzadeh S, Zakerin HR, Kobraee S, Valadabadi SA. Foliar application of iron, zinc and manganese nano-chelates improves physiological indicators and soybean yield under water deficit stress. **Journal of Plant Nutrition**. 2020;**43**(18):2740-2756.
- Venkatesh M, Babu KK, Prasanth P, Lakshminarayana D, Kumar SP. Study on effect of different levels of nitrogen in combination with nano urea on growth and yield of marigold (*Tagetes erecta* L.) Cv. PusaNarangiGainda. **The Pharma Innovation Journal**. 2022;**11**(11):1313-1317.
- WA Al-juthery H, HilalObaid Al-Maamouri E. Effect of urea and nano-nitrogen fertigation and foliar application of nano-boron and molybdenum on some growth and yield parameters of potato. Al-Qadisiyah, **Journal for Agriculture Sciences**. 2020;**10**(1):253-263.
- Yasser E, El-Ghobashy, Elmehy AA, El-Douby KA. Influence of Intercropping Cowpea with some maize hybrids and N nano mineral fertilization on productivity in salinity soil. **Egyptian Journal of Agronomy**. 2020;**42**(1):63-78.