

Assessing the Efficacy and Biological Benefits of Withanolide-Rich *Withania somnifera* ("Ashwagandha") Root Extract

Formatted: Centered

Formatted: Font: 16 pt

ABSTRACT:

The medicinal plant *Withania somnifera*, usually referred to as Ashwagandha, is a member of the Solanaceae family. The presence of Withanolides in the roots is responsible for a number of pharmacological effects in Ashwagandha. Withanolides have been demonstrated to be an effective neuronal, immune, anti-stress, and anti-cancer agent. However, Withanolides demonstrated limited permeability, lowering the bioavailability and efficacy of active compounds. The goal of the study was to ascertain the biological efficacy of Ashwagandha Composition, a blend of *W. somnifera* Ashwagandha milk extract and water extract (1:1) with a high concentration of Withanolides (3-5%), at lower dosages with improved bioabsorption compared to pure Ashwagandha Hydro ethanolic extracts fortified with 2.5% withanolides. The *W. somnifera* Ashwagandha Composition was assessed for its bioabsorption and bioefficacy by exploring its intestinal absorption capacity, Acetylcholinesterase Inhibition, antioxidant potential, glutathione reduction potential, anti-inflammatory activity, and anti-cancer activity. Additionally, Ashwagandha Composition was also evaluated for its safety profile. We found *W. somnifera* Ashwagandha Composition (1:1) is more bioavailable and showed higher biological activity than the Ashwagandha Hydro ethanolic extract fortified with 2.5% withanolides (Ashwagandha extract). Hence, the *W. somnifera* Ashwagandha Composition can be used as a therapeutic medication once its safety concerns are addressed by *in vivo* trials.

Commented [MOU1]: Give values

KEYWORDS: *Withania somnifera*, Withanolides, Bioabsorption, Bioactivity.

1. INTRODUCTION

Withania somnifera, popularly recognized as "Ashwagandha," "Indian Winter Cherry," or "Indian Ginseng," is a key therapeutic plant in Ayurveda, India's holistic medical system (1). For millennia, it has been utilized as a Rasayana for its wide spectrum of health virtues. This herbal concoction fosters physical and mental health and elevates oneself. Tonics of this type are administered to little children, and they are also used to promote life expectancy in the

middle-aged and elderly (Ref). Ashwagandha is the most important Rasayana herb used in Ayurvedic therapy (2). It is known as "Sattvic Kapha Rasayana" herb (3). A significant proportion of Rasayana herbs act as anxiety relievers (ref).

~~Withania somnifera~~ Ashwagandha is typically marketed as a churna, a finely ground powder that can be combined with ghee, water, or honey for ~~admisteration~~ administration. It is well-recognized for enhancing energy and stamina and for having sedative, diuretic, and anti-inflammatory properties (Ref). It promotes memory and improves neurological function (Ref). By enhancing the reproductive system's functionality, it promotes the harmony of sexual and reproductive activities. It boosts the body's tolerance to stress due to its high adaptogenic characteristics (Ref). ~~Withania somnifera~~ Ashwagandha strengthens the body's resistance ~~against to~~ illness through increasing cell-mediated immunity. Additionally, it has strong antioxidant qualities that aid in preventing cellular damage brought on by oxidants (2). ~~Withania-W~~ somnifera has a long history of being used as a powerful rejuvenator, tonic, and therapy for an array of diseases. It also functions as an adaptogen and has potent anti-stress and immunostimulatory properties (4). As a result, in the Indian Herbal System, *W. somnifera* is recognized as one of the most pivotal herbs and an ideal adaptogen.

~~Withania somnifera~~ Ashwagandha has a ~~number of several~~ pharmacological effects, which are mostly ascribed to the bioactive constituents, such as Withanolides, found in the roots of the herb. The primary Withanolides isolated from the roots of *W. somnifera* are Withanoside V (WS V), Withaferin A, Withanolide A (WN A), and Withanolide B (WN B), which aggregate to about 35 Withanolides in total (5-7). Withaferin A exhibits anti-inflammatory, anti-cancer, anti-angiogenesis, and anti-apoptotic properties. ~~Withanolide WN~~ have been shown to be useful in the treatment of an array of malignancies, such as breast cancer, pancreatic cancer, and skin cancer (8). ~~Withanolide WN~~ A is an effective neuronal, immune, and anti-stress drug. ~~Withanolide WS~~ IV and V are crucial in neuroregeneration. However, the permeability of the glycosylated and polar Withanolide glycosides WS IV and WS V is poor. As a result, the bioavailability and efficacy of active substances containing anolide glycosides are reduced (9). Therefore, the present invention relates to making ~~W. somnifera~~ Ashwagandha more bioavailable to the body and thereby enhancing its bioactivity.

The current study investigates the bioactivity of ~~W. somnifera~~ Ashwagandha Composition, which is a combination of ~~W. somnifera~~ Ashwagandha milk extract and water extract with a high concentration of Withanolides. The bioactivity of Ashwagandha Hydro ethanolic extract

Commented [MOU2]: Use scientific name

Commented [MOU3]: When scientific name at the beginning write full genus name

Commented [MOU4]: Use scientific name

Commented [MOU5]: Don't use abbreviation at the beginning of sentence

Commented [MOU6]: Don't use abbreviation at the beginning of sentence

Commented [MOU7]: Don't use abbreviation at the beginning of sentence

fortified with 2.5% withanolides (*W. somnifera* Ashwagandha extract) was used to compare the effectiveness of this *W. somnifera* Ashwagandha Composition. The goal of this study is to look into its effectiveness as a potent acetylcholinesterase (AChE) inhibitor, anti-oxidant, cyclooxygenase inhibitor, and anti-cancer agent.

2. METHODS

2.1 Sample collection and authentication

2.2 Sample preparation

An herbal composition was formulated with a mixture of *W. somnifera* Ashwagandha milk extract and water extract (1:1) with a high concentration of Withanolides by using the process discussed by Reddy et al (9). This composition was assessed for its biological activities and efficacy was compared with the bioactivity of ashwagandha Hydro ethanolic extract fortified with 2.5% withanolides (~~Ashwagandha extract~~).

2.1 Bioavailability Assessment of Withanolides in *W. somnifera* Ashwagandha Extracts using Caco-2 Model

Cytotoxicity tests determined the non-toxic concentration for absorption trials. In three separate experiments, the permeability test was carried out in the apical-basolateral (A-B) and basolateral-to-apical (B-A) directions at 37 °C for 2 hours in a shaking incubator with 100 rpm using *W. somnifera* Ashwagandha extracts dissolved in HBSS buffer. Reverse phase chromatography was used to analyze the transported materials using a C18 column (column length) and UV-VIS detector in order to quantify them (Waters HPLC system, Milford, MA, USA (give details about flow rate, temperature, mobile phase)). Drug absorption studies on Caco-2 cells involved Ashwagandha Composition administered at 1 mg/mL, analyzed over different time intervals. The apparent permeability coefficient of the compounds was computed (10). The apparent permeability coefficient (Papp) was computed using MS Excel using the standard formula (11, 12).

Commented [MOU8]: Give details about Sample collection and authentication

Formatted: Indent: Left: 0.54"

Commented [MOU9]: Give details about sample preparation

Formatted: Outline numbered + Level: 2 + Numbering Style: 1, 2, 3, ... + Start at: 1 + Alignment: Left + Aligned at: 0.25" + Indent at: 0.54"

Formatted: Strikethrough

Commented [MOU10]: Write expanded name and put HBSS abbreviation with the bracket

2.2 Acetylcholinesterase Inhibition Assay

The Acetylcholinesterase (AChE) Assay Kit, employing DTNB, measured AChE activity (13). Thiolcholine levels indicated AChE inhibition, following the manufacturer's protocol (Thermo Fisher, Scientific, India).

Commented [MOU11]: Give expanded name of DTNB and give DTNB abbreviation within bracket

2.3 Radical Scavenging Activity

Withania somnifera Ashwagandha Composition and extract were evaluated for antioxidant potential. DPPH assay at various concentrations (Give concentration range with units) and absorbance measurement at 517 nm determined percentage inhibition. All chemicals and standards were purchased from Thermo Fisher, Scientific, India (14).

Commented [MOU12]: Use scientific name

Commented [MOU13]: Give expanded name of DPPH and give DPPH abbreviation within bracket

Commented [MOU14]: What is the positive stand you used? Give details with concentration range

2.4 Glutathione Reduction Assay

Assessment of *W. somnifera* Ashwagandha Composition's ability to generate free radicals at different concentrations (Give concentration range with units) was measured using Glutathione Reduction Assay kit (Thermofisher, Scientifics, India). Glutathione was purchased from Sr. Lifesciences. Absorbance was measured at 412 nm for radical formation after incubation (15).
[What is your positive control?](#)

2.5 *In-vitro* Anti-inflammatory Activity

COX-2 inhibitor screening kit (Thermofisher, Scientifics, India) measured fluorescence during prostaglandin G2 production by COX-2 enzyme. prostaglandin G2 produced was assessed for the anti-inflammatory activity of *W. somnifera* Ashwagandha extract and Composition composition (16).

Formatted: Font: Italic

Commented [MOU15]: Give expanded name of COX-2 and give COX-2 abbreviation within bracket

2.6 *In-vitro* Anti-cancer Activity

MTT method was used to evaluate Ashwagandha compositions against pancreatic, breast, and colorectal cancer cells (17). A Tecan Microplate reader was employed to measure the absorbance at 570 nm. Megalan software (Tecan Instruments, Inc.) was used to collect the data, which was then exported in Microsoft Excel format for additional analysis (18).

Formatted: Font: Italic

Commented [MOU16]: Give expanded name of MTT and give MTT abbreviation within bracket

2.7 Toxicity Studies

In-vitro cytotoxicity of *W. somnifera* Ashwagandha extract and Composition against kidney epithelial (Hek293T) and Skin Fibroblast cells (HFF-1) was assessed using the MTT method. Absorbance at 570 nm measured cytotoxic effects at various concentrations (17, 18).

Formatted: Font: Italic

Statistical Analysis

Commented [MOU17]: Give details about statistics

3. RESULTS (Is this results or results and discussion)

3.1 Determining the bioavailability of Withanolides of ashwagandha Composition using an *in-vitro* absorption model system.

Withania somnifera Ashwagandha has a multitude of pharmacological properties that are mostly due to the bioactive constituents, such as Withanolides, found in the roots of the herb. The effectiveness of bioactive Withanolide components is reliant on their absorption and transit across the intestinal epithelium. The *in-vitro* cell culture method has been touted as a potential model system for efficiently determining a drug's bioavailability. We described the bioavailability profiles of primarily Withanolides from *W. somnifera* in this study using the Caco-2 cell culture method. An important requirement for a medicine to be taken orally is that it has adequate intestinal absorption in humans; in the early phases of drug discovery, this requirement is often tested using *in vitro* techniques. There are several *in-vitro* intestinal absorption models that fall under the categories of physicochemical and biological techniques (19).

Commented [MOU18]: Use scientific name

~~The~~ Table. 1 and Figure 1 show the results ~~for of~~ the analysis of *W. somnifera* Ashwagandha extract and ~~Ashwagandha~~ Composition's intestinal absorption in Caco-2 cells. When compared to pure Ashwagandha extracts, the Caco-2 *in-vitro* absorption study ~~clearly~~ demonstrated that the synergistic combination of *W. somnifera* Ashwagandha milk extract and act with a high concentration of Withanolides improved membrane permeability and absorption. The *W. somnifera* Ashwagandha component was completely absorbed by the Caco-2 cells from the medium after 5 minutes, as validated by HPLC qualitative evaluation.

Formatted: Strikethrough

At a concentration of 0.125 mg/mL, the transfer of active components from the apical to the basolateral (ab) was studied. Ashwagandha Composition was found to have the highest apparent permeability coefficient (Papp) in absorptive transport at $164.5 \pm 13.5 \text{ cm-s}^{-1}$ compared to the secretive direction at $89.6 \pm 11.4 \text{ cm-s}^{-1}$ and an efflux ratio of 0.54. Whereas Ashwagandha extract was found to have an apparent permeability coefficient (Papp) in absorptive transport as $142.2 \pm 16.3 \text{ cm-s}^{-1}$ compared to secretive direction as $71.2 \pm 7.2 \text{ cm-s}^{-1}$ and with an efflux ratio of 0.50.

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Formatted: Superscript

Figure 1: Graph representing the Intestinal absorption of Ashwagandha extract and Ashwagandha Composition in Caco-2 cells.

Commented [MOU19]: Put standard error bars




Figure 1: Graph representing the Intestinal absorption of *W. somnifera* extract and Composition in Caco-2 cells.

Commented [MOU20]: Label figures at the bottom

Table 1: Interpretation of the Intestinal absorption of Ashwagandha extract and Ashwagandha Composition in Caco-2 cells.

Papp X 10 ⁻⁶ (cm/s)			
Drugs	Concentration (mg/mL)	Apical to Basolateral transport	Basolateral to Apical transport

<u>W.</u> <u>somnifera</u> Ashwagandha extract	0.125	142.2 ± 16.3	71.2 ± 7.2
<u>W.</u> <u>somnifera</u> Ashwagandha Composition	0.125	164.5 ± 13.5	89.6 ± 11.4

At different time intervals, Caco-2 cell lysate and media samples of W. somniferaAshwagandha Composition were screened. The samples were qualitatively determined with an analytical standard by the HPLC analysis using Shimadzu's Lab Solutions software with a photo-diode array detector.

The above Caco-2 *in-vitro* absorption study (Fig. 1 and Table. 2) ~~clearly~~ revealed that Ashwagandha Composition which is a synergistic mixture of Ashwagandha milk extract and water extract with a high concentration of Withanolides exhibited good membrane permeability and absorption compared to the as such W. somniferaAshwagandha extracts.

At the 5-minute time interval, the Ashwagandha Composition was completely absorbed by the Caco-2 cells from the media, which was confirmed by HPLC qualitative determination (Fig. 2).

Commented [MOU21]: Resolution of the Figure 2 is not enough. Put an image with good quality




Fig.2: *Withania somnifera* Ashwagandha Composition HPLC chromatogram- qualitative determination of Withanolides in Caco-2 cells lysate and media at 5 min time interval.

Commented [MOU22]: Use scientific name

3.2 *Withania somnifera* Ashwagandha Root Extract's Anti-Acetylcholinesterase Activities:

Commented [MOU23]: Use scientific name

Withania somnifera Ashwagandha is given for the treatment of memory and cognitive deficiencies caused by accidents, sickness, or simply old age. Preclinical research has suggested that ashwagandha may act as a nootropic, boosting cognitive function and memory (20). The present study provides a novel synergistic mixture of ashwagandha milk extract and water extract with a high concentration of Withanolides, wherein the said *W. somnifera* Ashwagandha Composition proved to be a potent AChE inhibitor in comparison to *W. somnifera* Ashwagandha Hydro ethanolic extract and standards such as donepezil. The graph below interprets the data for the Acetylcholinesterase Inhibition Potential of *W. somnifera* Ashwagandha Composition in comparison to *W. somnifera* Ashwagandha extract (Fig. 3). Ashwagandha extract inhibited Acetylcholinesterase with an IC₅₀ of 4.4 mg/ml and a 62% inhibition. Whereas *W. somnifera* Ashwagandha extract was discovered to possibly inhibit Acetylcholinesterase with an IC₅₀ value of 3.57 mg/ml and 84% inhibition.

Commented [MOU24]: Use scientific name

Commented [MOU25]: Use scientific name

Commented [MOU26]: Use scientific name

Commented [MOU27]: Use scientific name

Commented [MOU28]: Use scientific name

Commented [MOU29]: Use scientific name




Figure 3: Graph representing the percentage of Acetylcholinesterase inhibition for *W. somnifera* Ashwagandha extract and Ashwagandha Composition

Commented [MOU30]: Use scientific name

3.3 Anti-oxidant Potential of the *W. somnifera* Ashwagandha Composition

Commented [MOU31]: Use scientific name

For ages, Ayurvedic medicine has employed ashwagandha to increase longevity and vitality. The roots of this plant have strong antioxidant capacity (14, 21). Through this study we assessed the antioxidant and free radical scavenging activity of *W. somnifera* Ashwagandha Composition. A DPPH assay was performed to screen for anti-oxidant activity of the *W. somnifera* Ashwagandha Composition which is a mixture of *W. somnifera* Ashwagandha milk extract and Ashwagandha-water extract with a high concentration of Withanolides in comparison with *W. somnifera* Ashwagandha Hydro ethanolic extract. The graph (Fig. 4) and table (Table. 2) below provides an interpretation of the findings on the anti-oxidant potential of the *W. somnifera* Ashwagandha Composition in comparison to the commercially available *W. somnifera* Ashwagandha extract and other standards such as ascorbic acid. The IC₅₀ value of Ashwagandha extract was found to be ~ 47.21 µg/mL. Unlike Ascorbic acid (Positive control), which had an IC₅₀ value of ~7.12 g/mL, Ashwagandha Composition demonstrated potential anti-oxidant activity with an IC₅₀ value of ~32.16 g/mL. The findings from this study clearly indicate that *W. somnifera* Ashwagandha Composition has strong antioxidant properties.

Commented [MOU32]: Use scientific name

Commented [MOU33]: Use scientific name

Commented [MOU34]: Use scientific name

Commented [MOU35]: Use scientific name

Commented [MOU36]: Use scientific name

Commented [MOU37]: Use scientific name

Commented [MOU38]: Are there any statistical different between test samples as well as in between test samples and positive control

Commented [MOU39]: Use scientific name




Figure 4: Graph representing the percentage of Radical scavenging activity for *W. somnifera* Ashwagandha extract and Ashwagandha Composition

Commented [MOU40]: Use scientific name

Table 2: DPPH free radical scavenging assay

S. No	Name of the compound	DPPH IC ₅₀ µg/mL
1	Ascorbic acid	~ 7.12
2	<i>W. somnifera</i> Ashwagandha extract	~ 47.21

Commented [MOU41]: Use scientific name

3	W. somnifera Ashwagandha Composition	~ 32.16
---	---	---------

Commented [MOU42]: Use scientific name

3.4 Glutathione Reduction assay of the Ashwagandha herbal Composition

[Withania somnifera](#) *W. somnifera*'s is also known for its capacity to repair oxidative damage (22). A glutathione reduction assay was performed to screen for the ability of [W. somnifera's Ashwagandha](#) Composition and [Ashwagandha](#) extract to generate ROS. The results for the Glutathione reduction potential of [W. somnifera's Ashwagandha](#) herbal Composition in comparison to the [W. somnifera's Ashwagandha](#) extract are interpreted in the following graph (Fig. 5). Ashwagandha extract was found to reduce glutathione with ~ 11.06% inhibition at 1 $\mu\text{g}/\text{mL}$. In comparison to ascorbic acid (the positive control), which showed only 7% glutathione reduction at 1 g/mL , [W. somnifera's Ashwagandha](#) Composition was found to reduce glutathione with 15.6% inhibition at 1 g/mL .

Commented [MOU43]: Use scientific name




Figure 5: Graph representing the percentage Glutathione Reduction for [W. somnifera](#) [Ashwagandha](#) extract and [Ashwagandha](#) Composition

Commented [MOU44]: Use scientific name

3.5 *In-vitro* anti-inflammatory activity of the [W. somnifera](#) [Ashwagandha](#) herbal Composition

Commented [MOU45]: Use scientific name

Withania W. somnifera, notably the roots with their distinct components, has been demonstrated to be beneficial against a variety of anti-inflammatory activity (23). In the present study, An anti-inflammatory assay was performed using a COX-2 inhibitor screening kit. The anti-inflammatory activity of *W. somnifera* Ashwagandha Composition and Ashwagandha extract is represented in the graph below (Fig. 6). At 1 mg/ml, Ashwagandha extract inhibited COX-2 with an 86.4 percent inhibition. Whereas celecoxib (positive control) exhibited only 69% COX-2 inhibition, and Ashwagandha Composition was found to potentially inhibit COX-2 with 87.2 percent inhibition at 1 mg/ml.

Commented [MOU46]: Use scientific name




Figure 6: Graph representing the percentage of COX-2 Inhibition for *W. somnifera* Ashwagandha extract and Ashwagandha Composition.


Commented [MOU47]: Use scientific name

3.6 In-vitro anti-cancer activity of the *W. somnifera* Ashwagandha herbal Composition

Commented [MOU48]: Use scientific name

The two main active components, Withanolides and withaferins, as well as a few additional metabolites such as withanone (WN) and Withanosides, have been shown to be effective against several cancer cell lines (25-27). MTT *in vitro* cell proliferation assays were performed on MIA PaCa-2, MBA-MB-231, MDA-MB-468, and HCT116 cell lines. *Withania somnifera*

Commented [MOU49]: Use scientific name



Ashwagandha extract and Ashwagandha Composition were checked at various concentrations, including 0.1, 0.25, 0.5, and 1 mg/mL, for their percentage inhibition of cell proliferation. A Tecan Multimode reader was used to record the data and plot a non-linear regression curve against the log concentration versus absorbance as shown in Figs. 7, 8, 9, and 10. Ashwagandha extract (Figs. 7A, 8A, 9A, and 10A) was found to effectively inhibit MIA PaCa-2, MBA-MB-231, MDA-MB-468 and HCT116 cells with an IC₅₀ value of ~ 12.67, ~ 2.93, ~ 6.88, and ~ 1.47 µg/mL, respectively. Whereas, Ashwagandha Composition (Figs. 7B, 8B, 9B, and 10B) was found to effectively inhibit MIA PaCa-2, MBA-MB-231, MDA-MB-468, and HCT116 cells with an IC₅₀ value of ~ 8.52, ~ 1.72, ~ 1.46, and ~ 1.68 µg/mL, respectively.

Figure 7: Graph illustrating the percentage inhibition of proliferation of MIA PaCa-2 cells on treatment with (A) Ashwagandha extract and (B) Ashwagandha Composition




Figure 8: Graph illustrating the percentage inhibition of proliferation of MDA-MB-231 cells on treatment with (A) *W. somnifera* Ashwagandha extract and (B) *W. somnifera* Ashwagandha Composition

Commented [MOU50]: Use scientific name

Commented [MOU51]: Use scientific name




Figure 9: Graph illustrating the percentage inhibition of proliferation of MDA-MB-468 cells on treatment with (A) *W. somnifera* Ashwagandha extract and (B) *W. somnifera* Ashwagandha Composition

Commented [MOU52]: Use scientific name

Commented [MOU53]: Use scientific name




Figure 10: Graph illustrating the percentage inhibition of proliferation of HCT116 cells on treatment with (A) *W. somnifera* Ashwagandha extract and (B) *W. somnifera* Ashwagandha Composition

Commented [MOU54]: Use scientific name

Commented [MOU55]: Use scientific name

3.7 Toxicity Profiling of *W. somnifera* Ashwagandha herbal Composition

In medication research, determining a molecule's non-clinical toxicity and pharmacokinetics is crucial since it sheds light on its safety and posology in individuals. Understanding the toxicity of *W. somnifera* extract has been the focus of intensive research by several scientists around the globe (28, 29). Although WA is the part of *W. somnifera* that is most physiologically active, there is little to no evidence released regarding its safety or toxicity on oral administration. MTT *in-vitro* cytotoxicity assays were performed on Hek293T and HFF-1 cell lines. *Withania somnifera* Ashwagandha extract and Ashwagandha Composition were checked at

Commented [MOU56]: Use scientific name

Commented [MOU57]: Use scientific name

various concentrations, including 0.1, 0.25, 0.5, and 1 mg/mL, for their percentage inhibition of cell proliferation. A Tecan Multimode reader was used to record the data and plot a non-linear regression curve against the log concentration versus absorbance, as shown in Figs. 11 and 12. *W. somnifera* Ashwagandha extract (Figs. 11A and 12A) was found to be non-toxic against Hek293T and HFF-1 with an IC₅₀ value of ~ >1 mg/mL and ~ 958.5 µg/mL, respectively. In contrast, *W. somnifera* Ashwagandha Composition (Figs. 11B and 12B) was found to be safe and non-toxic against Hek293T and HFF-1, with IC₅₀ values of ~ > 1 mg/mL and ~1 mg/mL, respectively.

Commented [MOU58]: Use scientific name

Commented [MOU59]: Use scientific name




Figure 11: Graph illustrating the percentage inhibition of proliferation of Hek293T cells on treatment with (A) *W. somnifera* Ashwagandha extract and (B) *W. somnifera* Ashwagandha Composition

Commented [MOU60]: Use scientific name

Commented [MOU61]: Use scientific name




Figure 12: Graph illustrating the percentage inhibition of proliferation of HFF-1 cells on treatment with (A) *W. somnifera* Ashwagandha extract and (B) *W. somnifera* Ashwagandha Composition

Withania somnifera Ashwagandha is a common folk treatment in India, where people utilize it as an aphrodisiac for young people and as the finest tonic for the elderly and young. Previous research has yielded pure Withanolides with little toxicity. Although the key downside of the previous study is its low bioavailability, Due to its decreased intestinal absorption, this limits its ability to perform therapeutically (30). The current study aims to increase the bioavailability of active ingredients in *W. somnifera* Ashwagandha extract by combining *W. somnifera* Ashwagandha milk water and Ashwagandha–water extract with a high concentration of Withanolides. This helps to fully harness its therapeutic potential. However, further *in vivo* studies should be explored to completely eliminate the toxicity concerns and confirm its safety for clinical use.

4. CONCLUSION

The growing incidence of cancer-related mortality necessitates the development of innovative chemopreventive medicines with minimal detrimental reactions. Distinct active constituents extracted from different parts of the *W. somnifera* Ashwagandha herb have been shown to exhibit potent anti-tumor and immunomodulating properties. A mixture of ashwagandha milk extract and water extract containing a high concentration of Withanolides was found to effectively inhibit the proliferation of MIA PaCa-2, MDA-MB-231, MDA-MB-468, and HCT116 cells, displaying significant anti-cancer properties. Additionally, the *W. somnifera* Ashwagandha Composition showed better bioabsorption and bioefficacy by exploring its intestinal absorption capacity as well as improved Acetylcholinesterase Inhibition, antioxidant potential, glutathione reduction potential and anti-inflammatory activity. Therefore, this extract can be employed as such or in the form of a concoction along with other therapeutic agents for *in-vivo* trials on humans.

REFERENCE

1. Misra L, Lal P, Sangwan RS, Sangwan NS, Uniyal GC, Tuli R. Unusually sulfated and oxygenated steroids from *Withania somnifera*. *Phytochemistry* 2005;66:2702-7.

Commented [MOU62]: Use scientific name

Commented [MOU63]: Use scientific name

Commented [MOU64]: Use scientific name

Commented [MOU65]: Use scientific name

Commented [MOU66]: Use scientific name

Commented [MOU67]: Use scientific name

Commented [MOU68]: Use scientific name

2. Singh N, Bhalla M, de Jager P, Gilca M. An overview on ashwagandha: a Rasayana (rejuvenator) of Ayurveda. *African Journal of Traditional, Complementary and Alternative Medicines* 2011;8(5S).
3. Sharma CG. *Ashwagandharishta–Rastantra Sar Evam Sidhyaprayog Sangrah–Krishna–Gopal Ayurveda Bhawan (Dharmarth Trust) 1938.*
4. Choudhary B, Shetty A, Langade DG. Efficacy of Ashwagandha (*Withania somnifera* [L.] Dunal) in improving cardiorespiratory endurance in healthy athletic adults. *Ayu* 2015;36:63.
5. Sangwan RS, Chaurasiya ND, Lal P, Misra L, Tuli R, Sangwan NS. Withanolide A is inherently de novo biosynthesized in roots of the medicinal plant Ashwagandha (*Withania somnifera*). *Physiologia plantarum* 2008;133:278-87.
6. Chatterjee S, Srivastava S, Khalid A, Singh N, Sangwan RS, Sidhu OP, et al. Comprehensive metabolic fingerprinting of *Withania somnifera* leaf and root extracts. *Phytochemistry* 2010;71:1085-94.
7. Devkar S, Badhe Y, Jagtap S, Hegde M. Quantification of major bioactive Withanolides in *Withania somnifera* (Ashwagandha) roots by HPTLC for rapid validation of Ayurvedic products. *JPC-Journal of Planar Chromatography-Modern TLC* 2012;25:290-4.
8. Malik F, Kumar A, Bhushan S, Mondhe DM, Pal HC, Sharma R, et al. Immune modulation and apoptosis induction: Two sides of antitumoural activity of a standardised herbal formulation of *Withania somnifera*. *European Journal of Cancer* 2009;45:1494-509.
9. Reddy, K. N. Devaraj, Aluri, Srilakshmi, Shetty, Prathvi, Udaya, Shreya, BK, Nalina, MS, Sudhanva, Rangappa, Shobith. "Formulation of A Simplified Process for Extraction of High Concentration of Bioavailable Withanolides from Ashwagandha Root." **Journal of Research in Pharmaceutical Science**, vol. 10, no. 3, 2024, pp. 15-20.
10. Haltner-Ukomadu E, Gureyeva S, Burmaka O, Goy A, Mueller L, Kostyuk G, Margitich V. In vitro bioavailability study of an antiviral compound enisamium iodide. *Scientia Pharmaceutica* 2018;86:3.
11. van Breemen RB, Li Y. Caco-2 cell permeability assays to measure drug absorption. *Expert opinion on drug metabolism & toxicology* 2005;1:175-85.

12. Hubatsch I, Ragnarsson EG, Artursson P. Determination of drug permeability and prediction of drug absorption in Caco-2 monolayers. *Nature protocols* 2007;2:2111-9.
13. Miao Y, He N, Zhu JJ. History and new developments of assays for cholinesterase activity and inhibition. *Chemical reviews* 2010;110:5216-34.
14. Manwar J, Mahadik K, Sathiyarayanan L, Paradkar A, Patil S. Comparative antioxidant potential of *Withania somnifera* based herbal formulation prepared by traditional and non-traditional fermentation processes. *Integrative Medicine Research* 2013;2:56-61.
15. Nuhu F, Gordon A, Sturme R, Seymour AM, Bhandari S. Measurement of glutathione as a tool for oxidative stress studies by high performance liquid chromatography. *Molecules* 2020;25:4196.
16. Jayaprakasam B, Nair MG. Cyclooxygenase-2 enzyme inhibitory Withanolides from *Withania somnifera* leaves. *Tetrahedron* 2003;59:841-9.
17. Badieyan ZS, Moallem SA, Mehri S, Shahsavand S, Hadizadeh F. Virtual screening for finding novel COX-2 inhibitors as antitumor agents. *The open medicinal chemistry journal* 2012;6:15.
18. Van Meerloo J, Kaspers GJ, Cloos J. Cell sensitivity assays: the MTT assay. *Cancer cell culture: methods and protocols* 2011;237-45.
19. Turco L, Catone T, Caloni F, Di Consiglio E, Testai E, Stamatii A. Caco-2/TC7 cell line characterization for intestinal absorption: how reliable is this in vitro model for the prediction of the oral dose fraction absorbed in human?. *Toxicology in vitro* 2011;25:13-20.
20. Choudhary D, Bhattacharyya S, Bose S. Efficacy and safety of Ashwagandha (*Withania somnifera* (L.) Dunal) root extract in improving memory and cognitive functions. *Journal of Dietary Supplements* 2017;14:599-612.
21. Tiwari P, Patel RK. Comparison of anti-hyperlipidemic activity in ashwagandharishta prepared by traditional and modern methods. *Asian Journal of Research in Chemistry* 2010;3:574-7.
22. Mishra LC, Singh BB, Dagenais S. Scientific basis for the therapeutic use of *Withania somnifera* (ashwagandha): a review. *Alternative medicine review* 2000;5:334-46.
23. Gupta SK, Jadhav S, Gohil D, Panigrahi GC, Kaushal RK, Gandhi K, et al. Safety, toxicity and pharmacokinetic assessment of oral Withaferin-A in mice. *Toxicology Reports* 2022.

24. Singh N, Yadav SS, Rao AS, Nandal A, Kumar S, Ganaie SA, et al. Review on anticancerous therapeutic potential of *Withania somnifera* (L.) Dunal. Journal of Ethnopharmacology. 2021;270:113704.
25. Malik F, Kumar A, Bhushan S, Khan S, Bhatia A, Suri KA, et al. Reactive oxygen species generation and mitochondrial dysfunction in the apoptotic cell death of human myeloid leukemia HL-60 cells by a dietary compound withaferin A with concomitant protection by N-acetyl cysteine. Apoptosis 2007;12:2115-33.
26. Rai M, Jogee PS, Agarkar G, Santos CA. Anticancer activities of *Withania somnifera*: Current research, formulations, and future perspectives. Pharmaceutical biology 2016;54:189-97.
27. Sehrawat A, Roy R, Pore SK, Hahm ER, Samanta SK, Singh KB, Kim SH, Singh K, Singh SV. Mitochondrial dysfunction in cancer chemoprevention by phytochemicals from dietary and medicinal plants. In Seminars in cancer biology 2017;47:147-153.
28. Mann JR, Backlund MG, DuBois RN. Mechanisms of disease: Inflammatory mediators and cancer prevention. Nature clinical practice Oncology 2005;2:202-10.
29. Prabu PC, Panchapakesan S and Raj CD. Acute and sub-acute oral toxicity assessment of the hydroalcoholic extract of *Withania somnifera* roots in Wistar rats. Phytotherapy Research 2013;27:1169-1178.
30. Patel SB, Rao NJ, Hingorani, LL. Safety assessment of *Withania somnifera* extract standardized for Withaferin A: Acute and sub-acute toxicity study. Journal of Ayurveda and integrative medicine 2016;7:30-37.