

VARIATION IN RENAL PARAMETERS IN DENGUE AND NON-DENGUE PATIENTS

Abstract

Background: Dengue virus infection (DVI) is a vector-transmitted illness that is widespread in tropical and subtropical **regions**. **DVI** has been linked to numerous renal completely such as proteinuria, glomerulonephritis, and acute kidney injury (AKI), which have been documented to occur concurrently with or immediately following acute dengue infection. The purpose of the study was the variation in renal parameters in dengue and non-dengue patients.

Methods: The retrospective study conducted at cooperative hospital **vadakara , kerala** Total 100 subjects (50 dengue patients and Non dengue **patients**). **All** dengue patients were identified by registration number using hospital record management system. Patients admitted with primary and confirmed diagnosis of DVI, irrespective of severity were included in this study. The investigation of the study was to compare urea, uric acid, creatinine, and serum electrolyte levels in dengue patients and non-dengue patients. The analysis was performed within 24 hours after collection with the help of ABBOTT ARCHITECT 4100 Analyzer.

Results: In this study 100 subjects were taken; 50 were selected as dengue patients and 50 were selected as non dengue patients. It has drastically expanded urea and creatinine in dengue patients compared with non dengue sufferers. Serum electrolytes showed slightly decreased in dengue patients compared with non-dengue sufferers

Conclusion: The observed change in creatinine and urea increases in dengue patients compared to non-dengue patients. Sodium, potassium and chloride are almost reduced in dengue patients compared to non-dengue patients. Therefore, we can predict acute kidney injury by evaluating the kidney characteristics of dengue patients.

Keywords: Acute kidney **injury** ,**DF** or DHF/DSS, thrombocytopenia.

1. INTRODUCTION

Dengue viral **infection(DVI)** is a mosquito-borne contamination especially normal in the tropics and subtropics and is taken into consideration a prime worldwide fitness hazard by means of the sector health enterprise (WHO). **approximately** one-third of the world's population is atrisk for dengue infection [1]. **the** global infection rate procedures **50-100million** each year

The dengue virus is an RNA virus from the genus Flaviviridae which is transmitted by the bite of Aedes aegypti mosquito. There are four serotypes of dengue virus; infection with one serotype produces lifelong immunity to that serotype, but the immunity lasts only a few months for other serotypes.

The dengue virus has an incubation length of **3-14[2] days**. DVI is manifested by way of a ramification of scientific displays such as asymptomatic contamination, undifferentiated fever, dengue fever (DF), dengue hemorrhagic fever (DHF), and existence-threatening dengue shock syndrome (DSS). Like other tropical infections, DVI is related to multiple organ dysfunction involving liver, muscular tissues, coronary heart, brain, and kidneys.[3],[4]

DVI has been related to a variety of renal manifestations which includes proteinuria, hematuria, glomerulonephritis, and acute kidney harm (AKI), which have been suggested at some stage in or quickly after acute dengue infection, The occurrence of the

above renal manifestations varies among **17% and 62%in** sufferers with DVI[5]

Acute Renal Failure (ARF) is a unprecedented however nicely known problem of Dengue **infection** . **that** is a retrospective take a look at on versions in one of a kind renal parameters in dengue patients.[6]

2. MATERIALS AND METHODS

This was a retrospective study with a cross-sectional design. Study conducted at **vadakara,cooperative,hospital,vadakara**. The data was analyzed between October 2023 to December **2023.Total** 100 **subjects(50** Dengue patients and 50 Non-dengue patients).

Inclusion criteria; Patients of every age group were included. Both male and female who are examined Suspected DVI cases were diagnosed by using at least one of the following criteria: 1. Positive reverse transcriptase polymerase chain reaction. **2.Presence** of dengue immunoglobulin M and G antibodies in acute-phase serum by enzyme-linked immunosorbent assay. The serum samples were also tested for dengue-specific NS1 **antigen.The** analysis was performed within 24 hours after collection with the help of ABBOTT ARCHITECT 4100 Analyzer.

Statistical Analysis

The collected data were summarized by using the Descriptive Statistics: frequency, mean and S.D. The Independent sample "t" test was used to compare urea, uric acid, creatinine and serum electrolytes between dengue and non-dengue cases. The "t" test was used to compare **urea ,uric acid ,creatinine** and serum electrolytes according to gender. The p value < 0.05 was considered as significant. Data were analyzed by using the SPSS software (SPSS Inc.; Chicago, IL) version 29.0.10.

1. RESULTS AND DISCUSSION

Table – 1: Level of **urea ,uric acid** ,creatinine and serum electrolytes in dengue and non dengue patients

PARAMETER	DENGUE PATIENT	NON-DENGUE PATIENTS
SODIUM (mEq/l)	130.45	139.05
POTASSIUM (mEq/l)	3.27	3.98
CHLORIDE (mEq/l)	92.88	102.06
UREA (mg/dl)	68.78	23.14
URIC ACID (mg/dl)	5.48	5.44
CREATININE (mg/dl)	2.41	0.83

Table 1 shows that sodium, potassium and chloride are slightly decreased in dengue patients compared to non-dengue patients. Urea and creatinine levels are significantly increased in dengue patients compared to non-dengue patients. The levels of uric acid did not show a significant change between the two groups..

Table 2: Comparison of urea, uric acid, creatinine and serum electrolytes between dengue and non-dengue cases

		Mean	S.D.	"t"	p value
Urea (mg/dl)	Dengue patients	68.78	35.70	7.43	< 0.001*
	Non Dengue patients	23.14	6.95		
Creatinine (mg/dl)	Dengue patients	2.41	1.78	5.64	< 0.001*
	Non Dengue patients	0.83	0.20		
Uric acid (mg/dl)	Dengue patients	5.48	2.26	0.08	0.940
	Non Dengue patients	5.44	1.14		
Sodium (mEq/l)	Dengue patients	130.45	4.16	-11.82	< 0.001*
	Non Dengue patients	139.05	2.39		
Potassium (mEq/l)	Dengue patients	3.27	0.46	-7.28	< 0.001*
	Non Dengue patients	3.98	0.43		
Chloride (mEq/l)	Dengue patients	92.88	4.54	-9.43	< 0.001*
	Non Dengue patients	102.06	2.85		

("t" = Independent sample "t" test; * Significant)

The Independent sample "t" test was used to compare urea, uric acid, creatinine and serum electrolytes between dengue and non-dengue cases. There was a difference ($p < 0.05$) in urea, creatinine and serum electrolytes between dengue and non-dengue cases. [Table – 2]

Table 3: Comparison of urea, uric acid, creatinine and serum electrolytes according to gender







		Mean	S.D.	"t"	p value
Urea (mg/dl)	Male	52.92	39.17	2.25	0.027*
	Female	33.96	17.62		
Creatinine (mg/dl)	Male	2.08	1.73	3.75	< 0.001*
	Female	0.90	0.37		
Uric acid (mg/dl)	Male	5.78	1.90	1.99	0.051
	Female	4.86	1.22		
Sodium (mEq/l)	Male	135.21	5.30	0.77	0.444
	Female	134.27	5.76		
Potassium (mEq/l)	Male	3.64	0.52	-0.09	0.930
	Female	3.65	0.64		
Chloride (mEq/l)	Male	97.94	5.68	-0.26	0.794
	Female	98.36	6.25		

("t" = Independent sample "t" test; * Significant)

The Independent sample "t" test was used to compare urea, uric acid, creatinine and serum electrolytes according to gender. There was a difference ($p < 0.05$) in urea as well as creatinine between males and females. [Table – 3]

Graph 1: Graph shows the comparison between urea ,creatinine ,uric acid in dengue and non dengue patients.

Graph 2 : Graph shows the comparison between serum electrolytes in dengue and non dengue patient



From the graph It is clearly seen that the mean increase in urea and creatinine is significantly higher in dengue patients than in non-dengue patients. Uric acid ranges confirmed that there was no significant trade between the two organizations.

Table 2 and graph number 1 definitely show that there was a difference ($p < 0.05$) between urea, creatinine in dengue cases..

The graph clearly shows that the mean sodium, potassium and chloride concentrations of dengue patients are slightly decreased compared to non-dengue patients.

Table 2 and graph number 2 clearly show that there was a difference ($p < 0.05$) in sodium, potassium and chloride between dengue and non-dengue cases

2. Discussion

The retrospective study evaluated various renal parameters in dengue patients compared to non-dengue patients. Out of 100 samples collected, 50 were from dengue patients and 50 were from non-dengue patients. Creatinine and urea levels were significantly higher in dengue patients than non-dengue patients. Sodium, potassium, and chloride levels were slightly lower in dengue patients compared to non-dengue patients. Uric acid levels did not significantly differ between dengue and non-dengue patients. A previous study on renal manifestations of dengue viral infections by Mahesh Eswarappa et al. (2019) found that AKI (acute kidney injury) is a complication of dengue **infection.** The findings of my study align with previous research published in the Journal of Clinical and Diagnostic Research on "Acute Renal Failure in Dengue Infection." That study found that acute renal failure in dengue infection is typically caused by extensive capillary leakage, hypotension, rhabdomyolysis, hemolysis, and severe disseminated intravascular coagulation. These factors lead to hypoxia, ischemia, and multiple organ dysfunction, as noted across various case reports. In some patients, acute tubular necrosis with interstitial edema and mononuclear cell infiltration can also

contribute to renal failure. The exact mechanisms by which dengue virus directly damages the kidneys, even in dengue fever patients without bleeding or hypotension, remains unclear. Similarly, a study by Karlo J Lizarragal and Ali Nayer on "Dengue- Associated Kidney Disease" found proteinuria in up to 74% of patients with dengue hemorrhagic fever. They described a 22-year-old woman with confirmed dengue infection who initially had a serum creatinine of 1.0 mg/dL.

3. Conclusion

The study examined the differences in renal parameters between dengue patients and non- dengue patients. It was observed that creatinine and urea levels were higher in dengue patients compared to non-dengue patients. sodium, potassium, and chloride levels were slightly lower in dengue patients than in non-dengue patients. Uric acid levels showed no significant change between the two groups. In conclusion, the study suggests that renal complications from dengue infections are generally mild. However, some patients may experience more severe renal damage, potentially resulting in acute renal failure. This highlights acute renal failure as a possible complication of dengue infections. By assessing renal function tests in **dengue**

patients, it may be possible to predict the development of acute kidney injury.

Informed consent

Written informed consent was taken from each participant before enrollment.

References

1. Guzmán MG, Kouri G. Dengue: An update. *Lancet Infect Dis* 2002;**2:33-42**.
2. World Health Organization. Dengue and Severe Dengue. World Health Organization. **3.Udwadia** FE. Multiple organ dysfunction syndrome due to tropical infections. *Indian J Crit Care Med* 2003;**7:233-6**.
4. Halstead SB. Dengue. *Lancet* 2007;**370:1644-52**.
5. Lizarraga KJ, Nayer A. Dengue-associated kidney disease. *J Nephrol* 2014;**3:57-62**.
6. Gubler D.J.: Dengue and dengue hemorrhagic fever. *Clin Microbiol Rev* 1998; 11: pp. 480.
7. Gibbons R.V., Vaughn D.W.: Dengue: an escalating problem. *Br Med J* 2002; 324: pp. 1563-1566.
8. Balmaseda A., Hammond S., Perez L., et. al.: Serotype-specific differences in clinical manifestations of dengue. *Am J Trop Med Hyg* 2006; 74: pp. 449-456.
9. Swaminathan and N. Khanna, "Experimental dengue vaccines," *Molecular Vaccines: From Prophylaxis to Therapy*, vol.1, pp. 135-151, 2013.
10. Bhatt, S., et al., The global distribution and burden of dengue. *Nature*, 2013.**496(7446)**: p. 504-507.
11. Siler J. F., Hall M. W., Hitchens **A.Dengue**, its history, epidemiology, mechanism of transmission, etiology, clinical manifestations, immunity and prevention. *Philipp. J. Sci.* 29 1926 1 **304** Google Scholar
12. Gubler D. J., Suharyono W., Tan R., Abidin M., Sie A. Viremia in patients with naturally acquired dengue infection. *Bull. W. H. O.* 59 1981 623 630.
13. Whitchorn J, Simmons CP. The pathogenesis of dengue. *Vaccine*. 2011;**29:7221-8**.
14. Guzman MG, Kouri G. Dengue and dengue hemorrhagic fever in the Americas: Lessons and challenges. *J Clin Virol*. 2003;**27:1-13**.
15. Revised and Expanded ed. New Delhi: WHO; 2011. World Health Organization. Comprehensive Guidelines for Prevention and Control of Dengue and Dengue Haemorrhagic Fever 16. Thomas EA, John M, Bhatia A. Mucocutaneous manifestations of dengue viral infection in Punjab. *Int J Dermatol*. 2007;**46:715-19**.
16. Arshad I, Malik FA, Hussain A, Shah SA. Dengue fever: Clinico-pathologic

correlations and their association with poor **outcome.**

Professional Med J.

17. Wu SJ, Grouard-Vigel G, Sun W, Mascola JR, Brachel E, Putvatana R, et al. Human skin langerhans cells are targets of dengue virus infection. *Nat Med.* 2000;6:816-20. 18. Bhamarapravati N. Pathology and Pathogenesis of DHIF. New Delhi: **WHOMeeting**; 1980.

19. Thomas SJ, et al. Dengue virus infection: Clinical manifestations and diagnosis.

20. Tjaden, N.B., et al., Extrinsic Incubation Period of Dengue:

Knowledge, Backlog, and Applications of Temperature Dependence. *Plos Neglected Tropical Diseases*, 2013. 7(6): p. 5.

21. Siler, J.F., M.W. Hall, and A.P. Hitchens, Dengue: Its history, epidemiology, mechanism of transmission, etiology, **clinical manifestations**, immunity and prevention. 1926, Manila: Bureau of Science.

22. Watts, D.M., et al., Effect of Temperature on the vector efficiency of *Aedes aegypti* for Dengue 2 virus. *American Journal of Tropical Medicine and Hygiene*, 1987.36(1): p. 143- 152.

23. Carington, **I.B,** et al, Fluctuations at low mean temperatures accelerate Dengue virus transmission by *Aedes aegypti*. *PLoS Neglected Tropical Diseases*, 2013. 7(4): p. e2190. **26 Lambrechts**, L, et al, Impact of daily temperature fluctuations on dengue virus transmission by *Aedes aegypti*. *Proceedings of the National Academy of Sciences of the United States of America*, 2011.108(18): p. 7460-7465.

24. Ye, Y.X.H., et al., Wolbachia Reduces the Transmission Potential of Dengue-Infected *Aedes aegypti*. *PLOS Neglected Tropical Diseases*, 2015. 9(6): p. e0003894.

25. Duong, V., et al., Asymptomatic humans transmit dengue virus to mosquitoes. *Proceedings of the National Academy of Sciences of the USA*, 2015. 112(47): p. 14688- 14693.

26. Nguyen, N.M., et al., Host and viral features of human dengue cases shape the population of infected and infectious *Aedes aegypti* mosquitoes. *Proceedings of the National Academy of Sciences of the United States of America*, 2013. **110(22): p.9072- 9077.**

27. Gubler, D.J., et al., Viraemia in patients with naturally acquired dengue infection. *Bulletin of the World Health Organisation*, 1981.

59: P. 623-630 28. Wilder-Smith A, Renhorn KE, Tissera H, Abu Bakar S, Alphey L, Kittayapong P, Lindsay S, Logan J, Hatz C, Reiter P, Rocklov J, Byass P, Louis VR, Tozan Y, Massad E, Tenorio A, Lagneau C, L'Ambert G, Brooks D, Wegerdt J, Gubler D. 2012. Dengue Tools: innovative tools

29. Lanciotti RS, Calisher CH, Gubler DJ, Chang GJ, Vorndam AV.

1992. Rapid detection and typing of dengue viruses from clinical samples by using reverse **transcriptase-polymerase chain reaction. J Clin Microbiol 30:545-551.**

30. Seah CL, Chow VT, Tan HC, Can YC. 1995. Rapid, single-step RT-PCR typing of dengue viruses using five NS3 gene primers. J Virol Methods 51:193-200.