

## Original Research Article

### **Growth rate and nutrient uptake of *Stevia Rebaudiana* Bertoni influenced by organic manures in laterite soils of Kerala, India**

**Abstract:** A field experiment was conducted to understand the effect of nutrition through organic sources on the crop growth rate (CGR), relative growth rates (RGR), and nutrient uptake of stevia (*Stevia rebaudiana* Bertoni) during April to August of 2022 and 2023 at the Agronomy farm, Department of Agronomy, College of Agriculture, Vellanikkara, Thrissur. The experiment was laid out in randomized block design (RBD) with 7 treatments replicated thrice which consisted of different levels of organic manures viz., three levels of farmyard manure (5, 10, 15 t ha<sup>-1</sup>), three levels of vermicompost (2.5, 5, 7.5 kg ha<sup>-1</sup>) and a control. The organic manure treatments significantly affected crop growth rate (CGR) and relative growth rate (RGR) in all the growth periods. The two-year pooled mean of CGR of stevia was significantly superior in FYM@15 t ha<sup>-1</sup> over control in all the three growth periods i.e., at 0-30 DAP, 30-45 DAP and 45 DAP to harvest. Whereas, the two-year mean of RGR of stevia was found statistically higher in FYM@15 t ha<sup>-1</sup> over control in the first two growth periods only. The organic manure treatments had a significant influence on the plant nutrient (NPK) uptake as well in both the years. The two-year pooled mean data regarding plant uptake of nitrogen and phosphorus was found significantly higher in FYM@15 t ha<sup>-1</sup> over control whereas statistically higher pooled mean potassium uptake by stevia was found in VC@7.5 kg ha<sup>-1</sup> over control.

*Keywords:* *Stevia*, *farmyard manure*, *vermicompost*, *growth rate*, *nutrient uptake*

#### **1. Introduction:**

*Stevia rebaudiana* Bertoni has emerged as a leading natural sweetener, renowned for its exceptional ability to deliver high sweetness without the calories associated with traditional sugars. This perennial herb, native to South America, has been utilized for centuries by indigenous communities for its sweet leaves (Brandle & Rosa, 1992). The main sweetening compounds in *Stevia rebaudiana* Bertoni are stevioside, rebaudioside-A, rebaudioside-C, and dulcoside-A, with their relative sweetness compared to sucrose being 210, 242, 30, and 30 times sweeter, respectively (Kingham, 1987). Its global appeal has grown significantly following the approval of stevioside and other steviol glycosides in key markets, including the United States and the European Union (Risso *et al.*, 2014). As consumers increasingly seek healthier alternatives to sugar and artificial sweeteners, Stevia offers a compelling option, being not only low in calories but also rich in antioxidants and beneficial phytochemicals. Research has consistently demonstrated that stevia leaf products are safe for individuals with diabetes, high blood pressure, and obesity, potentially aiding in the management or prevention of complications linked to these conditions (Carrera-Lanestosa *et al.*, 2017). Furthermore, studies have confirmed that stevia is safe for children as well (Aguero *et al.*, 2014). The popularity of stevia, driven by its pleasant taste and zero-calorie natural sweetener status, has encouraged Indian farmers to commercially cultivate it. As a result, stevia is now successfully grown in several Indian states, including Rajasthan, Maharashtra, Punjab, and Orissa (Goyal *et al.*, 2010).

The cultivation of *Stevia rebaudiana* Bertoni is influenced by environmental conditions and agricultural practices, especially nutrient management, which enhances growth and yield. As the crop is highly adaptable to the poor-quality Paraguayan soils, which is its natural habitat, stevia requires a low to moderate level of nutrition (Goenadi, 1987). However, commercial cultivation of Stevia needs exogenous fertilization to achieve sustainable yield (Rashid *et al.*, 2013). Although the specific nutrient requirements of Stevia, particularly in the context of South Indian soils are still not well understood, the RDF is estimated at 60:30:45 kg NPK ha<sup>-1</sup> for Bengaluru conditions (Farooqi and Sreeramu, 2004). With the global population projected to reach 9.2 billion by 2050 (Gruhn *et al.*, 2000), a shift to sustainable agricultural practices is essential. Moreover, indiscriminate use of inorganic fertilizers in modern farming, especially in medicinal crops, can leave residues in the medicinal preparations that can harm human health and the environment hence an urgent transition towards safe and sustainable organic farming practices is necessary (Umesha *et al.*, 2011). There has been limited research on supplementing inorganic nutrient sources with organics without compromising the productivity and quality of Stevia, prompting this study on organic nutrition. Organic manures like farmyard manure and vermicompost provide viable, low-input alternatives, supplying essential nutrients and improving soil structure and microbial activity for healthier crop growth.

The crop growth rate and relative growth rates of a crop are vital indicators for assessing the viability and productivity of its cultivation. These metrics provide valuable insights into how well the plant adapts to varying agricultural practices and environmental conditions, essential for maximizing yield. Moreover, analysing growth rates at various developmental stages provides deeper insights into the plant's nutritional requirements and its ability to withstand abiotic stress, which is essential for effective crop management. A comprehensive understanding of nutrient uptake is essential for developing effective fertilization strategies that align nutrient availability with the specific needs of *Stevia rebaudiana* (Angelini & Tavarini, 2014). Key nutrients such as nitrogen (N), phosphorus (P), and potassium (K) are crucial for physiological processes, including photosynthesis, energy transfer, and overall plant growth. Additionally, a thorough understanding of NPK dynamics in plants can help identify specific nutrient deficiencies or imbalances, facilitating the optimization of fertilization practices for enhanced plant health and productivity.

Against this backdrop, this research was carried out to explore the growth rates and nutrient uptake of *Stevia rebaudiana* as influenced by organic manures, providing insights to optimize sustainable nutrient management.

## **2. Materials and methods:**

The field experiment entitled 'Effect of organic manures on growth, yield and quality of *Stevia rebaudiana* B.' was conducted at the Agronomy farm, Department of Agronomy, College of Agriculture, Vellanikkara, Thrissur, Kerala from April to August of 2022 and 2023. The site is located at 13° 32' N latitude and 76° 26' E longitude, at an altitude of 40 m above mean sea level and the annual rainfall in 2022 was 3128.3 mm and 2697.3 mm in 2023. The texture of the experimental field was sandy clay loam and was acidic in reaction with pH 4.68, EC (0.07 ds m<sup>-1</sup>, 1.33 % OC, available N (146 kg ha<sup>-1</sup>), P (33 kg ha<sup>-1</sup>) and K (188 kg ha<sup>-1</sup>). The experiment was laid out in randomized block design (RBD) with 7 treatments replicated thrice which consisted of different levels of organic manures *viz.*, three levels of farmyard manure, three levels of vermicompost and a control. The treatments were T<sub>1</sub>-no manure; T<sub>2</sub>- FYM@ 5 t ha<sup>-1</sup>; T<sub>3</sub>- FYM@ 10 t ha<sup>-1</sup>; T<sub>4</sub>- FYM@ 15 t ha<sup>-1</sup>; T<sub>5</sub>- VC@ 2.5 t ha<sup>-1</sup>; T<sub>6</sub>- VC @ 2.5 t ha<sup>-1</sup>; and T<sub>7</sub>- VC @ 2.5 t ha<sup>-1</sup>. After thorough field preparation by ploughing and discing, beds of size 3m x 3m were prepared and mulched with

30  $\mu$  plastic mulch. A green colour shade net with 25 per cent sunlight permeability was used to provide artificial shading over the whole experimental area. The individual beds were fertilized according to the treatments using well-decomposed farmyard manure and vermicompost during preparation, before mulching as a basal application two weeks before transplantation. One-month-old rooted stem cuttings with 4-5 nodes raised in the nursery were transplanted uniformly in all the beds at a spacing of 30cm x 30cm in both years. Regular intercultural operations such as irrigation, weeding etc. were carried out during the experimental period and a single harvest was done in both years 110 days after planting (DAP) by uprooting the plants.

### 2.1 Crop growth rate (CGR)

Crop growth rate illustrates the dry matter accumulation by the crop per unit land area per unit time and it was estimated at 0-30 DAP, 30-45 DAP and at 45 DAP to harvest in both the years. CGR was determined using the formula provided by Watson (1952) and expressed as grams of dry matter produced per day.

$$\text{CGR}(\text{gm}^{-2}\text{day}^{-1}) = \frac{W_2 - W_1}{A(t_2 - t_1)}$$

Where  $W_1$  and  $W_2$  are dry weights of plants at time  $t_1$  and  $t_2$ , respectively,  $A$  is the land area.

### 2.2 Relative growth rate (RGR)

Relative growth rate explains the rate of growth of the crop per unit of dry matter per unit of time and it was calculated at 0-30 DAP, 30-45 DAP and 45 DAP to harvest in both years. RGR was determined using the formula proposed by Blackman (1919) and expressed as grams of dry matter produced by a gram of existing dry matter in a day.

$$\text{RGR}(\text{g g}^{-1}\text{day}^{-1}) = \frac{\log_e W_2 - \log_e W_1}{t_2 - t_1}$$

Where  $W_1$  and  $W_2$  are dry weights of plants at time  $t_1$  and  $t_2$ , respectively.

### 2.3 Plant uptake of N, P and K at harvest

The replication-wise collected plant samples were cleaned, shade-dried and dried further in a hot air oven at  $65 \pm 5^\circ\text{C}$  to achieve a constant weight and ground to fine powder. These powdered samples were used to analyse plant NPK content by following the standard procedures. The nutrient uptake was then determined by multiplying dry matter production and nutrient content in per cent and expressed as  $\text{kg ha}^{-1}$ .

## 3. Results and discussion:

### 3.1 Crop growth rate ( $\text{g m}^{-2}\text{day}^{-1}$ )

Crop growth rates (CGR) of stevia at three different growth stages in two years and the pooled mean are presented in Table 1. In the first two growth periods of both the years (2022 and 2023), *i.e.*, from 0-30 DAP and 30-45 DAP, the organic manures had a significant effect on the crop growth rate of stevia whereas, in the pooled mean, the CGR was significantly

affected by the organic manures in all the three growth periods. Despite the treatments, a steep increase in the CGR was observed from the first to the second growth period, whereas a steady increase was observed from the second to third growth period in both years. At 0-30 DAP, the two-year pooled mean of CGR was found significantly higher in FYM@15 t ha<sup>-1</sup> over control and was on par with FYM@10 t ha<sup>-1</sup> followed by VC@7.5 t ha<sup>-1</sup>. Whereas, at 30-45 DAP, significantly highest pooled mean of CGR was found in FYM@15 t ha<sup>-1</sup> over control followed by FYM@10 t ha<sup>-1</sup> and VC@7.5 t ha<sup>-1</sup>. However, at 45 DAP to harvest, the two-year mean data of CGR was found significantly higher in FYM@15 t ha<sup>-1</sup> over control and was on par with VC@7.5 t ha<sup>-1</sup>, VC@5 t ha<sup>-1</sup> and FYM@10 t ha<sup>-1</sup> treatments. The crop growth rate of stevia was found to be increasing with increasing quantity of manure (both farmyard manure and vermicompost) applied. This can be attributed to the improved soil physical chemical and biological properties by the application of well-decomposed organic manures. Better soil aeration, microbial growth and enhanced nutrient availability might have caused better root proliferation and establishment and nutrient absorption thereby increasing the growth parameters and leading to better herbage production and growth rate (Kumar *et al.*, 2023).

**Table 1. Effect of organic manures on the crop growth rate of *stevia rebaudiana* Bertoni at (0-30 DAP), (30-45 DAP) and (45 DAP – Harvest)**

CGR(g m <sup>-2</sup> day <sup>-1</sup> )	2022			2023			Pooled mean		
	0-30 DAP	30-45 DAP	45DAP-Harvest	0-30 DAP	30-45 DAP	45DAP-Harvest	0-30 DAP	30-45 DAP	45DAP-Harvest
No manure	0.231	1.188	1.989	0.271	1.389	2.066	0.260	1.316	2.121
FYM@ 5 t ha <sup>-1</sup>	0.286	1.573	2.334	0.364	1.735	2.458	0.325	1.654	2.396
FYM@ 10 t ha <sup>-1</sup>	0.357	2.705	2.585	0.415	2.767	2.866	0.386	2.736	2.725
FYM@15 t ha <sup>-1</sup>	0.381	2.883	2.582	0.460	3.030	3.006	0.421	2.957	2.794
VC@ 2.5 t ha <sup>-1</sup>	0.282	1.653	2.359	0.331	1.616	2.615	0.307	1.635	2.487
VC@ 5 t ha <sup>-1</sup>	0.304	1.630	2.705	0.390	1.852	2.823	0.347	1.741	2.764
VC@ 7.5 t ha <sup>-1</sup>	0.339	2.068	2.708	0.408	2.260	2.823	0.374	2.164	2.766
CD (0.05)	0.058	0.449	NS	0.087	0.606	NS	0.038	0.152	0.291
CV (%)	10.39	12.89	12.31	12.94	13.27	13.23	7.88	5.29	8.00

### 3.2 Relative growth rate (g g<sup>-1</sup> day<sup>-1</sup>)

Relative growth rates (RGR) of stevia at three different growth stages in two years and the pooled mean are presented in Table 2. In the first and third growth periods of 2022 *i.e.*, at 0-30 DAP and 45 DAP to harvest, the organic manures had a significant effect on the relative growth rate of stevia whereas, in 2023, the RGR was not significantly affected by the treatments in all the growth periods except 0-30 DAP. However, the pooled mean of the RGR was significantly affected by the organic manures in all three growth periods. The two-year mean RGR was found significantly higher in FYM@15 t ha<sup>-1</sup> over control and was on par with FYM@10 t ha<sup>-1</sup> followed by VC@7.5 t ha<sup>-1</sup> at 0-30 Dap and 30-45 DAP growth periods. However, the pooled mean of RGR was found significantly higher in control and stood on par with FYM@5 t ha<sup>-1</sup>, VC@2.5 t ha<sup>-1</sup>, and VC@5 t ha<sup>-1</sup>. Irrespective of the treatments, a slight increase of RGR was observed in all the treatments from the first to the second growth period and a steep decrease was found from the second to the third growth stage in both years. The trend in relative growth rate can be viewed as normal, as biomass production efficiency may decline over time due to plant senescence and an increase in the growth of non-photosynthetic woody tissues (Jarma-Orozco *et al.*, 2020).

### 3.3 Plant nutrient uptake (NPK in kg ha<sup>-1</sup>)

Majornutrient uptake (NPK) by stevia was significantly influenced by the organic manure treatments in both the years, 2022 and 2023 and the data is presented in Table 3. Two-year pooled mean data regarding uptake of nitrogen was found significantly higher in FYM@15 t ha<sup>-1</sup> over control but was on par with FYM@10 t ha<sup>-1</sup>, VC@7.5 t ha<sup>-1</sup>, and VC@5 t ha<sup>-1</sup> treatments. Plots treated with either farmyard manure or vermicompost, alone or in combination with apple pomace, showed a greater accumulation of nitrogen in the soil increasing N availability and thereby higher N uptake by plants, likely due to the slow mineralization of N from the manures, as noted by Anwar *et al.* (2005). The two-year pooled mean uptake of phosphorous by stevia was found significantly greater in FYM@15 t ha<sup>-1</sup> over control and was on par with FYM@10 t ha<sup>-1</sup> treatment. The increased concentration of phosphorus in stevia from organic manures attributing to higher P uptake may result from enhanced solubilization of P, either through the activation of microorganisms that excrete organic acids (Suba Rao, 1982) or due to increased phosphatase activity (Sainz *et al.*, 1998). When it comes to the pooled mean of potassium uptake by stevia, a significantly higher quantity was found in VC@7.5 t ha<sup>-1</sup> over control and was on par with FYM@15 t ha<sup>-1</sup> treatment. The increase in potassium uptake by stevia from vermicompost treatments may be attributed to improved K availability, which shifts the equilibrium in the soil from relatively exchangeable K to more soluble forms (Basker *et al.*, 1992). It was noted that the uptake of nitrogen and potassium exceeded that of phosphorus and these findings were in line with those of Kumar *et al.* (2013). The higher uptake of NPK across various treatment combinations can be attributed to the combined effects of increased nutrient concentrations and dry matter yield associated with these treatments. Similar patterns have been documented in previous studies involving basil (Anwar *et al.*, 2005), mint (Patra *et al.*, 2000), and wheat (Dudhat *et al.*, 1997).

RGR(g g <sup>-1</sup> day <sup>-1</sup> )	2022			2023			Pooled mean		
	Treatments	0-30 DAP	30-45 DAP	45DAP-Harvest	0-30 DAP	30-45 DAP	45DAP-Harvest	0-30 DAP	30-45 DAP
No manure	0.029	0.034	0.013	0.031	0.034	0.012	0.030	0.033	0.012
FYM@ 5 t ha <sup>-1</sup>	0.031	0.036	0.012	0.034	0.033	0.011	0.033	0.034	0.012
FYM@ 10 t ha <sup>-1</sup>	0.034	0.043	0.010	0.036	0.041	0.010	0.035	0.042	0.010
FYM@15 t ha <sup>-1</sup>	0.035	0.043	0.010	0.038	0.041	0.010	0.036	0.042	0.010
VC@ 2.5 t ha <sup>-1</sup>	0.031	0.037	0.012	0.033	0.034	0.012	0.032	0.035	0.012
VC@ 5 t ha <sup>-1</sup>	0.032	0.035	0.013	0.035	0.033	0.012	0.034	0.034	0.012
VC@ 7.5 t ha <sup>-1</sup>	0.034	0.038	0.011	0.036	0.037	0.011	0.035	0.037	0.011
CD (0.05)	0.002	NS	0.002	0.003	NS	NS	0.001	0.002	0.001
CV (%)	3.94	10.47	9.40	4.66	13.15	11.05	2.54	4.61	5.94

**Table 2. Effect of organic manures on the relative growth rate of *stevia rebaudiana* Bertoni at (0-30 DAP), (30-45 DAP) and (45 DAP – Harvest)**

Uptake (kg ha <sup>-1</sup> ) Treatments	N			P			K		
	2022	2023	Pooled	2022	2023	Pooled	2022	2023	Pooled
No manure	15.39	16.64	18.41	3.19	3.54	3.57	24.82	29.12	29.07
FYM@ 5 t ha <sup>-1</sup>	28.08	30.48	29.28	4.16	5.01	4.59	35.36	39.71	37.53
FYM@ 10 t ha <sup>-1</sup>	34.65	38.14	36.39	5.19	6.58	5.89	44.60	55.39	50.00
FYM@15 t ha <sup>-1</sup>	35.82	41.16	38.49	5.90	7.46	6.68	52.05	62.35	57.20
VC@ 2.5 t ha <sup>-1</sup>	27.79	30.37	29.081	3.87	4.61	4.24	43.10	49.01	46.06
VC@ 5 t ha <sup>-1</sup>	32.71	35.28	33.99	4.73	5.67	5.20	51.42	55.64	53.53
VC@ 7.5 t ha <sup>-1</sup>	34.20	36.61	35.41	5.12	5.85	5.49	57.55	64.78	61.16
CD (0.05)	4.777	5.652	4.735	0.742	0.959	0.906	7.227	8.962	6.455
CV (%)	9.009	9.725	10.613	9.075	9.744	12.595	9.206	9.906	9.56

**Table 3. Effect of organic manures on the plant nutrient uptake of *stevia rebaudiana* Bertoni in 2022, 2023 and pooled mean in kg ha<sup>-1</sup>**

### Conclusion:

The current study indicates that applying organic manure in the form of farmyard manure at the rate of 15 t ha<sup>-1</sup> results in better overall growth of stevia as evidenced by the crop and relative growth rates. Along with the improved plant growth, the crop's nutrient uptake also showed significant improvement with the application of organic manures in higher quantities.

### References:

- Aguero, S.D., Onate, G., Rivera, H.P. Consumption of nonnutritive sweeteners and nutritional status in 10-16- year-old students. *Arch Argent Pediatr.* 2014;112:207– 14.
- Angelini, L. G., & Tavarini, S. (2014). Crop productivity, steviol glycoside yield, nutrient concentration and uptake of *Stevia rebaudiana* Bert. under Mediterranean field conditions. *Communications in soil science and plant analysis*, 45(19), 2577-2592.
- Anwar, M., Patra, D. D., Chand, S., Alpesh, K., Naqvi, A. A., & Khanuja, S. P. S. (2005). Effect of organic manures and inorganic fertilizer on growth, herb and oil yield, nutrient accumulation, and oil quality of French basil. *Communications in soil science and plant analysis*, 36(13-14), 1737-1746.
- Basker, A., Macgregor, A. N., & Kirkman, J. H. (1992). Influence of soil ingestion by earthworms on the availability of potassium in soil: An incubation experiment. *Biology and Fertility of Soils*, 14, 300-303.
- Blackman, V.H. 1919. The compound interest law and plant growth. *Ann. Bot.* 33(3):353-360.
- Brandle, J.E. and Rosa, N. 1992. Heritability for yield, leaf: stem ratio and stevioside content estimated from a landrace cultivar of *Stevia rebaudiana*. *Can. J. Plant Sci.* 72: 1263-1266.
- Carrera-Lanestosa A, Moguel-Ordóñez Y, SeguraCampos M. *Stevia rebaudiana* Bertoni: A natural alternative for treating diseases associated with metabolic syndrome. *J Med Food.* 2017;20:933–43.

- Dudhat, M. S., Malavia, D. D., Mathukia, R. K., & Khanpara, V. D. 1997. Effect of nutrient management through organic and inorganic sources on growth, yield, quality and nutrients uptake by wheat (*Triticum aestivum*). *Indian Journal of Agronomy*, 1997, 42(3), 455-458.
- Farooqi, A. A., & Sreeramu, B. S. 2004. Cultivation of medicinal and aromatic crops. *Universities Press*.
- Goenadi, D. H. 1987. Effect of slope position on the growth of Stevia in Indonesia. *Communications in soil science and plant analysis*, 18(11), 1317-1328.
- Goyal, S.K., Samsheer, and Goyal, R.K. 2010. Stevia (*Stevia rebaudiana*) a biosweetener. *Int. J. Food Sci. Nutri.* 61: 1-10.
- Gruhn, P., Golett, F. and Yudelman, M. 2000. Integrated Nutrient Management, Soil Fertility and Sustainable Agriculture: Current Issues and Future Challenges. Food, Agriculture and Environment Discussion Paper 32. International Food Policy Research Institute, Washington DC, USA.
- Jarma-Orozco, A., Combatt-Caballero, E., & Jaraba-Navas, J. 2020. Growth and development of *Stevia rebaudiana* Bert., in high and low levels of radiation. *Current plant biology*, 22, 100144.
- Kinghorn, A.D. 1987. Biologically Active Compounds from Plants with Reputed Medicinal and Sweetening Properties. *Journal National Production*, 50, 1009-1024.
- Kumar, P. M., Roy, D. K., Singh, S. K., & Kumar, M. 2023. Influence of Different Tillage and Nutrient Management Practices on Growth Performance of Direct Seeded Rice. *Int. J. Environ. Clim. Change*, 13(8), 1432-1439.
- Kumar, R., Sharma, S., & Prasad, R. 2013. Yield, nutrient uptake, and quality of stevia as affected by organic sources of nutrient. *Communications in soil science and plant analysis*, 44(21), 3137-3149.
- Patra, D. D., Anwar, M., & Chand, S. 2000. Integrated nutrient management and waste recycling for restoring soil fertility and productivity in Japanese mint and mustard sequence in Uttar Pradesh, India. *Agriculture, ecosystems & environment*, 80(3), 267-275.
- Rashid, Z., Rashid, M., Inamullah, S., Rassol, S., Bahar, F.A. 2013. Effect of different levels of farmyard manure and nitrogen on the yield and nitrogen uptake by stevia (*Stevia rebaudiana* Bertoni). *African J. Agric. Res.* 8 (29), 3941-3945.
- Risso, Davide, Gabriella Morini, Luca Pagani, Andrea Quagliariello, Cristina Giuliani, Sara De Fanti, Marco Sazzini, Donata Luiselli, and Sergio Tofanelli. 2014. "Genetic signature of differential sensitivity to stevioside in the Italian population." *Genes & nutrition*, 9 (2014): 1-9.
- Sainz, M. J., Taboada-Castro, M. T., & Vilarino, A. 1998. Growth, mineral nutrition and mycorrhizal colonization of red clover and cucumber plants grown in a soil amended with composted urban wastes. *Plant and soil*, 205, 85-92.
- Subba Rao, N. S. 1982. Utilization of farm wastes and residues in agriculture. *Advances in agricultural microbiology*, 1982, 509-521.

Umesha, K., Smitha, G. R., Sreeramu, B. S., & Waman, A. A. 2011. Organic manures and bio-fertilizers effectively improve yield and quality of stevia (*Stevia rebaudiana*). *Journal of Applied Horticulture*, 13(2), 157-162.

Watson, D.J. 1952. The physiological basis of variation in yield. *Adv. Agron.* 4: 101-145.

UNDER PEER REVIEW