

Cell Therapy in Rabbits: Advances, Applications and Perspectives in Regenerative Medicine

ABSTRACT

Cell therapy represents an innovative and promising approach to treating debilitating conditions, and animal testing on rabbits has played a crucial role in advancing this therapy. The objective of this manuscript aims to review the literature on the use of lagomorphs in cell therapy studies. The Pubmed and Science Direct databases were used, with the descriptors: Cell therapy and experimentation in rabbits; rabbits and cell therapy. Of the articles found, those that deviated from the topic or did not address the use of rabbits in cell therapy were excluded. The articles that addressed the topic were included after reading them completely. The present work presents a broad and distinct view of cell therapy and its different applicability, presenting a promising framework for animal experimentation in rabbits, and the advantages of using this animal model.

Keywords: Autologous Therapy; Allogeneic therapy; Debilitating Conditions; Mesenchymal Stem Cells; Tissue Engineering.

1. INTRODUCTION

Cell therapy constitutes a branch of regenerative medicine that aims to utilize cells, often derived from the patient themselves, to treat various pathologies [1]. Animal experimentation in rabbits has played a crucial role in the advancement of this therapy. Based on the intrinsic capacity of cells to self-renew and differentiate into multiple cell types, cell therapy represents an innovative and promising approach for the treatment of debilitating conditions [2]. Notable examples include its application in detrusor hypocontractility [3], the reduction of the inflammatory process associated with osteoarthritis [4], modulation of tissue repair in injured vocal folds [5], in tissue engineering for blood vessels [6], in the treatment of neurological sequelae resulting from distemper [7], in rare genetic disorders [8], in ligament reconstruction [9], in the treatment of spinal cord injuries [10], and in the management of multiple myeloma [11]. These interventions have the potential to significantly improve patients' quality of life, as damaged or dysfunctional cells can be replaced by healthy ones, thereby restoring the function of affected tissues and organs.

Additionally, cell therapy can be classified into two types based on the origin of the cells: autologous, when obtained from the patient themselves, and allogenic, when obtained from a donor [12]. Autologous cell therapy represents a remarkable advancement in the field of regenerative medicine and advanced therapies, with the primary advantage of eliminating the risk of immune rejection, as the implanted cells are genetically identical to those of the

recipient. In contrast, allogenic therapy employs cells derived from a donor genetically distinct from the recipient, offering substantial advantages, particularly in situations where autologous cells are unfeasible for therapeutic purposes due to pathologies or genetic defects [13]. This procedure has paved the way for the development of treatments for various hematological and immunological diseases. However, allogenic therapy introduces an increased risk of immune rejection and complications, such as graft-versus-host disease (GVHD) [14].


Moreover, animal experimentation has demonstrated extremely promising potential and is continuously expanding. Investigations focused on rabbits have shown significant potential in regenerative medicine, as these animals exhibit physiological and biological characteristics that are similar to those of humans, thereby rendering them relevant as experimental models for tissue regeneration research [15]. Additionally, this therapy represents an emerging frontier in biomedical research, with significant implications for regenerative medicine, addressing challenges such as the validity of data obtained when compared to other animals [16].

The European rabbit (*Oryctolagus cuniculus*) has been widely used for human disease studies and has served as a suitable experimental model for the treatment of various diseases since the early 1900s, due to its similarity in brain development processes, including neuronal development. In light of the advances and challenges of cell therapy, this manuscript aims to review the literature on the influence and effectiveness of cell therapy in rabbits.

2. METHODOLOGY

This work is a literature review, which used PubMed and Science Direct as databases with the following descriptors: *Cell therapy and experimentation in rabbits; rabbits and cell therapy*. A total of 1,659 articles were found in the PubMed database and 140 articles in the Science Direct database. Inclusion criteria included articles in English and Portuguese published between 2019 and 2024. The selection was based on the reading of titles, abstracts, and introductions; additionally, only articles involving rabbits were included. Figure 1 shows the number of articles included and excluded. The criteria for selecting scientific records were applied by two independent authors, and discrepancies were resolved by consensus between them. The screening of articles was based on reading titles, abstracts, and introductions, and those classified as clinical trials, preclinical trials, and systematic reviews mentioning the use of rabbits in cell therapy were included.

Figure 1 - Flowchart demonstrating the classification of included and excluded articles during the review: Authors, 2024.



3. RESULTS AND DISCUSSION

The advances in cell therapy in this species have sparked growing interest, not only due to its direct applicability in the treatment of debilitating conditions, but also for its contribution to the understanding of the cellular mechanisms underlying tissue repair. With the potential to significantly impact regenerative medicine, this line of research continues to explore new clinical and scientific frontiers, aiming to improve patients' quality of life by replacing dysfunctional cells with healthy ones, thus restoring the function of affected tissues [4].

A study conducted by Zolocinska [17] in 2020 compared animal models for research with adipose tissue-derived stem cells. The investigation indicated that the adipose tissue mass of rabbits was significantly greater than that collected from rats. In addition to this advantage, rabbits are larger, which allows for more complex surgeries. In summary, they present low maintenance costs, ease of handling and housing, allowing for a greater number of surgical procedures, especially in orthopedic, cardiovascular, and neurological experiments. Recent research has shown a promising future in regenerative and replacement therapies involving mesenchymal stem cells (MSCs) and cellular engineering, including tissue-engineered blood vessels (TEBVs). Rodrigues [6] improved TEBV techniques by differentiating adipose tissue-derived mesenchymal stem cells into endothelium and smooth muscle in both an animal model and in vitro, demonstrating good cell colonization and differentiation, as confirmed by morphological characterization. This points to a promising future for the use of TEBV in vivo models. In the continued study of stem cells, Khanmohammadi [18] indicated that the use of menstrual blood-derived stem cells (MenSCs) encapsulated in fibrin glue (FG) in the healing of osteochondral defects shows adequate in vivo regenerative capacity, suggesting they could be used in future clinical trial applications.

Liu [4], in their study, demonstrated that piezoelectric stimulation induced by exercise could promote chondrogenesis and cartilage regeneration in cases of osteoarthritis. The authors suggest that this treatment approach may be potentially applicable to the regeneration of other injured tissues.

De Melo [10] reported preliminary results indicating that the developed orthosis significantly improves patients' mobility and autonomy, facilitating activities such as feeding and personal

hygiene. This advancement is crucial, as the ability to perform such activities independently positively impacts the quality of life and self-esteem of individuals with spinal cord injuries. Despite the promising findings, the study identifies certain limitations, such as the small sample size and the need for an extended evaluation period to validate the long-term efficacy of the orthosis. Furthermore, Coelho [3] conducted a study demonstrating that the administration of mesenchymal stem cells (MSCs) in patients with detrusor underactivity resulted in a significant improvement in the contractile function of the detrusor muscle. However, despite the promising results, the study has some limitations that should be considered. Firstly, the sample size was relatively small, which may limit the generalizability of the findings. Future studies with larger samples are needed to validate these results. Secondly, the follow-up period for the patients was relatively short. Long-term evaluations are essential to determine the durability of the therapeutic effects of MSCs and to monitor for potential late adverse effects. MSCs may reduce neurological sequelae resulting from distemper. Villalba-Briones [7] addressed the host-parasite relationship and how stem cells improved the immune system of rabbits, which responded positively to treatment.

Additionally, Barbosa [11] introduced innovative CAR-T therapy for the treatment of multiple myeloma as a personalized cell therapy for cancer in rabbits. However, challenges such as associated toxicities and the need to improve CAR-T production and administration remain significant. Future research should explore ethical therapeutic combinations and resistance mechanisms to expand the clinical use in humans effectively.

Mesenchymal stromal cells (MSCs) have gained attention due to their immunomodulatory effects, evidenced both *in vitro* and *in vivo*. Nagubothu [5] aimed to determine whether local injection of these cells could modulate the inflammatory response. The study demonstrated that MSCs were able to positively modulate the early wound healing response by resolving the inflammatory phase and promoting tissue repair. However, a low level of injected MSCs persisted in the VF tissue.

Additionally, a study evaluated the effect of hyperbaric oxygen therapy (HBOT) on graft healing after anterior cruciate ligament (ACL) reconstruction, suggesting that the adjuvant use of HBOT may improve graft maturity and integration, along with knee stability, providing better healing outcomes after ACL reconstruction [9]. However, the study presented limitations regarding the ligamentization phase, which is a more prolonged process, making it impossible to assess late events related to the healing process. Furthermore, specific biological mechanisms related to the effects of HBOT on graft healing after ACL reconstruction were not explored. Nevertheless, the study provides relevant insights into the adjuvant use of hyperbaric oxygen in ACL reconstruction, showing therapeutic potential.

In the context of rare genetic diseases, gene editing techniques such as CRISPR-Cas9 represent a significant breakthrough. However, technical challenges persist, such as efficient delivery and safety, as well as ethical and accessibility concerns, which need to be addressed. The implementation of strict regulations and international collaboration are essential to ensure the safety and efficacy of these therapies. As highlighted by Azevedo [8], it is imperative to increase investments in technology to enable the application of these therapies in rabbits.

4. CONCLUSION

The present study offers a promising perspective on animal experimentation in rabbits, particularly highlighting its reduced applicability in terms of invasiveness, simplicity, and low cost, as evidenced by the authors. The reviewed articles and theses provided a comprehensive and diverse analysis of cell therapy and its various applications. Most

studies demonstrated significant efficacy when rabbits were subjected to cell therapies, whether autologous or allogeneic. Therefore, further investigations are needed to elucidate scientific advancements in the treatment and management of diseases made possible through the ethical and safe use of these animal models.

REFERENCES

1. Castro APR, Brito JÁ, de Souza Silva N, de Oliveira ALF, Barreto RGS, Andrade ÍB, et al. Stem cell treatment in amyotrophic lateral sclerosis (ALS): a narrative review of the literature. *Rev EletrônicaAcervoSaúde*. 2022;15(11)
2. Torre BEF, Pinto GHNF, de Oliveira LAC, Moraes MR, Rodrigues MC, Rodrigues VF. Applicability of dental stem cells in tissue regeneration. *Libertas Odontol*. 2023;2(1).
3. Coelho HR. Effect of mesenchymal stem cells in patients with detrusor hypocontractility. 2022.
4. Liu Y, Dzidotor G, Le TT, Vinikoor T, Morgan K, Curry EJ, et al. Exercise-induced piezoelectric stimulation for cartilage regeneration in rabbits. *Sci Transl Med*. 2022;14(627).
5. Nagubothu SR, Sugars RV, Tudzarovski N, Andrén AT, Bottai M, Davies LC, et al. Mesenchymal stromal cells modulate tissue repair responses within the injured vocal fold. *Laryngoscope*. 2020;130(1)
6. Rodrigues LDS. Tissue engineering in blood vessels: an in vitro rabbit model. 2023.
7. Villalba-Briones R, Barros-Diaz C, Gallo-Pérez A, Blasco-Carlos M, Molineros EB. First description of sarcoptic mange in a wild coati (*Nasua narica*) in Ecuador and co-occurrence of canine distemper virus. *Rev Bras Parasitol Veter*. 2022;31.
8. Azevedo KJPLR, de Barros AM, Pinheiro GO, de Brito MBR, de Moura Vardasca GH, Cadête MLX, et al. Advances in gene therapy for rare genetic diseases. *Rev Ibero-Americana HumanidadesCiênc Educ*. 2023;9(8):1484-90.
9. Leite CBG. Effect of hyperbaric oxygen therapy on anterior cruciate ligament reconstruction in an experimental model in rabbits [doctoral thesis]. Universidade de São Paulo; 2023.
10. De Melo AC, Malcher ELS, da Silva PKE, Pereira NCS. Development of a multifunctional orthosis adapted for upper limbs of patients with spinal cord injury. *Rev CPAQV*. 2023;15(3).
11. Barbosa RS, Coelho A, Neto S, Nassar YL. CAR-T cells in the treatment of multiple myeloma: recent advances and prospects. *Rev Estud Multidisc UNDB*. 2023;3(3).
12. Li Y, Shi G, Han Y, Shang H, Li H, Liang W, et al. Therapeutic potential of human umbilical cord mesenchymal stem cells on aortic atherosclerotic plaque in a high-fat diet rabbit model. *Stem Cell Res Ther*. 2021;12:1-14.
13. Park EJ, Kim H, Lim Y, Lee SY, Lee WY. Effect of motivated physicians and elderly patients with hypertension or type 2 diabetes mellitus in prepared communities on health behaviors and outcomes: A population-based PS matched retrospective cohort study during five-year follow-up period. *PLoS One*. 2024;19(2).

14. Kızıldağ A, Tasdemir U, Arabacı T, Kızıldağ CA, Albayrak M, Şahin B. Effects of autogenous tooth bone graft and platelet-rich fibrin in peri-implant defects: an experimental study in an animal model. *J Oral Implantol.* 2020;46(3):221-6.

15. Pepelassi E, Perrea D, Dontas I, Ulm C, Vrotsos I, Tangl S. Porous titanium granules in comparison with autogenous bone graft in femoral osseous defects: a histomorphometric study of bone regeneration and osseointegration in rabbits. *Biomed Res Int.* 2019;2019:8105351.

16. Kumar R, Kolloli A, Singh P, Vinnard C, Kaplan G, Subbian S. Thalidomide and phosphodiesterase 4 inhibitors as host directed therapeutics for tuberculous meningitis: insights from the rabbit model. *Front Cell Infect Microbiol.* 2020;9:450.

17. Zolocinska A, Siennicka K, Debski T, Gut G, Mazur S, Gajewska M, et al. Comparison of mouse, rat and rabbit models for adipose-derived stem cells (ASC) research. *Curr Res Transl Med* 2020;68(4):205-10.

18. Khanmohammadi M, Golshahi H, Saffarian Z, Montazeri S, Khorasani S, Kazemnejad S. Repair of osteochondral defects in rabbit knee using menstrual blood stem cells encapsulated in fibrin glue: a good stem cell candidate for the treatment of osteochondral defects. *Tissue Eng Regen Med* 2019;16:311-24.

UNDER PEER REVIEW