

Popularization—Scalling Up of improved Dual purpose chickens (Koekoek chickens) in T/maichew District Central Zone of Tigray, Ethiopia

ABSTRACT

Formatted: Space After: 10 pt

In spite of central zone of Tigray have has potential and poultry stock, however, the productivity of their in terms of egg and meat is low since the majority of household depends on local breed chickens and rearing of their chickens through a traditional production practices. Participatory research has become increasingly relevant in public agricultural research to improve agricultural productivity. Then, the aim objectives of the popularization scalling up of Koekoek breed was i) To popularize scale up and evaluate the production performance the Koekoek breed chickens in the small holder communities and ii) To analyze farmers feedback towards this improved breed Koekoek chickens breed at smallholder farmers' management. T/maichew Woreda District, My-sie kebelles kebelle was selected based on the project financial support of Irish Aid Operational Research (OR) purposely. A total of 17 household heads willingly participated who were willingly to adopt the Koekoek breeds. Data record sheet and semi-structured questionnaire was were prepared to collect the required data. Then, using simple descriptive and Likert measurement level used to achieve result. The finding of the research shows that an annual average yield was found of to be 204 eggs per year/ chicken. Similarly, the respondents agreed on that the chickens have well adapted, good growing ability, relatively resistance to diseases, have good scavenging ability, and productivity as well have medium eggs size and good egg and meat tasty of Koekoek chickens. The improved chickens' breed (Koekoek) is better to be transferred to other similar mandate areas of Axum Agricultural Research Center. In addition, collaboration of stakeholders needs to be strengthening so that large scale demonstration community production system is going to be functional. Strengthen the human resource and institutional innovations are important point for dissemination of the improved poultry breeds to all beneficiaries of small holder farmers.

Key word: Popularization Scalling up; Improved improved chicken; Koekoek; and Smallholder

1. Introduction

Poultry production is considered as an integral part of livestock production system which plays an important socio-economic role in developing countries [3,4]. Raising income and urbanization in many parts of the developing world caused a growing demand for animal products. The poultry sector has the potential to provide relatively cheap animal protein to the population and improve nutritional status, create both rural and urban employment for both rural and urban and generate income in time of economic difficulty. Village poultry is the first step on the ladder for poor households to climb out of poverty [1].

In Ethiopia, chicken production system is both traditional and modern production system in which most widespread and almost every rural family owns chicken, which provide valuable sources of family protein and income [5]. The total chicken population in the country is estimated to be 59.5 million and the breed composition indicated that 54.06, 2.6 and 2.8 million are indigenous, exotic and hybrid chickens, respectively which are mainly kept by small holder farmers in scavenging environments and most of (88.5%) the poultry products obtained from the conventional family poultry production system [5]. According to [7] revealed that, feed resource for rural poultry is obtained by scavenging in and around the homesteads and consists of household wastes, anything edible found in the immediate environment, together with a small amount of grain supplements provided by the household. Similarly, the potential supplementary feed resources used by small holders' poultry producers are maize grain, household scraps, cereal debris and wheat [8]. However, Majority of local chickens are kept in an by majority extensive production system which also characterized by low productive performance, late age of sexual maturity because of their poor-low genetic potential in Ethiopia [14]. Additionally, due to the varying the production and reproduction system and infrastructure challenges, the poultry sector's economic contribution is not proportional to the huge chicken population in the country [12].

In Ethiopia, Genetic improvement programs of chickens commenced since in Ethiopia introduced in 1952 were used that used to focus on improving temperate breeds to improve productivity and different breeds of improved breeds like egg and dual purpose and crossing the temperate breeds with indigenous breeds chickens [13]. As a result, an evaluation on the reproductive performance of 62.5% blood level crosses (local * white leghorn) at Debrezeit Agricultural Research Center (DZARC) showed that the highest egg production performance cross bred than the pure white leghorn and Local breeds indigenous chickens [9]. In this regard, Axum Agricultural Research Center under the Tigray Agricultural research Institute (TARI) has been done different research agendas on the areas of poultry productivity improvement to improve the poultry production system of smallholder producers in his mandate areas and taking the mission on excellence of poultry research in the region among the research centers. In this regard Thus, this study aimed designed to popularize scale up and evaluate the improved dual purpose Koekoek chickens on their reproductive performance in smallholder farmers and to analyze the perceptions towards the breeds.

2. MATERIALS AND METHODOLOGY

2.1 Study Area

This study was conducted in Tahtay Maychew district rural administration of the Central Zone of Tigray regional state, northern part of Ethiopia. The district is located in a geographic location of 38°32' and 14°07'E, and 13°15' and 14°39'N and altitude of 1500-2260meter above sea level. The district is located 262 kilometer away of the capital city of Region Mekelle to the shire-Endaslassie route. The annual rainfall of the district received 850.5mm that is mono modal type of rainfall concentrated July to August with relative an annual temperature of 20°C. The agro ecology of Tahtay maychew of the district is tepid to cool sub moist mid-highlands SM₂ 5D-2, 70% mid-land 'Weina-dega' and hot to warm sub moist lowlands or SM14D 30% 'Kola' [20]. The study area has a good climatic condition and compatible for growing crop and livestock husbandry. The majority of crops grown in the area are Teff (epidemic to Ethiopian), Sorghum, Maize, Finger Millet and other horticultural crops whereas the dominant livestock species such as cattle, shoats, Chickens, Beehives and equines.

2.2. Sample size and technique

2.2.1. Sample size and technique

The study was done in Tahitay Maychew District namely Mysis Peasant Association (PA), selected purposively in the willingness of IRISH AID Operational Research (OR) project targeted watershed intervention area. Beneficiaries were also selected on their willingness to participate and adopt the dual purpose chicken breeds. The study was conducted on 17 farmers, and they were prepared chicken traditional houses of traditional house and metal cage. Koekoek chickens are improved breeds which are dual purpose of semi scavengers and reared for both egg and meat production. These breeds are semi scavenging and dominantly black and spotted white color feather helps from predators easily fit to rural poultry keepers and easily reared for poor low income generating farmers especially compared to other non-scavenging improved breeds. For each participant distributed unsexed 45 days old chickens. A total of 20-25 Koekoek chickens were given for each participant after taking after taking the training and prepared their house as well as feed.




Fig 1. Map showing study location

2.3 Capacity building

The farmers and DAs had trained and informed about the improved dual purpose [improved](#) chickens as well as recommended [a](#) management wise. Training was delivered [to](#) the [f](#)farmers and DA to building their knowledge and skillson all poultry management [practices](#) (housing, feeding and watering, health [condition-conditionsete.](#)). [The](#) [C](#)chickens breeds management carried out by the farmerswhereastechnical support and close follow [up](#)done by researchers. In recent years, participatory research has become increasingly relevant in public agricultural research. The degree to which a technology dissemination process is participatory and ensures the participation of all stakeholders, especially the poorest members of society, are frequently used to assess its effectiveness [18]

2.4 Data collection

[The necessary d](#)Data were [for thestudy was](#) collected using recording sheet and semi-structured questionnaires. Beginning from the housing [to](#) all technical information (housing, feed type, yield and perception attributes) data were collected by prepared semi-structured questionnaires. Additionally, the eggs[yield was were](#) collected [through using](#) prepared recording sheet which was kept by the participants for recording the daily egg yield. Recording sheet format had chicken age, number of layers, egg laid daily. Totally, the data [was were](#) collected for consecutive 20

~~weeks~~ ~~weeks~~ where the chickens started laying egg to maximum laid period. The farmers had maintained an average of 8 egg layers during data collection period. Rensis Likert was developed a Likert scale measurement and used to collect respondents' attitudes and opinions towards a product or parameters/items in the form of different agreement and disagreement levels of measurements. Accordingly, the data ~~was~~ ~~were~~ collected through prepared five Likert measurement scale method.

2.5 Statistical Analysis

The data ~~was~~ ~~were~~ collected ~~were~~ ~~and~~ analyzed ~~through~~ using methods of descriptive statistical methods (mean, minimum, maximum and standard deviation) and five scales Likert scale analysis using tools of SPSS. The Likert scales assigned from 1-5, namely strongly Disagree up to Strongly Agree used to analyzed the data. Simple descriptive statistics for the egg mean yield and mean score of agreement level and percentage for frequency perception measurement attribute parameters.



Fig. 2: Physical performance of Koekoek chickens under the farmers' management




Fig. 3: Recording sheet (2011 Ethio calendar) for daily egg layers chickens in each participant farmers

3. RESULTS AND DISCUSSION

3.1. Yield Estimation

The descriptive analysis result in (Table 1) indicated that the performance of Koekoek chicken under the farmers' management was found 204 eggs/year/hen. The result agreed with finding and reported by (Elias and Dagnachew-2023) that the performance production of Koekoek chicken breed gave average mean of egg of 145 per year per hen in famers' management. Similarly, [Yizengaw et al. \(2022\)](#) reported that the research results shows the performance of the Koekoek breeds exceed than the local egg production performance ~~which similar results reported by (Yizengaw et al.,~~

Formatted: Font: Italic

~~2022) and which found the~~ egg yield of local chickens 30-60 /year/hen compared the improved /exotic breeds gave an egg of 135 -180/year /hen. The management for their chickens among farmers makes a difference on the performance of these chickens as shown from the result. The maximum and minimum laid egg ranges from 104 up to 204 which shown almost 100% yield difference among the farmers. This could be due to the feed supplement in quantity and quality. The feed might have low quality to provide the required notorious requirements compared to concentrated feed. No matter the system of production and geographical location the availability, quality and feed price of ingredients among the major constraints of poultry production. The main constraints on chicken production are poor nutrition and health problems. Poultry production is similarly inhibited by reduced contact with goods, markets, and services, poor institution commitment, and absence of skills knowledge as reported by [19]. The farmers were supplied a feed of like maize, sorghum and cereal bran without any processing since there is no processing facility in the study area (rural). The farmers were allowed partially but not fully released as the local breeds to scavenge their chickens (see fig 2). A research report revealed by [10,17] maize, sorghum and sunflower the main feed supplements by poultry keepers and composed of a mixture various crops [16] that showed 95% of chicken keepers provided their products without any processing [15].

Table 1: Average distributed, survived and performance of Koekoek chickens (egg yield /hen/year)

Distributed	Survived	Min	Respondents (n=17)	
			Max	Mean \pm SD
25	22	104	292	204 \pm 67

3.2. Perception of farmers

The descriptive analysis of perception parameters ~~are~~ presented in (Table 2). ~~The analyzed result finding. The result analysis~~ showed that the majority (69%) of the sampled respondents agreed and positive perception on the prepared positive attributed sentences. The popularized dual purpose (meat and egg) chickens have accepted on well adapted to the farmers' management, growing ability and diseases resistant. In other hands, ~~these~~ ~~this~~ improved chickens poor in escaping from dangerous or predators, high in taking feed and less attractive in color by consumers especially in the market compared to the local breeds of red color preferred by consumers due to different cultural perceptions with mean score of 3.44 (68.8%). The respondents appreciated and agreed on the breeds have ability in scavenging additional feeds, gives good egg productivity on the farmers' management fitted. This is compatible a report revealed by [2] the perception of farmers showed that farmers response to the breed promised in terms of age to laying egg and egg market price but less preferred to their market price compared to the local breeds.

~~The response of the~~ ~~The~~ participants ~~respondents~~ agreed on the medium eggs size and good egg and meat tasty of Koekoek chickens. Generally, ~~raring~~ ~~rearing~~ and production of improved Koekoek chickens have to enhance the income farmers for food security of smallholder farmers which is ~~poultry is~~ the first step on the ladder for poor households to climb out of poverty.

Table 2: Participant farmers' feedback on the dual purpose Koekoek chicken and their attribute parameters

Positive attribute sentences	Perception Level (n=17)					Mean	SD
	SA	Agree	No - change	Dis-Ag	SD-Ag		
Koekoek breeds well adapted to farmers management		11(64.70)	3 (17.65)	2 (11.76)		3.5	0.71
Koekoek breeds have good growing ability	6 (35.3)	11 (64.70)				4.35	0.49
Koekoek breeds have disease resistant	2(11.76)	9 (52.94)	6 (35.3)			4.06	1.4
Koekoek breeds less exposed to danger		14(82.35)		3(17.65)		3.65	0.78
Koekoek breeds escaped easily from predators		5 (29.41)	2 (11.76)	10 (58.82)		2.71	0.92
Koekoek breeds less consumed feed		2(11.76)		8 (47.06)	7 (41.17)	1.88	0.93
Koekoek breeds good attractive in color		5 (29.41)		11(64.71)	1 (5.88)	2.53	1.01
Koekoek breeds have good scavenging habit		14 (82.35)	2 (11.76)	1 (5.88)		3.71	0.77
Koekoek breeds gives high yield		15 (88.24)	2 (11.76)			3.88	0.33
Koekoek breeds give having good egg size		9(52.94)	8 (47.06)			3.53	0.51
Koekoek breeds' egg are demanded by consumer		11 (100)				4.00	0.00
Koekoek breeds' meat is tasty		11 (64.71)	5 (29.41)	1 (5.88)		3.5	0.93
Mean weight						3.44	0.68

Note: SA= Strongly Agree (5), A= Agree (4), NC= No change (3), DA= Dis agree (2) and SD= strongly disagree (1)

4. Conclusions and Recommendation

The popularization of improved Koekoek in the smallholder farmers' management for dual purpose (egg and meat) has shown a good result. Farmers' provide locally available feed resources but not enough per recommended as observed in the practical during the study. Generally, from the improved chickens with an annual average yield of egg per chicken was found 204 eggs which are promised result. Similarly, the respondents agreed on that the chickens have well adapted, good growing ability, relatively resistance to diseases, have good scavenging ability, and productivity as well have medium eggs size and good egg and meat tasty of Koekoek chickens. Similarly, the respondents said that the chickens have poor escaping from predators, consumed more feed and less attractive in color compared to their local breeds. Generally, rearing and production of improved Koekoek chickens have to enhance the income farmers for food security of smallholder farmers. The improved chickens' breed (*Koekoek*) is better to be scale out to other similar small holder farmers poultry keepers. In addition, collaboration of stakeholders needs to be strengthening so that large scale demonstration community production system is going to be functional. Strengthen the human resource and institutional innovations are important point for dissemination of the improved varieties to all beneficiaries of the rural poultry keepers.

References

1. Akililu H, Almekinders CJM, Udo HMJ, Van der Zijpp AJ. Village Poultry Consumption and Marketing in Relation to Gender, Religious Festivals and Market access. *Tropical Animal Health and Production*. 2007; 39:165-177.
2. Abadi T, Gebretsadik D, Gebremedhin K, Tsigab T, Zenebe M. Evaluation of Insulin Resistance in Overweight and Obese Dogs. *Int J Vet Sci Res*.2020; 6(1): 064-067.
3. Alders R. (2004). Poultry for profit and pleasure. FAO diversification Booklet 3.FAO (Food and Agriculture Organizations of the United Nations), Rome, Italy.
4. Kondombo SR. (2005). Improvement of village chicken production in a mixed (chicken-ram) farming system in Burikina Faso. PhD thesis submitted to Wageningen University, The Netherlands 208.
5. Tadelle D, Million T, Alemu Y, Peters KJ. (2003). Village chicken production system in Ethiopia: use patterns and performance valuation and chicken products and socio-economic functions of chicken. *Livest. Res. Rural. Dev*. 2003; (15)1
6. CSA. Livestock and livestock characteristics (private peasant holdings). Statistical Bulletin 589, Addis Ababa, Ethiopia.2019
7. Mehari K. Poultry production systems and its feed resources in Ethiopia: a research review. *Sci. J. Ani. Sci*. 2016; 5(2): 220-227
8. Tekalegn Y, Etalem T, and Getinet A. (2017). Poultry Feed Resources and Coping Mechanisms of Challenges in Sidama Zone, Southern Ethiopia. *Food Science and Quality Management*, 2017;(60):77-86
9. DZARC (Debre Zeit Agricultural Research Centre), 1991. Annual Research Report. Debre Zeit Ethiopia. 1991.
10. Elias Gonta, Mengistu Urge , Meseret Girma. Assessment of smallholder chicken feeding practices, available feeds, and evaluation of locally used mixtures in two districts of South Omo Zone, Ethiopia. 2022.

11. Elias Gonta, DemerewGetaneh. Adaptation and performance evaluation of koekoek chicken breed under agro-pastoral management condition of South-OmoZone,Ethiopia. 2023.
12. Mogesse H. Phenotypic and genetic characterization of indigenous chickenpopulations in Northwest Ethiopia (Doctoral dissertation, University of the Free State.2007.
13. Nigussie D. Breeding programs for indigenous chicken in Ethiopia: analysis of diversity in production systems and chicken populations. PhD Thesis, Wageningen University, the Netherlands.2011.
14. Yizengaw Mengesha, Ewonetu Kebede & Ashenafi Getachew.Reviewof chicken productive and reproductive performance and its challenges in Ethiopia, All Life.2022.
15. Kidie HA. Characterization of chicken production system and on-farm evaluation of introduced exotic chicken breeds in Gondar Zuria and Kalu districts of Amhara Region, Ethiopia; 2019
16. Moges F, Mellesse A, Dessie T. Assessment of village chicken production system and evaluation of the productive and reproductive performance of local chicken ecotype in Bure district, North West Ethiopia. *Afr J Agric Res.* 2010;5:1739–48
17. Tsadik E, Tamir B, Goraga Z. Characterization of the adoption of village poultry technology package elements, chicken breeds and forms in the Central Oromia Region, Ethiopia. *J Econ Sust Dev.* 2015;6:2222–855
18. Binswanger-Mkhize HP, Bourguignon C, Brink RVD. Agricultural Land Redistribution: Towards Greater Consensus. Agricultural and Rural development: © World bank; 2009.
19. Ahiwe EU, Omede AA, Abdallh MB, Iji PA. 2018. Managing dietary energy intake by broiler chickens to reduce production costs and improve product quality. *AnimusbandryNutr.* 115
20. Axum Agricultural Research Center (AxARC).Agricultural Production and Farming System of Central Zone of Tigray. Unpublished Manual. 2004