

# Original Research Article

## Phytochemical Analysis and Evaluation of the Antifungal Activity of Total Extracts from Five Plants Against Four Dermatophytes Responsible for Superficial Infections

### ABSTRACT

This study is of great interest due to the rise in superficial infections caused by dermatophytes and the need to find natural alternatives to synthetic antifungals. The objective of this research was to determine the phytochemical composition and antifungal activity of total extracts from five plants: *Zanthoxylum gillettii*, *Distemonanthus benthamianus*, *Gmelina arborea*, *Justicia secunda*, and *Anacardium occidentale* against four dermatophytes, namely *Microsporum canis*, *Trichophyton mentagrophytes*, *Trichophyton rubrum*, and *Trichophyton soudanense*.

To achieve this, aqueous and hydroethanolic extracts were prepared and subjected to phytochemical analysis, followed by evaluations of antifungal activity using the slant double-tube method. Antifungal parameters ( $IC_{50}$  and MFC) were determined. The results revealed the presence of various bioactive compounds, such as flavonoids, tannins, and alkaloids, in most of the extracts. *Z. gillettii* and *D. benthamianus* showed strong efficacy, with  $IC_{50}$  (0.004 to 0.00609 mg/mL);  $IC_{50}$  (0.018 to 0.155) and MFC (0.0975 to 0.39 mg/mL); MFC (0.39 to 12.5 mg/mL) respectively, particularly against *M. canis* and *T. soudanense*. In contrast, *A. occidentale* showed no significant antifungal activity. This study thus highlights the potential of plant extracts in the treatment of skin infections and justifies further research to explore their clinical application.

**Keywords:** plant extracts, dermatophytes, Antifungal Activity

### 1. INTRODUCTION

Dermatophytes are filamentous fungi responsible for common dermatoses. They cause conditions such as ringworm, which affects hair, skin, and nails. These conditions, known as dermatophytoses, are particularly concerning as they affect approximately 20 to 25% of the global population, with a rising incidence in developing countries and increasing resistance to antifungal treatments [1]. The global prevalence of dermatophytoses varies by region, with higher rates in tropical and subtropical areas due to favorable climatic conditions for fungal growth [2].

In Africa, dermatophytoses are a major cause of fungal skin infections. Studies show that the prevalence of these infections is particularly high in rural areas, where hygiene conditions are often poor and access to medical care is limited [3]. The species *Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Trichophyton soudanense*, and *Microsporum canis* are among the most frequently isolated in sub-Saharan Africa. In Côte d'Ivoire, the prevalence of dermatophytoses is estimated at around 15%, primarily affecting children and young adults [3], [4], [5]. Traditional plant-based treatments are common, and these remedies are often the first line of defense against dermatoses.

The interest in studying the anti-dermatophytic activity of medicinal plants lies in several important aspects. First, the growing resistance to synthetic antifungals, such as terbinafine and azoles, makes treating dermatophytoses increasingly difficult. Recent studies show that nearly 20% of *Trichophyton rubrum* strains are resistant to terbinafine, which complicates the management of these [6]. Additionally, current antifungals can lead to undesirable side effects, justifying the exploration of natural alternatives [7], [8]. In light of this resistance to synthetic antifungals, it would be wise to explore medicinal plants, many of which have shown strong antimicrobial activity. Among these plants

are *Senna podocarpa*, *Piliostigma thonningii*, *Terminalia avicennoides*, *Terminalia ivorensis*, and *Terminalia catappa*, which are traditionally used to treat skin conditions in Africa [9].

This study aims to determine the phytochemical composition of aqueous and hydroethanolic extracts and evaluate the anti-dermatophytic activity of five medicinal plants against four major dermatophytes (*Trichophyton rubrum*, *Trichophyton mentagrophytes*, *Trichophyton soudanense*, and *Microsporum canis*) to offer a natural and effective alternative to current antifungal treatments.

## 2. MATERIAL AND METHODS

### 2.1 Plant material

The plant material consisted of organs collected from five plants. These different organs were collected in different areas of Côte d'Ivoire (Table 1). After identification of the different species at the national floristic centre of the Université Félix Houphouët-Boigny, the different organs were washed, cut into small pieces and dried in the shade, then reduced to a fine powder.

Table 1 : plants used

Plants	Collected organ	Harvesting area
<i>Anacardium occidentale</i>	Trunk bark	Abengourou
<i>Distemonanthus benthamianus</i>	Trunk bark	Abengourou
<i>Gmelina arborea</i>	Trunk bark	Abengourou
<i>Justicia secunda</i>	Aerial part	Abengourou
<i>Zanthoxylum gillettii</i>	Trunk bark	Abengourou

### 2.2 Fungal material

The fungal species tested were supplied by the Institut Pasteur de Côte d'Ivoire. These species were collected from various patients attending the Centre University Hospital of Cocody, Ivory Coast.

Table 2: Clinical information and profile of fungal species

Clinical information	Nature of the sample	isolated species	ANTIFUNGALPROFILE				
			5-FC	AMB	FCA	VRC	ITR
Moth	Hairline	<i>T. mentagrophytes</i>	S	S	R	S	R
Athlete's foot	Top of the foot	<i>T. rubrum</i>	S	S	S	S	R
Moth	Hairline	<i>M. canis</i>	S	S	S	S	S
Moth	Hairline	<i>T. soudanense</i>	S	S	S	S	R

### 2.3 Extraction method

Aqueous and hydroethanol extracts were prepared according to the method of Zirihi et al (2003), using distilled water and an ethanol-water mixture (70/30, v/v) as the solvent. One hundred grams (100 g) of plant powder was dissolved in one litre of solvent, giving a ratio of 1:10 (w/v). The mixture was then homogenised vigorously using a blender. The homogenate obtained was wrung out in a square of percale cloth and then filtered three times on cotton wool and then on filter paper. The filtrate was evaporated at 45°C using a Venticell® type ventilated oven for 24 h for the 70% ethanolic extract and 48 to 72 h for the aqueous extract.

### 2.4 Phytochemical sorting

Phytochemical sorting is a method used to characterise the main chemical groups such as sterols, polyterpenes, alkaloids, tannins, phenolic compounds, flavonoids, quinones and saponins. These compounds were identified using appropriate reagents that reacted with chemical compounds to give specific colours or precipitates that attested to the presence or absence of the desired molecules in the extracts. This study was carried out with the total extracts (aqueous and hydroethanolic) of each plant according to the identification protocol described by Békro et al. [10].

#### 2.4.1 Determination of sterols and polyterpenes

For each plant extract taken separately, five (5) mL of each of the two extracts (aqueous and hydroalcoholic) were evaporated over a sand bath. The residue was dissolved while hot in 1 mL of acetic anhydride; then 0.5 mL of concentrated sulphuric acid was added to the triturate. Whether or not a purple or violet ring appeared at interphase, turning blue and then green, indicated a positive or negative reaction.

#### **2.4.2 Identification of polyphenols**

A drop of 2% alcoholic ferric chloride solution was added to two (2) mL of each hydroethanolic and aqueous extract. The presence or absence of polyphenols was indicated by the appearance or absence of a more or less dark blue-black or green colour.

#### **2.4.3 Flavonoid analysis**

Two (2) mL of each extract was evaporated and the residue taken up in 5 mL of hydrochloric alcohol diluted 2-fold. By adding 2 to 3 magnesium chips, a release of heat followed by a pink-orange or purplish coloration could be observed. The addition of 3 drops of isoamyl alcohol may or may not intensify this colouration. This confirms the presence or absence of flavonoids.

#### **2.4.4 Identification of tannins**

Five (5) mL of each extract was evaporated to dryness. After adding 15 mL of Stiasny's reagent to the residue, the mixture was kept in a water bath at 80°C for 30 min. The presence or absence of catechin tannins was determined by whether or not a coarse flake precipitate was observed. For gallic tannins, the previous solution was filtered. The filtrate was collected and saturated with sodium acetate. The addition of 3 drops of FeCl<sub>3</sub> caused the appearance or absence of an intense blue-black colour, indicating the presence or absence of gallic tannins.

#### **2.4.5 Analysis of quinone substances**

Two (2) mL of each of the two extracts were evaporated to dryness. The residue was triturated in 5 mL of 1:5 hydrochloric acid. The triturate was transferred to a test tube and heated in a boiling water bath for 30 min. After cooling, it was extracted with 10 mL of chloroform. Twice-diluted ammonia (0.5 mL) was added to the chloroform solution. The presence or absence of quinones was indicated by a red or purple colouration or a lack of colour.

#### **2.4.6 Alkaloid analysis**

Six (6) mL of each solution was evaporated to dryness. The residue was taken up with 6 mL of 60° alcohol. The addition of 2 drops of Dragendorff's reagent to the alcoholic solution caused a precipitate or an orange coloration. The addition of 2 drops of Burchard's reagent to the alcoholic solution produced a precipitate or no reddish-brown coloration and indicated a positive or negative reaction.

#### **2.4.7 Identification of saponosides**

Ten (10) mL of the total extract was prepared in a test tube. The tube was shaken vertically for approximately 15 seconds and left to stand for 10 to 15 minutes. The height of the foam formed was measured after this period.

### **2.5 Antifungal Activity**

The tests were conducted separately on young cultures of dermatophytes grown on slant agar. The various plant extracts were incorporated into the agar before the fungal growth. The incorporation of these extracts into Sabouraud agar was performed using the double dilution method in slant tubes. Each series included 12 test tubes, of which 10 contained the plant extract and 2 were control tubes without the plant extract (one serving as a control for germ growth, the other without germ serving as a control for the sterility of the culture medium). The concentrations in the test tubes ranged from 1000 to 1.52 µg/mL. For the 10 tubes in each series, the concentrations varied according to a geometric progression with a ratio of ½, from tube No. 1 to tube No. 10. After the extract was incorporated, all 12 tubes in each series were autoclaved at 121°C for 15 minutes. The tubes were then tilted with a small base at room temperature to allow for cooling and solidification of the agar [11], [12].

Inoculums were prepared separately from young colonies (7 days old) of *Microsporium canis*, *Trichophyton mentagrophytes*, *Trichophyton rubrum*, and *Trichophyton soudanense*. A colony of each

germ was taken with a loop and homogenized in 10 mL of sterilized distilled water, resulting in the mother suspension ( $10^0$ ). From this suspension, a second suspension ( $10^{-1}$ ) was prepared by a tenfold dilution. Thus, for each of the test tubes (except for the control tube for the sterility of the culture medium), the germ culture was performed on the previously prepared media by inoculating 10  $\mu$ L of the  $10^{-1}$  suspension in cross streaks until exhaustion. The cultures were incubated at 30°C for seven days. After this incubation period, the colonies of each germ were counted. The growth in the ten experimental tubes of each series was evaluated as a percentage of survival, calculated against 100% survival in the growth control tube [13]. The treatment of experimental data allowed for the determination of the following antifungal parameters: the minimum fungicidal concentration (MFC) and the concentration for 50% inhibition (IC<sub>50</sub>), the latter being determined graphically [14].

### 3. RESULTS

#### 3.1 Phytochemical Study

Table 3 summarizes the presence of different chemical groups (sterols, terpenes, phenolic compounds, flavonoids, tannins, quinones, alkaloids, and saponins) in aqueous and hydroalcoholic (70% ethanol) extracts of five plants: *D. benthamianus*, *Z. gillettii*, *J. secunda*, *G. arborea*, and *A. occidentale*.

The results show that sterols, polyterpenes, phenolic compounds, flavonoids, and alkaloids are consistently present in almost all extracts, whether aqueous or ethanol-based. Quinones are absent in the majority of extracts, except in *Z. gillettii* and *G. arborea*, where they are present in both types of extracts. Saponins are predominantly found in the aqueous extracts but absent in the hydroalcoholic extracts, particularly in *D. benthamianus*, *Z. gillettii*, *J. secunda*, and *A. occidentale*.

**Table 3 : Secondary metabolites present in plant extracts**

Plants	Extracts	Chemical groups						
		Sterols, terpenes	Phenolic compounds	Flavonoids	Tannins	Quinones	Saponins	Saponins
<i>D. benthamianus</i>	aqueous	+	+	+	+	-	+	+
	70% ethanol	+	+	+	+	-	+	-
<i>Z. gillettii</i>	aqueous	+	+	+	+	+	+	+
	70% ethanol	+	+	+	+	+	+	-
<i>J. secunda</i>	aqueous	+	+	+	-	-	+	+
	70% ethanol	+	+	+	+	-	+	-
<i>G. arborea</i>	Aqueous	+	+	+	+	+	+	+
	70% ethanol	+	+	+	+	+	+	-
<i>A. occidentale</i>	aqueous	+	+	+	+	-	+	+
	70% ethanol	+	+	+	+	-	+	-

#### 3.2 Antifungal activity

The table 4 shows that ethanolic extracts are generally more effective than aqueous extracts in inhibiting and killing the fungal species tested. *Z. gillettii* and *D. benthamianus* stand out for their strong antifungal activity, with very low IC<sub>50</sub> and MFC, especially against *M. canis* and *T. soudanense*, while *A. occidentale* showed no measurable activity.

**Table 4:** Antifungal parameters of the various extracts

Plants extracts		Parameter values for fungal species tested (mg/mL)							
		<i>M. canis</i>		<i>T. mentagrophytes</i>		<i>T. rubrum</i>		<i>T. soudanense</i>	
		IC <sub>50</sub>	MFC	IC <sub>50</sub>	MFC	IC <sub>50</sub>	MFC	IC <sub>50</sub>	MFC
<b><i>A. occidentale</i></b>	aq	ND	ND	ND	ND	ND	ND	ND	ND
	eth	ND	ND	ND	ND	ND	ND	ND	ND
<b><i>D. benthamianus</i></b>	aq	0,035	1,56	0,198	6,25	0,292	50	0,035	1,56
	eth	0,018	0,39	0,155	3,125	0,13	12,5	0,0198	0,78
<b><i>G. arborea</i></b>	aq	0,046	3,125	0,3	12,5	0,26	25	0,045	3,125
	eth	0,0246	0,78	0,13	3,125	0,14	12,5	0,035	0,78
<b><i>J. secunda</i></b>	aq	0,0487	6,25	0,195	6,25	0,195	ND	0,0487	6,25
	eth	0,032	0,78	0,1	3,125	0,16	25	0,031	1,56
<b><i>Z. gillettii</i></b>	aq	0,021	0,39	0,13	1,56	0,1	1,56	0,026	0,39
	eth	0,00609	0,097 5	0,04	0,39	0,04	0,39	0,004	0,0975

#### 4. DISCUSSION

The chemical composition of the studied plants shows variation depending on the type of solvent used, particularly with respect to saponins, which are often more present in aqueous extracts than in 70% ethanol extracts. This result aligns with several previous studies that have shown saponins are more soluble in water than in organic solvents. A study on medicinal plants conducted by Rai et al. [15], reported a similar trend for species in the Fabaceae family, where aqueous extracts contained more saponins compared to hydroalcoholic extracts.

As for sterols, polyterpenes, flavonoids, and phenolic compounds, their presence in almost all extracts (aqueous and hydroalcoholic) is consistent with the work of Mahamane et al. [16], which showed that these groups of compounds are widely soluble in polar solvents, whether aqueous or hydroalcoholic. Phenolic compounds and flavonoids tend to be well extracted by polar solvents like water and ethanol, and this extraction is more efficient when ethanol is mixed with water, maximizing the solvent's polarity.

Quinones, present only in *Z. gillettii* and *G. arborea*, are relatively rare in other extracts. This could be explained by the variability in quinone concentrations depending on the plant species. Other authors, such as Akpo et al. [17], also reported differences in quinone presence between different plant species, suggesting that these secondary metabolites may be specific to certain plants or growth environments.

In comparison, some studies, including those by Bamba et al. [18], have shown that 70% ethanol effectively extracts tannins from various medicinal plants, which is confirmed by these results, where tannins are present in all extracts except *J. secunda*. However, tannins were absent in the aqueous extract of this plant, which could be related to a lower concentration or a different chemical structure that makes them less soluble in water.

The metabolites identified in these plant extracts exhibit significant pharmacological potential, supported by numerous studies. Sterols and polyterpenes have well-documented antimicrobial, anti-inflammatory, and cholesterol-lowering effects [19]. A recent study by Barkas et al. [20], shows that these compounds help reduce serum cholesterol and improve cardiovascular health by acting on cholesterol absorption in the intestines. Phenolic compounds and flavonoids, present in almost all

extracts, are powerful antioxidants. According to Rudrapal et al. [21], these metabolites reduce oxidative damage associated with various chronic diseases, such as diabetes, neurodegenerative diseases, and certain cancers, by neutralizing free radicals.

Tannins, known for their antimicrobial properties, have been recently studied by Trepa et al. [19], who showed they inhibit the growth of several pathogenic bacterial strains and viruses. This study suggests their potential use in developing treatments for gastrointestinal infections and viral diseases. Quinones, found in *Z. gillettii* and *G. arborea*, continue to be studied for their antitumor and antimicrobial properties. Oyenihni et al. [22], demonstrated that these compounds can induce apoptosis in cancer cells and inhibit the growth of drug-resistant pathogens.

Alkaloids, present in most extracts, are still widely recognized for their multiple pharmacological properties. Heinrich et al. [23], highlighted their anticancer, antimicrobial, and analgesic effects, emphasizing their crucial role in modern medicine, particularly due to the chemical diversity of these metabolites. Finally, saponins, abundant in aqueous extracts, are renowned for their anti-inflammatory and expectorant activities. Stan et al. [24], showed that these compounds are particularly effective in treating respiratory conditions and have antimicrobial properties, making them useful against certain skin infections.

The results obtained show significant variations in the antifungal activity of aqueous and ethanolic extracts of the tested plants, depending on the fungal species. Unlike the other plants, extracts of *Anacardium occidentale* showed no measurable antifungal activity (no data for IC<sub>50</sub> and MFC). This could be explained by the absence of active antifungal compounds in the parts used for extraction or a concentration too low to produce an effect. Other studies, such as that of Quejada et al. [25], reported moderate activity of some *A. occidentale* extracts on *Candida albicans* species, suggesting that antifungal activity might be specific to certain conditions or extraction methods.

Ethanolic extracts, in almost all cases, show superior antifungal activity compared to aqueous extracts. This observation is consistent with numerous studies, such as that of Sepehri et al. [26], which show that ethanol is a better solvent for extracting bioactive compounds, such as polyphenols and flavonoids, responsible for antifungal effects. Indeed, the CI<sub>50</sub> values of ethanolic extracts of *D. benthamianus* and *Z. gillettii* against *M. canis* and *T. soudanense* are significantly lower than those of aqueous extracts, indicating more potent activity.

The results show that *Z. gillettii* and *D. benthamianus* exhibit excellent antifungal activity, particularly with very low CI<sub>50</sub> values (e.g., 0.00609 mg/mL for *Z. gillettii* on *M. canis*) and MFC values that reflect fungicidal action at low concentrations. These results suggest that these plants may contain powerful antifungal compounds, such as quinones and flavonoids, already known for their effects on pathogenic fungi [27].

Antifungal activities vary depending on the fungal species. For instance, *G. arborea* is more effective against *C. albicans* and *T. rubrum*, while its activity on *T. mentagrophytes* and *M. canis* is more moderate. This suggests possible specificity of the bioactive compounds extracted, depending on the membrane or defense mechanisms specific to each fungal species. This phenomenon is often observed in plant pharmacology, as discussed by Zhou et al. [28], who found differences in fungal species' sensitivity to plant extracts.

These results highlight the therapeutic potential of the tested plants, particularly for fungal skin infections, such as those caused by *T. mentagrophytes* and *T. rubrum*, which are responsible for ringworm and other dermatophytoses. The fungicidal activity at low concentrations suggests that these plants could be used to develop natural treatments for these infections, thus reducing reliance

on synthetic drugs, which can lead to resistance. These findings are supported by recent studies, such as that of Ivanovet al.[29], which demonstrate that plant extracts offer a viable alternative in treating resistant fungal infections.

## 5. CONCLUSION

This study combined phytochemical analysis and antifungal activity. The results reveal a strong therapeutic potential for the tested plant extracts. Phytochemical analyses highlighted the presence of bioactive metabolites such as sterols, polyterpenes, phenolic compounds, flavonoids, tannins, alkaloids, and saponins in the majority of the extracts. These compounds are well-known for their antifungal properties, as confirmed by the IC<sub>50</sub> and MFC values obtained from tests on various fungal species.

Ethanollic extracts proved particularly effective, especially those of *Z. gillettii*, *D. benthamianus*, and *G. arborea*, which showed strong inhibition and fungicidal activity at low concentrations, particularly against *M. canis*, *T. soudanense*, and *T. mentagrophytes*. These results suggest that ethanol, as a solvent, allows for better extraction of the bioactive compounds responsible for antifungal activity, such as flavonoids and quinones. In contrast, *A. occidentale* showed no significant activity, which could be due to the absence of certain antifungal metabolites in the tested parts or a low concentration of these compounds.

## REFERENCES

1. Zhan P, Liu W. The changing face of dermatophytic infections worldwide. *Mycopathologia*. 2017; 182(1) : 77-86. PMID: 27783316, DOI: [10.1007/s11046-016-0082-8](https://doi.org/10.1007/s11046-016-0082-8)
2. Havlickova B, Czaika VA, Friedrich M. Epidemiological trends in skin mycoses worldwide. *Mycoses*. 2008, 4:2-15. PMID: 18783559, doi: [10.1111/j.1439-0507.2008.01606.x](https://doi.org/10.1111/j.1439-0507.2008.01606.x).
3. Bongomin F, Gago S, Oladele RO, Denning DW. Global and Multi-National Prevalence of Fungal Diseases-Estimate Precision. *Journal of Fungi (Basel)*. 2017, 18;3(4):57. PMID: 29371573, DOI: [10.3390/jof3040057](https://doi.org/10.3390/jof3040057).
4. Sule W, Okonko I, Omo-Ogun S, Nwanze J, Ojezele M, Ojezele O, Adeolu A, Soyemi E, Olaonipekun T. Phytochemical properties and in-vitro antifungal activity of *Senna alata* Linn. crude stem bark extract. *Journal of Medicinal Plants Research*. 2011, 5(2):176-183.
5. Yotsu RR, Kouadio K, Vagamon B, N'guessan K, Akpa AJ, Yao A, Aké J, Abbet Abbet R, Tchamba Agbor Agbor B, Bedimo R, Ishii N, Fuller LC, Hay R, Mitjà O, Drechsler H, Asiedu K. Skin disease prevalence study in schoolchildren in rural Côte d'Ivoire: Implications for integration of neglected skin diseases (skin NTDs). *PLoS Negl Trop Dis*. 2018; 17;12(5):e0006489. doi: [10.1371/journal.pntd.0006489](https://doi.org/10.1371/journal.pntd.0006489). PMID: 29771976; PMCID: [PMC5976208](https://pubmed.ncbi.nlm.nih.gov/PMC5976208/).
6. Falana M, Nurudeen Q, Salimon S, Abubakar I. Ethnopharmacological Survey of Medicinal Plants Used in the Management of Skin-Related Conditions in Ilorin, North-Central, Nigeria. *Trad Integr Med*. 2022;8(1):56-76. <https://doi.org/10.18502/tim.v8i1.12404>
7. Kisangau, Daniel P, Ken M. Hosea D, Herbert VM, Lyaruu F, Cosam C. Joseph, Zakaria H. et al. "Screening of Traditionally Used Tanzanian Medicinal Plants for Antifungal Activity." *Pharmaceutical Biology*. 2009 ; 47 (8): 708–16. doi:10.1080/13880200902933039.
8. Hui ST, Gifford H, Rhodes J. Emerging Antifungal Resistance in Fungal Pathogens. *Curr Clin Microbiol Rep*. 2024;11(2):43-50. doi: [10.1007/s40588-024-00219-8](https://doi.org/10.1007/s40588-024-00219-8). Epub 2024 Mar 18. PMID: [38725545](https://pubmed.ncbi.nlm.nih.gov/38725545/); PMCID: [PMC11076205](https://pubmed.ncbi.nlm.nih.gov/PMC11076205/).
9. VISHNU YP, Sharma TA, TARUN HF, Kumar K, ANIMA T, Sharma SA. Dermatophytes: Diagnosis of dermatophytosis and its treatment. *African Journal of Microbiology Research*, 2015 ; 9(19) :1286-1293. <https://doi.org/10.5897/AJMR2015.7374>.
10. Békro Y, Békro JAMA, Boua BB, Trabi FH, éhilé EE. Etude ethnobotanique et screening phytochimique de *Caesalpinia benthamiana* (Baill.) Herend. et Zarucchi (Caesalpiniaceae). *The Science of Nature*. 2007; 4, 217-225. <https://doi.org/10.4314/scinat.v4i2.42146>
11. Coulibaly K, Etien DT, Orso BA, Marie B, Kanga Y, Zirihi GN. Anti-hemorrhoidal medicinal plants of the department of issia: Inventory and cytotoxicity on HFF cells of the ethanolic extract 70% of *Landolphia utilis* A" Chev. (Apocynaceae). 2019;7:101 110. DOI: [10.4236/ijbm.2019.711009](https://doi.org/10.4236/ijbm.2019.711009)
12. Zirihi G.N., Datté J.Y., Kra-Adou K.M. & Grellier P., 2003. Phytochemical and pharmacological studies of the alcoholic extract (MFA) of *Fagara macrophylla* Oliv. Engl. (Rutaceae): the chemical

- structure of the active compound inducing antipaludic activity. *Journal of Chinese Clinical Medicine*, 2: 205-210.
13. Bagre I, Bahi C, Ouattara K, Guede NZ, Djaman AJ, Coulibaly A, N'Guessan JD, "Étude botanique et exploration de l'activité antifongique de *Morinda morindoides* (Baker) Milne-Redh sur la croissance *in vitro* de *Cryptococcus neoformans*". *Phytothérapie*. 2011; 136-141. DOI: [10.1007/s10298-011-0612-y](https://doi.org/10.1007/s10298-011-0612-y)
  14. Kra AKM, Ahon GM, Djo-Bi D, Ouattara S, Coulibaly A, Djaman AJ. Antifungal activities of medicinal plants extracts of Ivorian pharmacopoeia. *Journal of Intercult Ethnopharmacology*. 2014; 3: 159-166. doi: [10.5455/jice.20140627125512](https://doi.org/10.5455/jice.20140627125512). Epub 2014 Jul 10. PMID: 26401367; PMCID: [PMC4576806](https://pubmed.ncbi.nlm.nih.gov/PMC4576806/).
  15. Rai S, Kafle A, Devkota HP, Bhattarai A. Characterization of saponins from the leaves and stem bark of *Jatropha curcas* L. for surface-active properties. *Heliyon*. 2023 Apr 28;9(5):e15807. doi: [10.1016/j.heliyon.2023.e15807](https://doi.org/10.1016/j.heliyon.2023.e15807). PMID: 37187903; PMCID: [PMC10176063](https://pubmed.ncbi.nlm.nih.gov/PMC10176063/).
  16. Mahamane IIA, Hama HH, Alio SA, Nodjitoulou M, Moctar C, Hama GR, Bakasso S, Ilagouma AT. Ethno Botanical, Pharmacology and Phytochemistry of widely used medicinal plants in Niger: A Review. *Journal of Medicinal Plants Studies* 2022; 10(4): 46-60 DOI: <https://doi.org/10.22271/plants.2022.v10.i4a.1436>.
  17. Akpo KJM, Sangare OMM, Sacramento IT, Issotina AZ, Guinnin FF, Hounbeme A. Analyse phytochimique des extraits éthanoliques de la variété blanche d'*Hibiscus sabdariffa* Linn (Malvaceae) et évaluation de toxicité aiguë par voie orale chez des rats Wistar. *Int. J. Biol. Chem. Sci.* 2023. 17(7): 2909-2924,
  18. Bamba M, Bordage S, Sahuc ME, Moureu S, Samailie J, Roumy V, Vauchel P, Dimitrov K, Rouillé Y, Dubuisson J, Tra Bi FH, Séron K, Sahpaz S. Anti-HCV Tannins From Plants Traditionally Used in West Africa and Extracted With Green Solvents. *Front Pharmacol.* 2022 Jan 28;12:789688. doi: [10.3389/fphar.2021.789688](https://doi.org/10.3389/fphar.2021.789688). PMID: 35153750; PMCID: [PMC8831738](https://pubmed.ncbi.nlm.nih.gov/PMC8831738/).
  19. Trepa M, Sułkowska-Ziaja K, Kała K, Muszyńska B. Therapeutic Potential of Fungal Terpenes and Terpenoids: Application in Skin Diseases. *Molecules*. 2024; 29(5):1183. <https://doi.org/10.3390/molecules29051183>.
  20. Barkas F, Bathrellou E, Nomikos T, Panagiotakos D, Liberopoulos E, Kontogianni MD. Plant Sterols and Plant Stanols in Cholesterol Management and Cardiovascular Prevention. *Nutrients*. 2023 Jun 22;15(13):2845. doi: [10.3390/nu15132845](https://doi.org/10.3390/nu15132845). PMID: 37447172; PMCID: [PMC10343346](https://pubmed.ncbi.nlm.nih.gov/PMC10343346/).
  21. Rudrapal M, Khairnar SJ, Khan J, Dukhyil AB, Ansari MA, Alomary MN, Alshabrimi FM, Palai S, Deb PK, Devi R. Dietary Polyphenols and Their Role in Oxidative Stress-Induced Human Diseases: Insights Into Protective Effects, Antioxidant Potentials and Mechanism(s) of Action. *Front Pharmacol.* 2022 ; 14;13:806470. doi: [10.3389/fphar.2022.806470](https://doi.org/10.3389/fphar.2022.806470). PMID: 35237163; PMCID: [PMC8882865](https://pubmed.ncbi.nlm.nih.gov/PMC8882865/).
  22. Oyenihni OR, Oyenihni AB, Erhabor JO, Matsabisa MG, Oguntibeju OO. Unravelling the Anticancer Mechanisms of Traditional Herbal Medicines with Metabolomics. *Molecules*. 2021; 26(21):6541. <https://doi.org/10.3390/molecules26216541>.
  23. Heinrich M, Mah J, Amirkia V. Alkaloids Used as Medicines: Structural Phytochemistry Meets Biodiversity-An Update and Forward Look. *Molecules*. 2021 Mar 25;26(7):1836. doi: [10.3390/molecules26071836](https://doi.org/10.3390/molecules26071836). PMID: 33805869; PMCID: [PMC8036335](https://pubmed.ncbi.nlm.nih.gov/PMC8036335/).
  24. Stan D, Enciu AM, Mateescu AL, Ion AC, Brezeanu AC, Stan D, Tanase C. Natural Compounds With Antimicrobial and Antiviral Effect and Nanocarriers Used for Their Transportation. *Front Pharmacol.* 2021 Sep 6;12:723233. doi: [10.3389/fphar.2021.723233](https://doi.org/10.3389/fphar.2021.723233). PMID: 34552489; PMCID: [PMC8450524](https://pubmed.ncbi.nlm.nih.gov/PMC8450524/).
  25. Quejada LF, Hernandez AX, Chitiva LC, Bravo-Chaucanés CP, Vargas-Casanova Y, Faria RX, Costa GM, Parra-Giraldo CM. Unmasking the Antifungal Activity of *Anacardium occidentale* Leaf Extract against *Candida albicans*. *Journal of Fungi*. 2024; 10(7):464. <https://doi.org/10.3390/jof10070464>
  26. Sepehri Z, Javadian F, Khammari D, Hassanshahian M. Antifungal effects of the aqueous and ethanolic leaf extracts of *Echinophora platyloba* and *Rosmarinus officinalis*. *Curr Med Mycol.* 2016; 2(1):30-35. doi: [10.18869/acadpub.cmm.2.1.30](https://doi.org/10.18869/acadpub.cmm.2.1.30). PMID: 28681010; PMCID: [PMC5490295](https://pubmed.ncbi.nlm.nih.gov/PMC5490295/).
  27. Förster C, Handrick V, Ding Y, Nakamura Y, Paetz C, Schneider B, Castro-Falcón G, Hughes CC, Luck K, Pooapati S, Kunert G, Huffaker A, Gershenzon J, Schmelz EA, Köllner TG. Biosynthesis and antifungal activity of fungus-induced O-methylated flavonoids in maize. *Plant Physiol.* 2022 Jan 20;188(1):167-190. doi: [10.1093/plphys/kiab496](https://doi.org/10.1093/plphys/kiab496). PMID: 34718797; PMCID: [PMC8774720](https://pubmed.ncbi.nlm.nih.gov/PMC8774720/).
  28. Zhou X, Zeng M, Huang F, Qin G, Song Z, Liu F. The potential role of plant secondary metabolites on antifungal and immunomodulatory effect. *Appl Microbiol Biotechnol.* 2023; 107(14):4471-4492. doi: [10.1007/s00253-023-12601-5](https://doi.org/10.1007/s00253-023-12601-5). PMID: 37272939; PMCID: [PMC10240486](https://pubmed.ncbi.nlm.nih.gov/PMC10240486/).

29.Ivanov M, Ćirić A, Stojković D. Emerging Antifungal Targets and Strategies. Int J Mol Sci. 2022 ; 23(5):2756.[doi: 10.3390/ijms23052756](https://doi.org/10.3390/ijms23052756). PMID: 35269898; PMCID: [PMC8911111](https://pubmed.ncbi.nlm.nih.gov/PMC8911111/).

UNDER PEER REVIEW