

Original Research Article

Determination of Correlation and Path Coefficients ~~analysis~~ Analysis for Tomato Yield and Quality Attributes (*Solanum lycopersicum* L.)

Abstract

A study was conducted on 14 genotypes (10 Lines and 3 Testers and a check-‘Pant T1’) of tomato (*Solanum lycopersicum* L.) at the Vegetable Research Farm of the Department of Horticulture, Institute of Agricultural Sciences, Banaras Hindu University, Varanasi-221005, during the Rabi season of 2019-20, 2020-21 and 2021-22. The research focused on examining correlations and path coefficients among 15 yield-contributing traits. The experimental design employed was a Randomized Block Design with three replications. The yield per ha was significantly positively correlated with fruit yield per plant (0.943 and 0.876), fruits per plant (0.705 and 710), fruit width (0.266 and 0.296), number of primary branches (0.264 and 0.280), average fruit weight (0.246 and 0.246), pericarp thickness (0.245 and 0.283) and TSS (0.204 and 0.210) at phenotypic and genotypic respectively. Whereas the number of fruits per plant (0.710) showed ~~the~~ a significantly positive indirect effect with fruit yield (q/ha) and other traits as average fruit yield per plant (0.876), fruit width (0.296), pericarp thickness (0.283), primary branches per plant (0.280), average fruit weight (0.248), TSS (0.210) and plant height (0.188) are also showed positive indirect impact on fruit yield (q/ha) at genotypic level which specified that yield can be improved by enhancing these attributes.

Comment [BK1]: Check it

Keywords: tomato, genotypic correlation, phenotypic correlation, genotypic path analysis, yield and attributes.

Introduction

Tomato (*Solanum lycopersicum* L.) is a member of the Solanaceae family and is a significant warm season self-pollinating vegetable ~~which-that~~ is cultivated across both fresh as well as processed industries (Nwosu *et al.* 2014). Increasing popularity with consumers for ~~their~~ its nutritional benefits, farmers for their high demand, and investigators for its genetic as well as genomic traits. India produced 20.33 MT, with 0.84 million ha under cultivation (Anonymous, 2021-22, Choudhury *et al.* 2021). Tomato ~~is consisting~~ consists of phytonutrients such as ~~vit~~ Vit. A, Vit. C, GSH, carotenoids, and polyphenols, as well as minerals such as calcium, phosphorus, and iron. They are eaten fresh and in a variety of recipes. Tomato and ~~their-its~~ byproducts are the

most abundant dietary source of lycopene and antioxidants in the diet of human beings, while lycopene levels ~~increasing-increase~~ 500-times at the time ~~of~~ maturity. (Bai and Lindhout, 2007).

Systematic research and assessment of genotypes ~~is-are~~ critical for agricultural agronomic and genetics development now and in the coming years. Furthermore, ~~in-order~~ to determine the correlation coefficient and path analysis. A complicated attribute, yield consists of a series of both linear and non-line relationships between yield components that have differing degrees of influence which is known as ~~the~~ correlation coefficient. When formulating an effective breeding plan, genetic criteria such as the degree of association between different traits and the direct and indirect impacts of characters that contribute to the overall fruit yield are constantly significant called ~~as~~-path analysis (Sharma *et al.*; 2019, Tasnimet *al.* 2022).

Enhancement of self-pollinated plants such as tomatoes is typically accomplished by hybridization or the selection of cultivars containing desired natural trait variations (Salunke *et al.*; 2012). In light of the foregoing context, the study has been designed accordingly. The present research investigation was designed with all of this information considered ~~in-order~~ to evaluate ~~the~~ correlation coefficient and path analysis through direct and indirect association of characters with yield per ha in tomato.

Materials and Methods

During the 2019–2020 Rabi season, ongoing studies were conducted at the Vegetables Research Field of the Department of Horticulture, Faculty of Agriculture Sciences, B.H.U., Varanasi-221 005, Uttar Pradesh. Here ~~are-is~~ information on the materials and techniques utilized in the experiment. The genotype material was collected from ICAR Indian Institute of Vegetable Research, Jakhani, Varanasi, that ~~were-was~~ used in this investigation.

Three replications, each with a plot size of 3 x3 meters, were used in the Randomized Block Design (RBD) experiment layout. All genotype's plants were spaced 60 x 60 centimeters² apart. ~~In-order-t~~To grow the commercial crop of tomatoes for the trial in the present research, a total of sixteen genotypes were used. The experimental field was repeatedly ~~ploughed-plowed~~ with a disc ~~ploughplow~~, and varieties ~~are-showed~~were shown after followed by planking. The necessary thickness was acquired ~~prior-to~~before transplantation. Plots under designated areas were created. A base fertilizer dosage is administered.

Analytical statistics

Using the formulas of Johanson *et al.* (1955), the correlation coefficients of variation between genotype and phenotype were calculated. The path analysis was computed using the Dewey and Lu (1959) approach.

Result and discussion

Correlation coefficients

Fig. 1 and Table 3 present the correlation coefficients between every possible pair of ten quantitative characters. Correlation analysis can be used to identify the component traits that will serve as the basis for selection ~~in order~~ to increase yield. Both genotypic and phenotypic correlations ~~have been shown to bare~~ ~~significant~~ ~~significance~~; however, the genotypic correlations (rg) ~~have been found to bare~~ greater in magnitude than the phenotypic correlations (rp), suggesting that genes had an impact ~~in on~~ the expression of every characteristic as well as the traits under study were innately connected. According to Panda *et al.* (2016), there is a strong positive association between traits and fruit yield per ha. This correlation may be attributed to increased phenotypic correlation and decreased environmental variance, as well as genetic linkage.

All characters were positively or negatively correlated with fruit yield per ha, which was one of the important characteristics. The yield per ha was observed highly positive correlation with fruit yield per plant (0.943 and 0.876), number of fruits per plant (0.705, 710), fruit width (0.266, 0.296), number of primary branches (0.264 and 0.280) have been significantly positively correlated at phenotypic and genotypic respectively. Similar results were found by Souza *et al.* (2012) and Khan *et al.* (2020), similar findings were found by Behera *et al.* (2020), that ~~is~~ the fruit yield ~~were was~~ positively associated with ~~several number of~~ fruits per plant and Sahoo *et al.* (2022). While average fruit weight (0.246 and 0.246), pericarp thickness (0.245, 0.283), and TSS (0.204, 0.210) were also significantly positively ~~association~~ ~~associated~~ with yield per ha at phenotypic and genotypic levels ~~respectively~~, and height of plant (0.188) showed the positive genotypic correlation with fruit yield per ha and these similar results ~~was~~ ~~were~~ observed by Alamet *et al.* (2019), Basforee *et al.* (2020), Arya *et al.* (2023), whereas fruit yield ha⁻¹ phenotypically and genotypically negative correlated with pH (-0.210, -0.319) respectively and

days to 50 % flowering (-0.233) at genotypic level. Thus, a higher yield of that plant would inevitably come from selecting plants with a greater number of branches per plant and more number of fruits per plant. Meena and Bahadur's (2015) research showed a significant and positive correlation between fruit yield and both fruit weight and the total number of fruits per plant. Meitei *et al.* (2014) and Behera *et al.* (2017) ~~and~~ also reported similar findings in tomatoes.

UNDER PEER REVIEW

Table 1 Phenotypic correlation															
	DTF	PH	NPBP	FL	FW	AFW	FYPP	NFPP	PT	NLPF	TSS	AAC	PH	CC	FYPH
DTF	1.0000	0.0417	0.0494	-0.301**	0.204*	-0.0205	-0.1706	-0.1207	-0.209*	0.311**	-0.0728	0.0111	-0.253*	-	-0.1679
PH		1.0000	-0.0553	0.1552	-0.0427	0.307**	0.1493	-0.1128	0.0195	-0.1349	-0.1015	-0.1090	0.0134	0.0564	0.1512
NPBP			1.0000	0.1004	0.213*	0.286**	0.263*	0.0511	0.315**	0.0951	-0.0155	0.324**	-0.0326	0.1681	0.264*
FL				1.0000	0.222*	0.425**	0.1579	-0.1681	0.353**	-0.1266	-0.1347	-0.296**	-0.0095	0.232*	0.1582
FW					1.0000	0.206*	0.267*	0.0552	0.335**	0.411**	0.1365	-0.1262	-0.1202	0.1018	0.266*
AFW						1.0000	0.246*	-0.468**	0.312**	0.1166	-0.1640	-0.232*	-0.0230	0.0841	0.246*
FYPP							1.0000	0.706**	0.246*	-0.0705	0.204*	0.0804	-0.211*	0.1035	0.943**
NFPP								1.0000	-0.0394	-0.1317	0.341**	0.222*	-0.176*	0.0391	0.705**
PT									1.0000	0.0286	0.0802	0.0425	0.0176	0.205*	0.245*
NLPF										1.0000	0.1571	-0.0506	0.1007	0.0115	-0.0705
TSS											1.0000	-0.0066	0.1509	0.0754	0.204*
AAC												1.0000	-0.0365	0.1019	0.0849
PH													1.0000	-	-0.210*
CC														1.0000	0.1030
FYPH															1.0000

DTF- Daysto50%flowering, PH- Plant height (cm), NPBP- Numberofprimarybranches per plant, FL- Fruitlength(mm), FW- Fruitwidth(mm),AFW- Averagefruitweight(g), FYPP- Fruityieldperplant (kg), NFPP- Numberof fruit per plant, PT- Pericarp thickness(mm), NLPF- Number of locules/fruit, TSS- Total soluble solids (⁰Brix), AAC- Ascorbicacid content(mg/100ml), CC-Chlorophyllcontent(nmol/cm²),FYPH- Fruit yieldperha(Kg)

Table 2 Genotypic correlation

	DTF	PH	NPBP	FL	FW	AFW	FYPP	NFPP	PT	NLPF	TSS	AAC	PH	CC	FYPH
DTF	1.0000	-0.0578	0.1012	-0.412**	0.394**	-0.0430	-0.229*	-0.1474	-0.393**	0.446**	-0.1689	0.0071	-0.439**	0.0230	-0.233*
PH		1.0000	-0.0612	0.201*	-0.0870	0.370**	0.190*	-0.1317	0.0819	-0.1417	-0.1199	-0.1509	0.0840	0.1218	0.188*
NPBP			1.0000	0.1190	0.234*	0.307**	0.281*	0.0528	0.357**	0.1269	-0.0130	0.341**	-0.0046	0.223*	0.280*
FL				1.0000	0.246*	0.428**	0.1586	-0.1704	0.399**	-0.1354	-0.1400	-0.304**	0.0004	0.349**	0.1598
FW					1.0000	0.225*	0.295**	0.0606	0.442**	0.469**	0.175*	-0.1344	-0.187*	0.263*	0.296**
AFW						1.0000	0.248*	-0.466**	0.362**	0.1259	-0.172*	-0.240*	-0.0238	0.1271	0.248*
FYPP							1.0000	0.708**	0.283*	-0.0728	0.209*	0.0846	-0.319**	0.1470	0.876**
NFPP								1.0000	-0.0501	-0.1378	0.356**	0.237*	-0.273*	0.0496	0.710**
PT									1.0000	0.0352	0.0806	0.0517	0.0945	0.350**	0.283*
NLPF										1.0000	0.173*	-0.0581	0.1261	-0.0191	-0.0739
TSS											1.0000	-0.0104	0.269*	0.0814	0.210*
AAC												1.0000	-0.1101	0.1183	0.0819
PH													1.0000	-0.1049	-0.319**
CC														1.0000	0.1478
FYPH															1.0000

DTF- Daysto50%flowering, PH- Plant height (cm), NPBP- Numberofprimarybranches per plant, FL- Fruitlength(mm), FW- Fruitwidth(mm),AFW- Averagefruitweight(g), FYPP- Fruityieldperplant (kg), NFPP- Numberof fruit per plant, PT- Pericarp thickness(mm), NLPF- Number of locules/fruit, TSS- Total soluble solids (⁰Brix), AAC- Ascorbicacid content(mg/100ml), CC-Chlorophyllcontent(nmol/cm²),FYPH- Fruit yieldperha(Kg)

Table 3 Genotypic PATH matrix of Fruit yield per ha (kg)

	DTF	PH	NPBP	FL	FW	AFW	FYPP	NFPP	PT	NLPF	TSS	AAC	PH	CC	FYPH
DTF	0.1077	-0.0062	0.0109	-0.0444	0.0424	-0.0046	-0.0247	-0.0159	-0.0423	0.0480	-0.0182	0.0008	-0.0472	0.0025	-0.233*
PH	0.0017	-0.0297	0.0018	-0.0060	0.0026	-0.0110	-0.0056	0.0039	-0.0024	0.0042	0.0036	0.0045	-0.0025	-0.0036	0.188*
NPBP	-0.0031	0.0019	-0.0311	-0.0037	-0.0073	-0.0095	-0.0087	-0.0016	-0.0111	-0.0039	0.0004	-0.0106	0.0001	-0.0069	0.280*
FL	-0.0210	0.0102	0.0061	0.0510	0.0125	0.0218	0.0081	-0.0087	0.0203	-0.0069	-0.0071	-0.0155	0.0000	0.0178	0.1598
FW	-0.0278	0.0062	-0.0166	-0.0174	-0.0707	-0.0159	-0.0208	-0.0043	-0.0313	-0.0332	-0.0124	0.0095	0.0132	-0.0186	0.296**
AFW	0.0011	-0.0095	-0.0078	-0.0109	-0.0058	-0.0255	-0.0063	0.0119	-0.0093	-0.0032	0.0044	0.0061	0.0006	-0.0032	0.248*
FYPP	-0.2472	0.2049	0.3026	0.1710	0.3176	0.2669	1.0778	0.7631	0.3047	-0.0785	0.2255	0.0912	-0.3437	0.1584	0.876**
NFPP	0.0045	0.0041	-0.0016	0.0053	-0.0019	0.0144	-0.0219	-0.0309	0.0015	0.0043	-0.0110	-0.0073	0.0084	-0.0015	0.710**
PT	-0.0206	0.0043	0.0187	0.0209	0.0232	0.0190	0.0148	-0.0026	0.0524	0.0018	0.0042	0.0027	0.0049	0.0183	0.283*
NLPF	-0.0063	0.0020	-0.0018	0.0019	-0.0066	-0.0018	0.0010	0.0019	-0.0005	-0.0140	-0.0024	0.0008	-0.0018	0.0003	-0.0739
TSS	-0.0019	-0.0014	-0.0001	-0.0016	0.0020	-0.0020	0.0024	0.0041	0.0009	0.0020	0.0115	-0.0001	0.0031	0.0009	0.210*
AAC	0.0000	-0.0009	0.0021	-0.0019	-0.0008	-0.0015	0.0005	0.0015	0.0003	-0.0004	-0.0001	0.0062	-0.0007	0.0007	0.0819
PH	-0.0197	0.0038	-0.0002	0.0000	-0.0084	-0.0011	-0.0143	-0.0123	0.0042	0.0057	0.0121	-0.0050	0.0450	-0.0047	-0.319**
CC	-0.0003	-0.0015	-0.0028	-0.0044	-0.0033	-0.0016	-0.0018	-0.0006	-0.0044	0.0002	-0.0010	-0.0015	0.0013	-0.0125	0.1478
FYPH	-0.233*	0.188*	0.280*	0.1598	0.296**	0.248*	0.876**	0.710**	0.283*	-0.0739	0.210*	0.0819	-0.319**	0.1478	1.0000
Partial R²	-0.0251	-0.0056	-0.0087	0.0081	-0.0209	-0.0063	1.0781	-0.0219	0.0148	0.0010	0.0024	0.0005	-0.0143	-0.0019	

DTF- Daysto50%flowering, PH- Plant height(cm), NPBP- Numberofprimarybranches per plant, FL- Fruitlength(mm), FW- Fruitwidth(mm),AFW- Averagefruitweight(g), FYPP- Fruityieldperplant (kg), NFPP- Numberof fruit per plant, PT- Pericarp thickness(mm), NLPF- Number of locules/fruit, TSS- Total soluble solids (⁰Brix), AAC- Ascorbicacid content(mg/100ml), CC-Chlorophyllcontent(nmol/cm²),FYPH- Fruit yieldperha(Kg)




Fig. 1

Comment [BK2]: Add a suitable name

Path analysis

When dividing the correlation coefficients into the direct and indirect effects of independent variables on a dependent variable, path coefficient analysis is an important tool. The indirect association between more variables in a correlation study becomes more complicated. Two characters might show a connection, since they are corresponded with a normal third one. In such conditions, coefficient examination gives a successful method for a basic assessment of explicit powers activity to create a given relationship and measure the general significance of each component. The fruit yield per ha was used as the dependent variable in this analysis, and the other characters were considered unreliable variables.

The analysis pertaining to about genotypic path matrix for fruit yield (q/ha) revealed that number of fruits per plant (0.710) showed the highest significantly positive indirect effect followed by average fruit yield per plant (0.876), fruit width (0.296), pericarp thickness (0.283), primary branches per plant (0.280), average fruit weight (0.248), TSS (0.210) and plant height (0.188) and however, the highest direct negative impact on fruit yield per ha has been shown by days to 50% flowering (-0.233) and P^H (-0.319). Similar results were found by Meitei *et al.* (2014), Nagariya *et al.* (2015), Sudesh and Anita (2016) and Alam *et al.* (2019).

The fact that the residual factor (0.231) had no effect and did not affect the fruit yield per hectare suggests that there is no other significant yield component left over. The present research findings show that the fruit yield per plant showed a direct and significant positive effect on fruit yield per plant, days to 50% flowering, pericarp thickness, fruit length, pH, TSS and ascorbic acid, so we can conclude that to increase the fruit yield per ha, the traits fruit yield per plant, days to 50% flowering and pericarp thickness play a very important role and similar result was found by Islam *et al.* (2010). Additionally, we saw that Pericarp thickness showed positive direct impact on yield per ha, and similar results also observed by Nagariya *et al.* (2015), Alam *et al.* (2016) and Maurya *et al.* (2022).

Conclusion

In these findings, we can conclude that yield per plant, fruits per plant, fruit width, primary branches, average fruit weight, pericarp thickness, and TSS are contributed to increase

increasing yield and quality in tomatoes simultaneously because these characteristics directly influenced yield and quality. This concentrate likewise uncovered that enormous size tomato natural products are not simply great yielders in addition they are similarly healthfully rich in quality parameters.

References

1. Nwosu, D. J., Onakoya, O. A., Okere, A. U., Babatunde, A. O. and Popoola, A. F. (2014). Genetic variability and correlations in rainfed tomato (*Solanum* spp.) accessions in Ibadan, Nigeria. *Greener Journal of Agricultural Sciences*, 4(5), 211-219.
2. Islam, B. M. R., Ivy, N. A., Rasul, M. G. and Zakaria, M. (2010). Character association and path analysis of exotic tomato (*Solanum lycopersicum* L.) Genotypes. *Bangladesh Journal of Plant Breeding and Genetics*, 23(1), 13–18.
3. Maurya, D., Akhtar, S., Chattopadhyay, T., Kumar, R., Sahay, S., Sangam, S., Kumari, N. and Siddiqui, M.W., 2022. Genetic Variability and Character Association in Tomato (*Solanum lycopersicum* L.). *Bangladesh Journal of Botany*, 51(4), 747-757.
4. Choudhury MAR, Mondal MF, Khan AU, Hossain MS, Azad MOK, Prodhan MDH, Uddain J, Rahman MS, Ahmed N, Choi KY, and Naznin MT. 2021. Evaluation of Biological Approaches for Controlling Shoot and Fruit Borer (*Eariasvitella* F.) of Okra Grown in Peri-Urban Area in Bangladesh. *Horticulturae*. 7(7): 1-8. <https://doi.org/10.3390/horticulturae7010007>.
5. Bai, Y. and Lindhout, P. (2007). Domestication and breeding of tomatoes: what have we gained and what can we gain in the future? *Annals of botany*, 100(5), 1085-1094.
6. Anonymous. Statistical data. Directorate of Agriculture and Corporation, Ministry of Agriculture and Farmer Welfare, New Delhi. 2022, 1-3.
7. Salunke, D. S., Jadhav, A. S., Pawar, B. D., Kale, A. A. and Chimote, V. P. (2012). Diversity analysis of tomato genotypes using RAPD markers and high-performance liquid chromatography in relation to β -carotene content. *Vegetos*, 25(2), 95-101.

8. Tasnim S, Ivy NA, Raihan MS, Akter N, Islam MZ, Jahan N, and Khan AU. 2022. Evaluation of Genetic Variation in Segregating Population of Pea (*Pisum sativum* L.). SAARC Journal of Agriculture. 20(1): 29-39 (2022) Doi: <https://doi.org/10.3329/sja.v20i1.60545>.
9. Sharma, P., Dhillon, N. S., Kumar, V. and Kumar, P. (2019). Correlation and path analysis for yield and its contributing traits in tomato (*Solanum lycopersicum* L.) under the protected environment. *Journal of Pharmacognosy and Phytochemistry*, 8(3), 447-450.
10. de Souza, L. M., Melo, P. C. T., Luders, R. R. and Melo, A. M. (2012). Correlations between yield and fruit quality characteristics of fresh market tomatoes. *Horticultura Brasileira*, 30, 627-631.
11. Panda, A., Paul, A. and Mohapatra, P. (2016). A study on variability, character association and path analysis for pod yield in French bean (*Phaseolus vulgaris* L.). *International Journal of Bio-resource and Stress Management*, 7, 033-039.
12. Arya, D., Pal, A. K., Singh, A. K., Kumar, B. and Rathour, A. (2023). Studies of Correlation and Path Coefficients for Tomato Yield and Quality Attributes (*Solanum lycopersicum* L.). *International Journal of Environment and Climate Change*, 13(10), 381-388.
13. Alam, M. S., Huda, M. N., Rahman, M. S., Azad, A. K. M., Rahman, M. M. and Molla, M. M. (2019). Character association and path analysis of tomato (*Solanum lycopersicum* L.). *Journal of Bioscience and Agriculture Research*, 22(01), 1815-1822.
14. Basfore, S., Sikder, S., Das, B., KV, M. and Chatterjee, R. (2020). Genetic variability, character associations and path coefficient studies in tomato (*Solanum lycopersicum* L.) grown under terai region of West Bengal. *International Journal of Chemical Studies*, 8(2), 569-573.
15. Behera, M., Jagadev, P. N., Das, S., Pradhan, K. and Sahoo, B. B. (2020). Character association and path coefficient studies in tomato. *International Journal of Current Microbiology and Applied Sciences*, 9(9), 2770-2775.
16. Sahoo, B., Panda, A., Mohanty, S. and Senapati, N. (2022). Gene-gene correlation, path coefficients and the genetic variability of tomato (*Solanum lycopersicum* L.) genotypes. *Journal of Crop Weed*, 18(2), 171-184.

17. Meena, O. P. and Bahadur, V. (2015). Genetic associations analysis for fruit yield and its contributing traits of indeterminate tomato (*Solanum lycopersicum* L.) germplasm under open field condition. *Journal of Agricultural Science*, 7(3), 148.
18. Meitei, K. M., Bora, G. C., Singh, S. J. and Sinha, A. K. (2014). Morphology based genetic variability analysis and identification of important characters for tomato (*Solanum lycopersicum* L.) crop improvement. *American-Eurasian Journal of Agricultural and Environmental Sciences*, 14 (3), 1105-1111.
19. Behera, M., Jagadev, P. N., Das, S., Sahoo, B. B., Pradhan, K. and Senapati, N. Character association study on tomato advance lines in Odisha. *International Journal of Chemical Science*, 3 (4), 93-96.
20. Johnson, H. W., Robinson, H. F. and Comstock, R. E. (1955). Estimates of genetic and environmental variability in soybeans. *Agronomy Journal*, 47(7), 314-318.
21. Dewey, D. R. and Lu, K. (1959). A correlation and path coefficient analysis of components of crested wheatgrass seed production. *Agronomy Journal*, 51(9), 515-518.
22. Nagariya, N. K., Bhardwaj, R., Sharma, N., Mukherjee, S. and Umesh (2015). Correlation and path analysis in tomato (*Solanum lycopersicon* L.). *International Journal of Farm Sciences*, 5(4), 111-117.
23. Alam, M. S., Rahman, M. S., Rahman, M. G., Rahman, M. M., Yesmin, S. and Uddin, M. Z. (2016). Correlation and path-coefficient analysis of pummelo. *Journal of Bioscience and Agriculture Research*, 8(1), 718-725.
24. Sudesh, K. and Anita, S. (2016). Assessment of genetic variability, character association and path analysis in tomato (*S. lycopersicum*L.) under tarai condition of Uttarakhand. *International Journal of Agriculture Sciences*, 34(8), 1706-1709.
25. Khan AU, Choudhury MAR, Dash CK, Khan UHS, and Ehsanullah M. 2020. Insect Pests of Country Bean and Their Relationships with Temperature. *Bangladesh Journal of Ecology*. 2 (1): 43-46.