

Biological properties influenced by different tree species in deep Chambal ravine of Madhya Pradesh

Abstract-An experiment site planted with different multipurpose trees and their impact on soil biological properties were measured and analyzed. Based on overall growth on biological properties showed that preferred ranking tree species in the following order viz., maximum soil microbial biomass carbon, dehydrogenase activity and soil respiration were recorded under *Moringa oleifera*, respectively followed by *Albizia lebeck*, *Azadirachta indica*, *Acacia nilotica*, *Dalbergia sissoo*, *Terminalia arjuna*, *Millettia pinnata* and *Gmelina arborea* while *Justicia adhatoda* was recorded minimum, respectively. Tree stand growth significantly improved the soil biological properties. Thus, *A. Moringa oleifera* and *Albizia lebeck* species are most suitable multipurpose hardy tree species for rehabilitation of medium and deep ravines of Chambal ravines in India.

Keywords- Ravine lands, multipurpose trees, biological properties, microbial biomass and soil respiration

Introduction-The land degradation due to water induced erosion is a major global problem and in India it affects more than 28% of total geographical area of the country Dhruvanarayana(1993) and ICAR(2010). Among various forms of land degradation, rain water erosion affects 1.37 million ha area under ravine (GOI, 2003 and ICAR, 2010) and this occurs along several river systems in the alluvial zones of India. In Indian context, the word ravine refers to a network of gullies which are generally spread along any river system. The gully is an erosion channel developed by ephemeral streams with steep banks and a nearly vertical deep gully head enough to create hindrance to the normal tillage operations. The rate of gully erosion depends on the run-off producing characteristics of the watershed which are governed by the size and shape of drainage area, soil characteristics, the alignment, size, shape and gradient of the gully channel. The

gully are widening and ravines are extending at the rate of 8 to 9 m per annum with average soil loss of more than 17 tons ha⁻¹year⁻¹ (Singh *et al.*, 2014). It causes potential threat to nearby cultivated land, but also led to other processes destructive to national economy, viz. floods in river basins, siltation in water reservoirs/tanks and the consequent loss in their storage capacity, choking of estuaries and harbors, damage to railway lines, roads and other public utility properties. In India, Madhya Pradesh state of India alone has 0.68 million ha (Tomaret *et al.*, 2015). These have encroached upon the inhabited villages; many of them had to be shifted to new sites to avoid loss of lives (Verma *et al.*, 2012). The objective of this study was to determine the success of reforestation with multipurpose trees for higher biomass production above & below ground carbon stock and improvement in soil biological properties.

Materials and methods-The present study was conducted at university experimental site at the village Aisah of Morena district of Madhya Pradesh. The geographical location of study area is 26° 40'40.84 N latitude and 78° 06'29.21E longitude with an altitude ranges 150 to 240 m above mean sea level. In both these studies, a heavy emphasis was laid on plantation from the very beginning several species of native forest trees and fruit trees plants were raised during 2012 at 3x3 m spacing evaluated on different sloppy and leveled patches of ravine land. The experiments were started in 2020-21 to 2021-22. Soil sample three hundred twenty four (162 first year and 162 second year) surface and subsurface soil (0-15, 15-30, 30-60, 60-90, 90-120 and 120-150 cm depths) samples collected from different nine trees (6 samples of each tree and one type tree three replications) of site. The experimental design is randomized block design (RBD).

Microbial biomass carbon was determined by the modified fumigation-incubation method Jackinson and Ladd (1981).

$$\text{MBC } (\mu\text{g g}^{-1} \text{ of soil}) = \frac{\text{ECF} - \text{ECNF}}{\text{KEC}}$$

Where, KEC= 0.25 ± 0.05 and it represents the efficiency of extraction of microbial biomass carbon and ECF and ECNF are extractable C in the fumigated and non-fumigated solutions, respectively.

Dehydrogenase activity, as a determinant of total microbial activity in the soil under the tree species, was measured by following the procedure described by Tabatabai (1994). Dehydrogenase activity in soil was calculated as =

$$\frac{\text{Concentration} \times \text{Dilution}}{\text{Incubation time} \times \text{Soil weight}}$$

Where, Concentration = Absorbance value \times Y value of standard curve (78.58).

Soil respiration was estimated by closed jar method Isermeyer (1952).

The rate of soil respiration was calculated by the following formula = $\frac{(V_0 - V) \times 1.1}{Dwt \times T}$

Where, T: Time of incubation in hrs., V_0 : HCl used for blank, V: HCl used for soil sample, Dwt: Dry wt. of 1 g moist soil,

1.1: Conversion factor (1 ml 0.05 M NaOH equals 1.1 mg CO₂).

Results and discussion- Soil microbial biomass carbon ($\mu\text{g/g}$) pooled mean analysis two years experiments positive and significant effect of the treatments on soil organic carbon at different depths 0-15, 5-30, 30-60, 60-90, 90-120 and 120-150 cm. According to the data presented in the table, it is obvious that the maximum soil microbial biomass carbon was recorded under *Moringa oleifera* (65.08, 62.69, 62.24, 56.96, 52.26 and 42.38 $\mu\text{g/g}$), respectively followed by *Albizia lebeck*, *Azadirachta indica*, *Acacia nilotica*, *Dalbergia sissoo*, *Terminalia arjuna*, *Millettia pinnata* and *Gmelina arborea* while *Justicia adhatoda* (53.26, 51.15, 50.99, 42.32, 35.12 and 32.21 $\mu\text{g/g}$) was recorded minimum, respectively. Soil microbial biomass carbon which signifies very labile carbon fraction of active carbon pool which is easily available for plant uptake through mineralization was higher under different tree species as compared to control. The soil microbial biomass carbon was reported to be higher under tree species due to the relatively higher microbial population, the addition of a large quantity of organic matter to the soil. A similar case was with MBP which had direct and significant relation with OC ($r = 0.657^{**}$), available N ($r = 0.455^{**}$), CO₂

($r=0.310^{**}$) and DHA ($r=0.557^{**}$) and significantly negative relationship with pH ($r=-0.134^*$) and APEA ($r=-0.642^{**}$) content in soil of Gird zone. Since, organic matter and microbial activity are typically related to each other, the significant decrease in organic carbon probably intensifies the adverse effects of salinity on soil microbial biomass (Muhammad *et al.*, 2006). As the pH and EC are significantly correlated with biological properties, the special effects of pH and EC were more pronounced. Thus, the study emphasizes that soil EC and pH should be considered as an indicator of changes in biological soil health in soil salinized by saline irrigation water (Singh *et al.*, 2018). The results obtained for soil microbial biomass carbon are verified from the findings of Sharma *et al.*, (2009) and Yadav *et al.*, (2011) who reported higher soil MBC under agro forestry and lower under control. The main reason is the addition of organic inputs in the form of litter fall, recycling of fine root biomass and pruning debris contributed considerably towards organic carbon thus positively affecting microbial population and activity. However, Munoz *et al.*, (2007) reported that microbial biomass carbon was considerably less up to a depth of 10 cm in all agro forestry land use systems compared to treeless.

Dehydrogenase activity (microgram tpf/gm soil/day) the presence of tree species also significantly improves the soil enzymatic activity as represented in table and figure. Dehydrogenase is a good indicator of biological activity taking place in the soil. Dehydrogenase is an important enzyme in the Krebs cycle and electron transport system hence used for studying biological activity. The pooled analysis depicts positive and significant effect of the treatments on dehydrogenase activity at different depths 0-15, 5-30, 30-60, 60-90, 90-120 and 120-150 cm. According to the data presented in the above table, it is obvious that the maximum dehydrogenase activity was recorded under *Moringa oleifera* (90.23, 82.83, 82.47, 79.50, 70.68 and 69.27 microgram tpf/gm soil/day), respectively followed by *Albizia lebeck*, *Azadirachta indica*, *Acacia nilotica*, *Dalbergia sissoo*, *Terminalia arjuna*, *Millettia pinnata* and *Gmelina arborea* while *Justicia adhatoda* (55.34, 53.19,

52.89, 49.50, 47.76 and 45.56 microgram tpf/gm soil/day) was recorded minimum, respectively. Prasad and Mertia (2005) reported considerably high dehydrogenase activity in tree rhizosphere as compared to non-rhizosphere soil owing to a greater supply of carbon and nutrients from dead root cells and rhizodepositions. Yadav *et al.*, (2011) and Lalita (2004) observed that the presence of an agri-silvi-horticultural system exerts a positive impact on the soil by facilitating the build-up of organic carbon thus improving the microbial population and enzymatic activity. Similar results were found by Batra (2004). An Alkaline phosphatase enzyme activity was inversely related and this relationship remains significant with OC ($r=-0.657^{**}$), AN ($r=-0.440^{**}$), MBC ($r=-0.796^{**}$), MBN ($r=-0.772^{**}$), MBP ($r=-0.642^{**}$) and DHA ($r=-0.545^{**}$) and such relations have been also reported by Kumar *et al.* (2013).

Soil respiration the pooled analysis depicts positive and significant effect of the treatments on soil respiration at different depths 0-15, 5-30, 30-60, 60-90, 90-120 and 120-150 cm. According to the data presented in the above table, it is obvious that the maximum soil respiration was recorded under *Moringa oleifera* (31.94, 26.71, 21.74, 12.69, 10.98 and 7.24 CO₂ mg/g of soil/hrs.), respectively. Followed by *Albizia lebeck*, *Azadirachta indica*, *Acacia nilotica*, *Dalbergia sissoo*, *Terminalia arjuna*, *Millettia pinnata* and *Gmelina arborea* while *Justicia adhatoda* (23.19, 14.99, 13.84, 6.44, 6.14 and 5.21 CO₂ mg/g of soil/hrs.) was recorded minimum, respectively. A similar result was found by Greenwood (1961) and Fornelio *et al.*, (2017).

Table 1. Effect of tree species on SMBC, DHA and soil respiration at different soil depths (cm)

S.No.	Treatment	SMBC($\mu\text{g/g}$)						DHA(microgram tpf/gm soil/day)						Soil respiration (CO_2 mg/g of soil/hrs.)					
		0-15	15-30	30-60	60-90	90-120	120-150	0-15	15-30	30-60	60-90	90-120	120-150	0-15	15-30	30-60	60-90	90-120	120-150
T ₁	<i>Moringa oleifera</i>	65.08	62.69	62.24	56.96	52.26	42.38	90.23	82.83	82.47	79.50	70.68	69.27	31.94	26.71	21.74	12.69	10.98	7.24
T ₂	<i>Terminalia arjuna</i>	60.82	54.96	54.52	49.49	41.51	37.73	75.88	74.24	65.68	61.98	57.67	53.54	26.41	20.89	17.94	9.44	6.79	6.52
T ₃	<i>Azadirachta indica</i>	63.30	59.27	59.76	51.78	43.56	40.90	80.65	80.54	80.03	78.43	68.55	66.72	28.25	23.65	19.37	10.53	9.45	6.92
T ₄	<i>Gmelina arborea</i>	56.20	53.81	51.59	45.05	36.32	33.69	67.06	60.94	62.18	52.00	51.82	46.04	24.04	17.44	15.42	8.49	6.41	5.46
T ₅	<i>Millettia pinnata</i>	57.93	54.31	53.03	47.39	39.33	35.39	73.58	74.04	63.53	54.94	55.80	50.41	24.23	20.04	16.83	8.60	6.73	5.87
T ₆	<i>Albizia lebbeck</i>	63.31	60.14	61.66	53.96	43.87	42.01	83.04	80.59	81.80	78.92	70.48	67.97	29.10	25.41	20.80	10.87	10.27	7.06
T ₇	<i>Acacia nilotica</i>	62.47	56.00	56.91	51.29	42.10	40.18	79.67	79.82	77.98	77.83	67.89	65.50	27.62	22.16	19.00	10.36	8.16	6.77
T ₈	<i>Dalbergia sissoo</i>	61.70	53.96	56.49	50.59	41.93	39.36	78.61	77.64	76.21	73.77	67.01	56.38	27.24	21.05	18.78	9.93	7.87	6.65
T ₉	<i>Justicia adhatoda</i>	53.26	51.15	50.99	42.32	35.12	32.21	55.34	53.19	52.89	49.50	47.76	45.56	23.19	14.99	13.84	6.44	6.14	5.21
S.Em. \pm		0.95	1.17	0.97	0.76	0.71	0.56	0.88	1.01	1.00	0.53	0.51	0.94	0.57	0.62	0.27	0.20	0.19	0.19
C.D.		2.73	3.36	2.80	2.18	2.05	1.62	2.55	2.92	2.88	1.52	1.46	2.71	1.64	1.80	0.77	0.57	0.56	0.54




Figure 1. Effect of tree species on SMBC, DHA and soil respiration at different soil depths (cm)

Conclusion-The conclusion of our study results indicated that *Moringa oleifera* (Drumstick) has the most promising tree species followed by *Albizia lebbek*, *Azadirachta indica*, *Acacia nilotica*, *Dalbergia sissoo*, *Terminalia arjuna*, *Millettia pinnata* and *Gmelina arborea* while *Justicia adhatoda* was recorded minimum biological activity. Simultaneously, development of aboveground vegetation canopy due tree suitable planting influence on addition of letter residues, decomposition of leaf litter and fine roots turnover may causes improvement in soil biological properties in long run. Thus, the ravine lands of India have a huge potential for biomass production, livelihood security, C-stocking, improvement in soil environment with adopting tree species to achieve both economic and ecological aim of quick green cover, ecological restoration, carbon balance in sustaining economic and environmental benefits.

References-

- Batra, L. 2004. Dehydrogenase activity of normal, saline and alkali soils under different Agricultural management system. *Journal of the Indian Society of Soil Science*, **52**: 160-163.
- Dhruvnarayna VV and Ram Babu. 1983. Estimation of soil erosion in India. *J. Irrigation Drainage Engg*, ASCE **109**:419-34.
- Fornelio AC, Expósito CV, Pérez TM, and Vilrriales CM, Bermejo H, Copa-Patino JL, Soliveri J, and Serrano JP. 2017. Laboratory analysis of soil respiration using oxygen-sensitive micro plates. *Geoderma***305**: 12–20.
- Greenwood DJ. 1961. The effect of oxygen concentration on the decomposition of organic materials in soil. *Plant and Soil* **14**: 360-376.
- Isermeyer H. 1952. Einzeintache method ezurbestimmung der bodenatmung unter karbonate in Boden Z. Pflanzenernahrung Bodenkd, **56**:26-38.
- Jackinson, D.S. and Ladd, J.N. 1981. Microbial biomass in soil: Measurement and turnover. In: Paul, E.A., Ladd, J.N.eds. *Soil Biochemistry*. Dekker and New York. pp. 415-471.
- Kumar, M.V., Lakshmi, G.V. and Madhuvani, P. 2013. Appraisal of soil fertility status in salt affected soils of ongole division, Prakasam district, Andhra Pradesh. *Journal of the Indian Society of Soil Science*, **61**: 333-340.
- Lalita, B. 2004. Dehydrogenase activity of normal, saline and alkali soils under different agricultural management systems. *J. Indian Soc. Soil Sci.* **52**(2): 160-163.
- Muhammad, S., Müller, T., & Joergensen, R. G. 2006. Decomposition of pea and maize straw in Pakistani soils along a gradient in salinity. *Biology and Fertility of Soils*, **43**, 93-101.

Munoz, C., Zagal, E and Ovalle, C. 2007. Influence of trees on soil organic matter in Mediterranean agroforestry systems: an example from the 'Espinal' of central Chile. *European Journal of Soil Science*. **58**: 728–735.

Prasad, R. and Mertia, R.S. 2005. Dehydrogenase activity and VAM fungi in tree - rhizosphere of a agroforestry systems in Indian arid zone. *Agroforest. Forum*. **63**: 219-22.

Sharma, K.L., Raju, K. R., Das, S. K., Prasad Rao, B. R. C., Kulkarni, B. S., Srinivas, K. Grace, J. K. Madhavi, M. and Gajbhiye, P. N. 2009. Soil Fertility and Quality Assessment under tree, crop, and pasture-based land-use systems in a rainfed Environment. *Communications in Soil Science and Plant Analysis*. **40**: 1436–1461.

Singh A., Verma S.K., Khambalkar P.A., Yadav S.S. and Rajput S. 2018. Biomass Carbon, Carbon Sequestration Potential and Soil Properties as Influence by Different Modules for Management of Chambal Ravines. *Current World Environment*, ISSN: 0973-4929, Vol. **13**, No. (3) 2018, Pg. 464-470.

Singh Y.P., Raghubanshi B.P.S., Tomar R.S., Verma S.K. and Dubey S.K. 2014. Soil Fertility Status and Correlation of Available Macro and Micronutrients in Chambal Region of Madhya Pradesh. *Journal of the Indian Society of Soil Science*, Vol. **62**, No. 4, pp 369-375.

Tabatabai, M.A. 1994. Soil enzymes. In: Methods of Soil analysis. Part 2. Weaver, R.W., Angle, S., Bottomley, P., Bezdicek, D., Smith, S., Tabatabai, A. and Wollum, A. eds. *Soil Sciences Society of America*, Madison, Wisconsin, USA. pp. 775-833.

Tomar, V.S., Verma, G.P. and Singh, Y.P. 2015. Land degradation in India: Strategies to manage Chambal ravines in Madhya Pradesh. *JNKVV Res. J.*, **49**: 366-373.

Verma, G.P., Singh, Y.P. and Dubey, S.K. 2012. *Watershed based reclamation and control of Chambal ravines*. Conservation of Natural Resources for Food and Environmental Security In: Dubey *et al.* (Eds), pp 329–341.

Yadav, R. S., Yadav, B. L., Chhipa, B. R., Dhyani, S. K. and Munna, R. 2011. Soil biological properties under different tree based traditional agroforestry systems in a semi-arid region of Rajasthan, India. *Agroforest Syst.* **81**: 195–202.