

Review Article

Rice Fallow Pulses for Agricultural Potential for South Asian Region

Abstract:

Rice fallows refer to lowland areas where rice is cultivated during the monsoon (*khariif* season) is left unplanted during the winter season (*rabi*) due to various factors. These include the early cessation of monsoon rains which resulted in soil moisture stress during winter season; waterlogging and excessive moisture in November/December, led to unsuitable for winter crops; the absence of suitable varieties for late planting at winter season and socio-economic issues such as damage by stray cattle or blue bulls. India accounts for a significant majority of about 79% of South Asia's total rice fallows which amount to 11.65 million hectares out of the region's total of 15.0 million hectares. The review assess the scenario of rice fallow pulses, production technologies and its constraints in the present trend.

Keywords: Rice fallow, monsoon, pulses, technologies

Introduction:

Rice fallow pulses is a rainfed lowland agro-ecology which is presently seeking attention for a sustainable cropping system in the Southern parts of Asia where rice is grown as a major staple crop. Rice fallows refer to the lowland regions used for rainfed rice cultivation is left unused or uncropped after the rice harvest (Ghosh *et al.*, 2016). There is great potential to increase the cultivation of pulses in rice-fallow conditions, which could enhance food and nutritional security. To achieve this, it is crucial to conserve the remaining soil moisture in these fallow lands. This conservation will allow for the successful introduction of pulses into rice-fallows, leading to increased productivity and profitability. Data estimated that around 11.70 million hectares of land in India remain fallow after the rice harvest. Around 82% of

this fallow land is located in Eastern India, with the remaining area distributed among three southern states: Tamil Nadu, Karnataka and Andhra Pradesh (Singh *et al.*, 2016).

Pulses play an important role in the human diet as they are more rich in protein sources than any other cereals and millets. Thereby, pulses is known as poor man's meat. Pulses are used in day-to-day food in the Asian diet. It has gained an irreplaceable locus in Indian cuisine.

The mineral content consists of calcium (286 mg), iron (15.67 mg), magnesium (553 mg), phosphorus (785 mg), potassium (2035 mg), sodium (79 mg), zinc (6.93 mg), copper (2.03 mg), manganese (3.16 mg) and protein (52.18 g). Additionally, it contains various vitamins such as vitamin A (2.14 mg), C (2.62 mg), B1 (0.27 mg), B2 (0.25 mg), B3 (1.45 mg), B6 (0.91 mg), B9 (0.28 mg), B12 (216 mg) and vitamin B15 (0.91 mg) (Suneja *et al.* 2011).

Pulses, which are part of the Leguminosae family are widely distributed throughout the Indian subcontinent and are commonly referred to as "dal." If pulses are grown as rice fallow crops it supplements soil by implanting nitrogen on the soil and utilizes the unused fertilizer left in the soil at the time of crop cultivation. Pulses enhance the physical, chemical and biological characteristics of the soil, which leads to a decrease in the fertilizer needs of subsequent crops (Garg and Geetanjali, 2009). Recognizing the significance of pulses in human nutrition, the year 2016 was designated as the International Year of Pulses.

With all such significance, the production of pulses is now attaining a stage of decline due to various constraints faced in the present scenario. Due to ever increase in population, rise in income of people and pulses being the major source of dietary protein in India, the demand for pulses continues to grow obviously (Chauhan *et al.*, 2016). About 20% of the total pulse demands are met by imports only. The recommendation for pulses (43 g/day) by the Indian Council of Medical Research but as per the Directorate of Pulses Development in 2016, the recommended daily intake for adult males is 60 g/day and for adult females is 55 g/day. However, the per capita availability in India is only 42 g/day. Kadakia and Jacob, (2009)

estimated that to overcome protein deficiency, pulses alone in India will require 38 million in 2017-18 and to produce this quantity domestically it would be essential to either double its area at current yield levels or double the productivity keeping the acreage constant.

Scope of rice fallow pulses

“There is a significant opportunity to expand total cropland by strategically utilizing rice fallows for cultivation” (Kar and Kumar, 2009; Ghosh *et al.*, 2014). “Pulses stand out due to their high protein content (20 to 25%), capacity to fix atmospheric nitrogen (around 30-150 kg/ha) and their role as a reliable source of income and employment for small-scale and marginal farmers. This places them in a prominent position within global agriculture” (Ali and Gupta 2012). Iriti and Varoni (2017) “stated that throughout human history, pulses have been the cornerstone of staple food preparation, serving as the most vital food grain. They have been extensively relied upon to fulfil basic protein and energy requirements, making them indispensable in sustaining populations across time”. “The accelerated maturation of crops in Direct Seeded Rice presents a chance for the enhancement and broadening of rice fallow lands” (Joshi *et al.*, 2013). “Poor farmers often rely on incorporating perennial grasses or legumes with varying growth durations into crop rotations as a primary method to enhance soil fertility and increase the availability of food and fodder. This approach is crucial for sustaining agricultural productivity while ensuring sufficient nutrition for both humans and livestock” (Sathishkumar *et al.*, 2023).

Rice fallow pulses for various regions

“Pulses are well-suited for the rice-fallow conditions as it demand less water for cultivation and possess deep-rooted systems capable of accessing soil moisture up to 0.4 meters deep. This makes them an excellent choice for maximizing land productivity under rice fallow situations” (Hazra and Bohra, 2021). “Pulse crops characterized by fast growth, early

maturation and the ability to establish extensive ground cover with deep roots are highly sought after rice harvested fields, where soil moisture is typically scarce after harvest of rice” (Hariharan *et al.*, 2022). “Rice fallows in areas of Andhra Pradesh, Tamil Nadu and Karnataka are covered under coastal peninsular zone. The conventional rice–pulse relay systems (lentil, lathyrus, mung bean and urd bean) are being practiced in Odisha, West Bengal, Chattishgarh, Jharkhand, Andhra Pradesh and Tamil Nadu” (Ghosh *et al.*, 2016). “The rice fallows of central region covers Chhattisgarh, Madhya Pradesh and Maharashtra. Soils are generally clay in nature (Vertisols), become hard and develop deep cracks on drying and poor in soil nutrients” (Ali *et al.*, 2014). “Fallow Pulses meets up to 80% of its nitrogen fixation from air and leaves behind substantial amount of residual nitrogen and organic matter for subsequent crops. This makes them valuable not only for their own cultivation but also for enhancing soil fertility and productivity over the long term”. (Singh *et al.*, 1995). The Utera cultivation of pulses can be made more efficient by using short duration and high yielding varieties of rice, as rice will be harvested from the field during September-October (Pande *et al.*, 2012). Maji *et al.*, (2019) suggested Cool-season food legumes such as lentils, chickpeas, field peas and grass peas show great promise in improving productivity and sustainability within the rice-fallow systems of eastern India and the central plateau region.

Suitable pulse crop for rice fallow

“Pulses like lentil and chickpea are the most candidate crops for Temperate regions” (Anon, 2013). “In locations where competitive crops such as linseed and sesame are cultivated in rice fallows, the practice of intercropping these crops with pulses, facilitated by appropriate agricultural machineries, provides an additional avenue to increase crop yield per unit area” (Paharaj and Blaise, 2016). “Pulse crops that show rapid growth, early maturity and form rapid ground cover with deep root systems are much desired for rice fallows as the soil water content is limited after the rice harvest” (Hariharan *et al.*, 2022). “The desirable traits of

pulses, including their deep root system, capacity to access moisture from deeper soil layers, minimal water requirements and ability to establish crops through surface seeding, make them well-suited for cultivation in moisture-limited conditions typically found in rice fallows. Similarly, oilseed crops such as linseed and mustard can also be incorporated into rice-fallow ecosystems to enhance system intensification”. (Hazra *et al.*, 2014; Kumar *et al.*, 2016).

Table1: Pulse Varieties suitable for rice fallows in southern India

S.no.	State	Crop	Rice-fallow crops(varieties)	Reference
1	Tamil Nadu	Black gram	ADT 3, ADT 6	Ramesh and Rathika (2015)
		Green gram	ADT 3	Umamageswari <i>et al.</i> , (2019)
		Horse gram	CRIDA 18R	Vijayakumar and Jalaluddin (2016), Hanif <i>et al.</i> , (2017)
2	Karnataka	Black gram	TAU 1	Shashikumar <i>et al.</i> , (2013)
3	Andhra Pradesh	Green gram	LGG 460’, ‘TM 96-2’, ‘LGG 410’, ‘LGG 407	Rao <i>et al.</i> , (2015)
4	Telangana	Black gram	LBG 17’, ‘LBG 602’, ‘LBG 623’, ‘LBG 402’, ‘LBG 611’, ‘LBG 22’, ‘LBG 648’, ‘LBG 685’, ‘LBG 645’, (2017) ‘LBG 709’ and ‘LBG 752	Reddy and Reddy, (2017)
		Green gram	Pusa 9072’, ‘NARM 1’, ‘NARM 2’, ‘NARM - 18’, ‘LGG460’, ‘LGG 410’, ‘LGG-450’, ‘LGG-	

			407' and 'IM 96-3'	
		Groundnut	Kadiri 4', 'Kadiri 6', 'TAG 24', 'Greshma', 'Rohini', 'Tripura 4' and 'Narayani'	
5	Kerala	Cowpea	PGCP 6	Adarsh and Jacob,2022
		Black Gram	Syama, Sumanjana,	https://pop.kau.in/pulses.htm
		Red Gram	Sa 1	

Plant population and establishment

Amuthaselvi *et al.*, (2019) observed that “in rice fallow black gram cultivation, broadcasting method resulted in a higher plant population per unit area compared to sowing with seed drill. Ensuring an adequate plant population is essential for maximizing returns from rice fallow pulse cultivation” (Subrahmaniyan *et al.*, 2023). “Practicing diversification such as combining rice with pulses in farming systems, enables farmers to mitigate risks and attain a steadier more varied income. This approach enhances their resilience and ability to adapt over time. In the utera production system, the primary difficulty lies in sustaining the ideal plant density” (Ghosh *et al.*, 2016). The effect of organic amendments on yield tends to be either positive or neutral, primarily due to their role in improving soil fertility and nutrient cycling, which are crucial factors for crop productivity (Chen *et al.*, 2018). Iqbal, (2015) observed that 5 per cent solution of moringa leaf extract promoted both the rate and overall percentage of germination in pulse seeds.




Fig:1 Illustration of Rice Fallow Crop Rotation

Nutrient application in rice fallow pulses

“Foliar nutrients are known to influence a wide array of physiological parameters like alteration of plant archetype, assimilate partitioning, promotion of photosynthesis, uptake of mineral ions, enhancing nitrogen metabolism, promotion of flowering, uniform pod formation, increased mobilization of assimilates to defined sinks, improved seed quality, induction of synchrony in flowering and delayed senescence of leaves” (Sharma *et al.*, 2013). In addition to other management practices such as irrigation and plant protection, blackgram responds markedly to précised application of plant nutrients in combination with foliar nutrition especially when applied in a balanced amounts and at the appropriate time (Swaminathan *et al.*, 2021). Gupta *et al.*, (2011) found that 20 kg N ha⁻¹ + rhizobium + PSB + PGPR + 2% urea spray at flowering and 10 days produced a maximum number of nodules and nodule dry weight in chickpeas. Devaraju and Senthivel, (2018) revealed that “application of pulse wonder at 5 kg ha⁻¹ through foliar spray led to increased plant height, increased accumulation of dry matter and enhanced yield characteristics and resulted in a higher yield of 870 kg ha⁻¹ and maximum net returns of Rs. 22149 ha⁻¹, with a benefit-cost ratio of 2.19 in blackgram cultivation”. Vinoth, (2013) concluded that “soil application of recommended NPKS and ZnSO₄ at basal and foliar spray of 1% TNAU pulse wonder at

flowering and 15 days later is the profitable nutrient management package to black gram for getting higher income in terms of higher yield and benefit cost ratio”. Sakthivel *et al.*, (2012) “reported *Pongamia pinnata* as green leaf manure and foliar sprays of *Moringa oleifera* 5% increased number of pods, number of seeds per pod and yield in blackgram”.

Microbial status in rice fallow pulses

Organic matter content of the soil can increase the presence of microbial population in the soil. Residue incorporation results more microbial activity than residue removal or burning (Mandal *et al.*, 2004). Growing of legume crops under rice fallow conditions increase the nitrogen content of soil due to their root nodulations. Maintenance of microbial diversity of soil is necessary for soil health and interactions of rhizosphere with higher plants (Morgan *et al.*, 2005). After harvesting of pulses the beneficial microbial load in the soil may help to maintain better growth of succeeding crop. Pulses grown in dry/post-rainy season after rice harvest may increase the activity of soil enzymes around 10–25 % (Gautam *et al.*, 2021). Intensification of rice fallows with the incorporation of legumes in dry season can be helpful in improving soil health via nitrogen fixation (Dabin *et al.*, 2016). Increase in essential soil nutrients indicated that the combination of rock phosphate and microbial inocula significantly increased the soil fertility among all treatments (Abbasi *et al.*, 2015; Brahim *et al.*, 2017). Venkatesh *et al.*, (2013) reported that 10–15 % increase in soil microbial content and 10–11 % increase in total organic carbon with the inclusion of legumes in rice-based cropping systems.

Challenges for low productivity of rice fallow pulses

Mostly rice fallow pulses are grown under rainfed, low fertile, problematic soils and unpredictable environmental conditions in India with a high degree of risk and often face many constraints. Despite immense scope, the extensive use of rice fallows for pulse

cultivation is mostly restricted because of several biotic, abiotic and socio-economic constraints (Pande *et al.*, 2000).

●Abiotic stress

Insufficient moisture levels during the flowering and harvest stages of winter crops can be detrimental to their growth and yield are common abiotic factors that often results in drought conditions (Pande *et al.*, 2012). Due to the absence of tillage operation weeds pose a significant challenge in relay cropping systems (Ali *et al.*, 2014). Previous season rice crop itself may act as weed and is neglectable. Severe deficiencies in zinc, iron, boron and molybdenum as well as secondary nutrients like sulphur are particularly notable in traditional pulse cultivation (Singh *et al.*, 2013d). More than 90% of farmers suffer lack of suitable rice fallow variety which act as the prime reason for a large area that remains fallow in rabi season (Ghosh *et al.*, 2016). “If ploughing is done after the harvest of rice to remove stubbles, the sowing of rabi pulses are delayed and germination will also be affected due to the formation of large size clods” (Ali *et al.*, 2014). “Rice harvesting through combine harvester in relay cropping resulted in seedling mortality of about 10-15 per cent. Pulses sown after rice under zero tillage conditions is affected due to improper seed dropping in the presence of stubbles and fast drying of surface soil. Regrowing of rice stubbles after harvest is another major problem in the rice fallow relay cropping system in many parts of the country. After harvest of rice soil moisture content gradually declines causing mid and terminal drought at flowering and pod-filling stages which adversely affects the productivity of pulses in rice fallows and sometimes causes 50% reduction in seed yield” (Kumar *et al.*, 2018). “In rice fallows, application of manure or fertilizer is denied due to no-tillage under relay planting and consequently, crops face nutrient stress” (Ali and Masood, 2014). “Unlike manual harvest, machine harvesting of paddy damages the establishing pulse plants due to the trampling effect of wheels on the field traverse. With the wider adoption of mechanised

harvesting, it has emerged as a major abiotic stress in rice fallow cultivation” (Subrahmaniyan *et al.*, 2023).

●Biotic stress

Powdery mildew and yellow mosaic virus are serious diseases of rabi-planted urd bean and mung bean. Legumes are vulnerable to attack by Blue Bulls in the Indo-Gangetic Plains (Singh *et al.*, 2015). Among the biotic constraints, several fungal and viral diseases, insect pests and nematodes constraint both cool season and warm season pulse production in Rainfed rice fallow lands (Pande *et al.*, 2012).

Table 2: Constraints and technologies to Boost Rice Fallow pulses productivity

S.no	Properties	Constraints	Technologies to overcome constraints
1.	Population management	Poor crop stand and establishment	Tillage machines, sowing methods, seed priming, higher seed rate, timely planting, seed treatment with fungicides
2.	Nutrient management	Poor crop growth	Foliar spray of urea/DAP to supplement N and P
3.	Dry spell	Terminal drought during critical stages of crop growth results in reduction in crop growth and yield	Residue mulching and spraying of TNAU crop boosters may help to mitigate stress conditions
4.	Weed menace	Early season rice crop and other weed flora interventions	Post-emergence herbicides like Quizalofop ethyl and Imazethapyr
5.	Pest and disease management	Pest and disease incidence	Development of IPM modules
6.	Lack of	Lack of labours and	Tillage machines, zero-till planter and

	mechanization	mechanization in rice fallows	harvester
--	----------------------	-------------------------------	-----------

Advantages of rice fallows

Under organic farming practices, the soil organic carbon content significantly increases from 0.54% to 1.17% due to the cultivation of a legume crop (Nivethadevi *et al.*, 2022). “Legumes provide nitrogen (N) to agroecosystem through their exclusive capability to fix atmospheric N in symbiotic relationship with soil rhizobia” (Raza *et al.*, 2020). Due to its ability to establish with surface seeding, appropriateness for relay/para cropping and resilience against soil moisture and temperature stress, pulses are the best crops to plant in the spaces left empty after rice (Kumar *et al.*, 2018).

SWOT analysis

•Strengths

The major strength of the rice fallow pulse crops is the existence of large rabi fallow lands that virtually have zero cost. Besides, pulses have better adaptability to marginal lands. Yields of pulses(chickpea) in Chattisgarh and Madhya Pradesh are also higher than the national average. The pulse cultivation on the residual moisture after kharif rice helps in better moisture conservation.

Unavailability of quality seed, poor accessibility to markets, practice of cultivation of long duration rice varieties, lack of short-duration chickpea varieties, weak extension system, drought-like situations at the time of crop maturity and uncertain rainfall are some of the obstacles to production in Rainfed rice fallow lands. Besides water-saving/harvesting technologies and improved tools and implements required for crop establishment.

● Opportunities

The Government of India is committed to introduce the 'Food Security Act (FSA)' in the near future and the success of FSA will depend on augmentation of agricultural production by raising agricultural productivity or cropping intensities of mono-cropped, rainfed and marginal lands apart from other measures. Pulses complement cereals in both production and consumption. These improve soil fertility, requires less water in comparison to cereals and control diseases and pests in rotation with cereals. This indicates scope for legume production in which chickpea plays a significant role.

● Threats

Imbedded production and price risk is the most serious threat to chickpea production. Year to year variation in pulse production is a regular feature. Prices also show erratic behaviour. When farmer cultivates rice fallow pulses in isolated areas, menace of stray animals such as neelgai (blue bull) and monkeys are acute but substantial. Problems of *Helicoverpa* and wilt are the other persistent threats that need immediate attention.

● Conclusion

From this review, it can be concluded that cultivation of rice fallow pulses offer a significant advantage for farmers in terms of incorporating pulses into rice fallows is a sustainable and profitable practice. It also enhances the agricultural productivity, soil health and per capita availability of pulses. Policy and technology driven approach is essential to promote rice fallow pulses area and productivity.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

References

1. Ghosh, P. K., Hazra, K. K., Nath, C. P., Das, A., & Acharya, C. L. (2016). Scope, constraints and challenges of intensifying rice (*Oryza sativa*) fallows through pulses. *Indian Journal of Agronomy*, 61(4), 122-128.
2. Singh, N. P., Praharaaj, C. S., & Sandhu, J. S. (2016). Utilizing untapped potential of rice fallow of East and North-east India through pulse production. *Indian Journal of Genetics and Plant Breeding*, 76(04), 388-398.
3. Suneja, Y., Kaur, S., Gupta, A. K., & Kaur, N. (2011). Levels of nutritional constituents and antinutritional factors in black gram (*Vigna mungo* L. Hepper). *Food Research International*, 44(2), 621-628.
4. Garg, N., & Geetanjali. (2009). Symbiotic nitrogen fixation in legume nodules: process and signaling: a review. *Sustainable Agriculture*, 519-531.

5. Chauhan, J. S., Singh, B. B., & Gupta, S. (2016). Enhancing pulses production in India through improving seed and variety replacement rates. *Indian Journal of Genetics and Plant Breeding*, 76(04), 410-419.
6. Pande, S., Sharma, M., Ghosh, R., Rao, S. K., Sharma, R. N., & Jha, A. K. (2012). Opportunities for Chickpea Production in Rainfed Rice Fallows of India. Baseline Survey Report.
7. Kar, G., & Kumar, A. (2009). Evaluation of post-rainy season crops with residual soil moisture and different tillage methods in rice fallow of eastern India. *Agricultural Water Management*, 96(6), 931-938.
8. Ghosh, P. K., & Hazra, K. K. (2014). Constraints, issues and opportunities of RCT in pulse based cropping systems. *Resour Conserv Technol Pulses*, 18-30.
9. Ali, M., & Gupta, S. (2012). Carrying capacity of Indian agriculture: pulse crops. *Current Science*, 874-881.
10. Iriti, M., & Varoni, E. M. (2017). Pulses, healthy, and sustainable food sources for feeding the planet. *International Journal of Molecular Sciences*, 18(2), 255.
11. Joshi, E., Kumar, D., Lal, B., Nepalia, V., Gautam, P., & Vyas, A. K. (2013). Management of direct seeded rice for enhanced resource-use efficiency. *Plant Knowledge Journal*, 2(3), 119-134.
12. Sathishkumar, A., Subramanian, E., & Selvarani, G. (2023). Ley Farming: A Way to Restore Soil Fertility and Green Fodder Production in Dry Land Areas. *AgroScience Today*, 4(10), 0694-0696.
13. Hariharan, G., Abhayapala, R., Karunarathna, B., & Zakeel, M. C. M. (2022). Efficient utilization of rice fallow through pulse cultivation. In *Advances in Legumes for Sustainable Intensification* (pp. 71-92). Academic Press.

14. Ghosh, P. K., Hazra, K. K., Nath, C. P., Das, A., & Acharya, C. L. (2016). Scope, constraints and challenges of intensifying rice (*Oryza sativa*) fallows through pulses. *Indian Journal of Agronomy*, 61(4), 122-128.
15. Ali, M., Ghosh, P. K., & Hazra, K. K. (2014). Resource conservation technologies in rice fallow. *Resource conservation technology in pulses*, 83-88.
16. Praharaj, C. S., & Blaise, D. (2016). Intercropping: An approach for area expansion of pulses. *Indian J. Agron*, 61(4th IAC Special issue), S113-S121.
17. Hazra, K. K., Venkatesh, M. S., Ghosh, P. K., Ganeshamurthy, A. N., Kumar, N., Nadarajan, N., & Singh, A. B. (2014). Long-term effect of pulse crops inclusion on soil-plant nutrient dynamics in puddled rice (*Oryza sativa* L.)-wheat (*Triticum aestivum* L.) cropping system on an Inceptisol of Indo-Gangetic plain zone of India. *Nutrient cycling in agroecosystems*, 100, 95-110.
18. Kumar, N., Hazra, K. K., Singh, S., & Nadarajan, N. (2016). Constraints and Prospects of. *Indian Farming*, 66(6), 13-16.
19. Pande, S., Sharma, S. B., & Ramakrishna, A. (2000). Biotic stresses affecting legumes production in the Indo-Gangetic Plain.
20. Pande, S., Sharma, M., Ghosh, R., Rao, S. K., Sharma, R. N., & Jha, A. K. (2012). Opportunities for Chickpea Production in Rainfed Rice Fallows of India. Baseline Survey Report.
21. Ali, M., Ghosh, P. K., & Hazra, K. K. (2014). Resource conservation technologies in rice fallow. *Resource conservation technology in pulses*, 83-88.
22. Singh, A. K., Meena, M. K., Bharati, R. C., & Gade, R. M. (2013). Effect of sulphur and zinc management on yield, nutrient uptake, changes in soil fertility and economics in rice (*Oryza sativa*)-lentil (*Lens culinaris*) cropping system. *Indian J. Agril. Sci*, 83(3), 344-348.

23. Singh, D. P. (1995). Pulses in India. *Production of pulse crops in Asia and the Pacific region (Sinha SK and Paroda RS, eds). Bangkok, Thailand: Regional Office for Asia and the Pacific (RAP A), Food and Agriculture Organization of the United Nations*, 101-112.
24. Pande, S., Sharma, M., & Ghosh, R. (2012). Role of pulses in sustaining agricultural productivity in the rainfed rice-fallow lands of India in changing climatic scenario.
25. Kumar, N., Yadav, A., Singh, S., & Yadav, S. L. (2018). Growing pulses in rice fallow: Ensuring nutritional security in India. *Conservation Agriculture for Advancing Food Security in Changing Climate*, 1, 107-122.
26. Ali, M. (2014). Improving productivity of pulses in rice fallows. *Indian Farming*, 63(12).
27. Maji, S., Das, A., Nath, R., Bandopadhyay, P., Das, R., & Gupta, S. (2019). Cool season food legumes in rice fallows: an Indian perspective. *Agronomic Crops: Volume 1: Production Technologies*, 561-605.
28. Ramesh, T., & Rathika, S. (2015). Weed management in rice fallow black gram through post-emergence herbicides. *Madras Agricultural Journal*, 102(oct-dec), 1.
29. Umamageswari, C., Manimaran, R., & Iyanar, K. (2019). Impact of improved production technologies on yield of rice fallow pulses in Cauvery delta zone. *Journal of Pharmacognosy and Phytochemistry*, 8(2S), 963-967.
30. Vijayakumar, M., & Jalaluddin, S. M. (2016). Physiological approaches for maximization of yield in horsegram (*Macrotyloma uniflorum L.*). *Journal of Chemical, Biological and Physical Sciences*, 6(3), 997-1.
31. Hanif, N. A., Ramasamy, R., Davidson, S. J., & Pandiyan, M. (2017). Popularization of Horse Gram (*Macrotyloma uniflorum*) in Vellore District of Tamil Nadu. *Journal of Krishi Vigyan*, 6(1), 148-150.

32. Basavarajappa, R., Salakinkop, S. R., Hebbar, M., Basavarajappa, M. P., & Patil, H. Y. (2013). Influence of foliar nutrition on performance of blackgram (*Vigna mungo* L.), nutrient uptake and economics under dry land ecosystems. *Legume Research-An International Journal*, 36(5), 422-428.
33. Rao, D., Naidu, T. C. M., & Rani, Y. A. (2015). Effect of Foliar Nutrition on Water Potential, Photosynthetic Rate, Dry Matter Production and Yield of Mung Bean Under Receding Soil Moisture Condition.
34. Adarsh, S., & John, J. (2022). Performance of different pulse crops in summer rice fallow of northern Kerala. *Crop Research*, 57(1and2), 8-14.
35. Kerala agricultural university, crop production guide, <https://pop.kau.in/pulses.htm>
36. Anonymous. 2013. Improving Productivity of rice fallows. Policy Paper No. 64, National Academy of Agricultural Sciences, New Delhi. 16 p.
37. Sharma, P. U. S. H. P., Sardana, V. I. R. E. N. D. E. R., & Singh, S. K. (2013). Dry matter partitioning and source-sink relationship as influenced by foliar sprays in groundnut. *The Bioscan*, 8(4), 1171-1176.
38. Swaminathan, C., Surya, R., Subramanian, E., & Arunachalam, P. (2023). Challenges in pulses productivity and agronomic opportunities for enhancing growth and yield in blackgram [*Vigna mungo* (L.) Hepper]: a review. *Legume Research-An International Journal*, 46(1), 1-9.
39. Gupta, S. C., Kumar, S., & Khandwe, R. (2011). Effect of bio-fertilizers and foliar spray of urea on symbiotic traits, nitrogen uptake and productivity of chickpea. *Journal of food legumes*, 24(2), 155-157.
40. Devaraju, B., & Senthivel, T. (2018). Effect of foliar application of different sources of nutrients on growth and yield of blackgram under irrigated

- conditions. *International Journal of Current Microbiology and Applied Sciences*, 7(1), 3105-3109.
41. Vinoth V. Effect of foliar nutrition on the growth and yield of blackgram (*Vigna mungo L.*). (M.Sc). (Ag.) Thesis), Tamil Nadu Agricultural University, 2013.
42. Sakthivel, A., Swaminathan, C., & Rajalakshmi, B. (2012). Influence of *Pongamia pinnata* green leaf manure on growth and yield of blackgram (*Vigna mungo*). *Madras Agric. J*, 99, 83-84.
43. Mandal, K. G., Misra, A. K., Hati, K. M., Bandyopadhyay, K. K., Ghosh, P. K., & Mohanty, M. (2004). Rice residue-management options and effects on soil properties and crop productivity. *Journal of Food Agriculture and Environment*, 2, 224-231.
44. Morgan, J. A. W., Bending, G. D., & White, P. J. (2005). Biological costs and benefits to plant–microbe interactions in the rhizosphere. *Journal of experimental botany*, 56(417), 1729-1739.
45. Gautam, P., Lal, B., Panda, B. B., Bihari, P., Chatterjee, D., Singh, T., ... & Nayak, A. K. (2021). Alteration in agronomic practices to utilize rice fallows for higher system productivity and sustainability. *Field Crops Research*, 260, 108005.
46. Dabin, Z., Pengwei, Y., Na, Z., Changwei, Y., Weidong, C., & Yajun, G. (2016). Contribution of green manure legumes to nitrogen dynamics in traditional winter wheat cropping system in the Loess Plateau of China. *European Journal of Agronomy*, 72, 47-55.
47. Abbasi, M. K., Musa, N., & Manzoor, M. (2015). Mineralization of soluble P fertilizers and insoluble rock phosphate in response to phosphate-solubilizing bacteria and poultry manure and their effect on the growth and P utilization efficiency of chilli (*Capsicum annum L.*). *Biogeosciences*, 12(15), 4607-4619.

48. Brahim, S., Niess, A., Pflipsen, M., Neuhoff, D., & Scherer, H. (2017). Effect of combined fertilization with rock phosphate and elemental sulphur on yield and nutrient uptake of soybean.
49. Venkatesh, M. S., Hazra, K. K., Ghosh, P. K., Praharaj, C. S., & Kumar, N. (2013). Long-term effect of pulses and nutrient management on soil carbon sequestration in Indo-Gangetic plains of India. *Canadian Journal of Soil Science*, 93(1), 127-136.
50. Amuthaselvi, G., Dhanushkodi, V., & Eswaran, S. (2019). Performance of zero till seed drill in raising blackgram under rice fallow.
51. Hazra, K. K., & Bohra, A. (2021). Increasing relevance of pulse crops to sustainable intensification of Indian agriculture.
52. Ghosh, P. K., Hazra, K. K., Nath, C. P., Das, A., & Acharya, C. L. (2016). Scope, constraints and challenges of intensifying rice (*Oryza sativa*) fallows through pulses. *Indian Journal of Agronomy*, 61(4), 122-128.
53. Chen, Y., Camps-Arbestain, M., Shen, Q., Singh, B., & Cayuela, M. L. (2018). The long-term role of organic amendments in building soil nutrient fertility: a meta-analysis and review. *Nutrient Cycling in Agroecosystems*, 111, 103-125.
54. Iqbal, M. A. (2015). Improving germination and seedling vigour of cowpea (*Vigna unguiculata* L.) with different priming techniques. *Am-Eur J Agric Environ Sci*, 15, 265-270.
55. Hariharan, G., Abhayapala, R., Karunarathna, B., & Zakeel, M. C. M. (2022). Efficient utilization of rice fallow through pulse cultivation. In *Advances in Legumes for Sustainable Intensification* (pp. 71-92). Academic Press.

56. Subrahmaniyan, K., Kumar, G. S., Subramanian, E., Raju, M., Veeramani, P., & Ravi, V. (2023). Crop establishment methods and moisture mitigation practices in rice fallow blackgram for productivity enhancement in Cauvery delta zone of Tamil Nadu. *Legume Research-An International Journal*, 46(4), 502-505.
57. Singh, A. K., Singh, S. S., Prakash, V. E. D., Kumar, S., & Dwivedi, S. K. (2015). Pulses production in india: Present status, sent status, bottleneck and way forward. *Journal of AgriSearch*, 2(2), 75-83.
58. Nivethadevi, P., Swaminathan, C., Kannan, P., & Tamilselvi, E. (2022). Seed fortification and foliar spraying with *Moringa oleifera* leaf extract enhances yield and yield attributes in blackgram [*Vigna mungo* (L.) hepper]. *Legume Research-An International Journal*, 45(3), 352-356.
59. Raza, A., Zahra, N., Hafeez, M. B., Ahmad, M., Iqbal, S., Shaukat, K., & Ahmad, G. (2020). Nitrogen fixation of legumes: Biology and Physiology. The plant family Fabaceae: biology and physiological responses to environmental stresses, 43-74.
60. Kumar, N., Yadav, A., Singh, S., & Yadav, S. L. (2018). Growing pulses in rice fallow: Ensuring nutritional security in India. *Conservation Agriculture for Advancing Food Security in Changing Climate*, 1, 107-122.