

Yield and quality of brinjal (*Solanum melongena* L. var. Pusa uttam) as influenced by organic manures and micronutrients

Comment [H1]: I suggest rewrite the title to be: Influence of organic manure and micronutrients on yield and quality of brinjal, *Solanum melongena* L. var. Pusa uttam

ABSTRACT

A field study was conducted at farm of S.K.N., Agriculture University, Jobner, (Jaipur) during kharif season 2018 to investigate effect of organic manures and micronutrients on yield and quality of brinjal (*Solanum melongena* L. var. Pusa Uttam) at farm of S.K.N., Agriculture University, Jobner (Jaipur) during kharif season 2018. The treatments included organic manures (control, FYM @ 40 t/ha), vermicompost @ 6.6 t/ha and poultry manure @ 6.6 t/ha,) and micronutrients (control, Zn @ 0.5%, Fe @ 0.5% and B @ 0.3%). The application of vermicompost @ 6.6 t/ha non significantly affected TSS content and significantly increased the ascorbic acid content, protein content, fruit yield per plot and fruit yield per hectare which was statistically at par with poultry manure @ 6.6 t/ha and micronutrients also non significantly affected TSS content and significantly increased the ascorbic acid content, protein content, fruit yield per plot and fruit yield per hectare as compared to control in brinjal.

Comment [H2]: Please add the country name.

Comment [H3]: ?

Comment [H4]: Is this commercial or common name?

Comment [H5]: ?

Comment [H6]: ?

Key Words – micronutrients, organic manures, TSS, protein content, yield, brinjal.

Comment [H7]: Short abstract, please support the abstract with more results.

INTRODUCTION

Brinjal (*Solanum melongena* L.) is cultivated as one of the leading vegetable crops grown in India. It is popularly known as eggplant belongs to family solanaceae and India is it's center of origin and diversity (Saravaiya *et al.*, 2010). It is highly productive and usually finds a place as "poor man's crop". It is a good source of minerals like phosphorus, iron and vitamins especially the B complex.

Comment [H8]: Add more recent references.

Brinjal is also valued for its medicinal properties and has got decholesterolizing property, primarily due to the presence of 65.1% linoleic and lenolenic poly-unsaturated fatty acids present in flesh and seeds. Presence of magnesium and potassium salts in fruits also impact of decholesterolizing action (Bhat, 2011).

Comment [H9]: Add a new statistic for cultivated area and production (Yield).

FYM consist of partially rotted straw of plants, urine and feces. It improves soil physical properties such as water holding capacity, erosion stability and gas exchange (Nyangani, 2010). FYM not only supplies a lot of macro and micro nutrients to the soil, but also improve

the soil physical, chemical and biological properties. Conventional FYM contains about 0.73% N, 0.18% P and 0.71% K (Tolessa and Friesen, 2001).

Vermicompost increases the surface area provides strong absorbability and retention of, nutrients as well and retains more nutrients for a longer period of time. [vermicompost](#) [Vermicompost](#) enhance the nutrient uptake by the plants by increasing the permeability of root cell membrane, stimulating root growth and increasing proliferation of root hairs (Pramanik *et al.*, 2007). Vermicompost being a stable fine granular organic matter was done based on plant requirements once a week. When added to clay soil loosens the soil and improves the passage to the enter of air for increase the fruit yield. The increase in nitrogen as found in poultry manure has its profound effect on the vegetative development of plants and ensures healthy and vigorous growth (Aliyu, 2000). Similarly Dauda *et al.* (2005) reported that the application of nitrogen, a major component of poultry manure has been reported to improve the yield of egg plant.

Zinc is an essential component of a number of enzymes *i.e.*, dehydrogenase, aldolase, isomerases, proteinase, peptidase and phosphohydrolase (Mousavi, 2011). It is directly involved in the synthesis of indol acetic acid (IAA) and proteins. The principal function is a metal activator of enzymes in [plants](#). Zinc deficiency may be related to weather conditions, as it increases in cold and wet weather, which might be due to the limited root growth in cool soils, or reduced activity of microorganisms and release of zinc from organic materials (Alam *et al.*, 2010 and Abdou *et al.*, 2011).

Comment [H10]: Add reference

Boron helps in the absorption of water and carbohydrate metabolism translocation of carbohydrates in plants, DNA synthesis in meristem, cell division and elongation, active salt absorption, fertilization, water relation and photosynthesis and involves indirectly in metabolism of nitrogen, phosphorous, fat and hormones (Haque *et al.*, 2011). Due to the lack of boron, there is hypertrophy, degeneration and disintegration of cambium cells in the meristematic tissues. Its deficiency may cause sterility, small fruit size, and poor yield (Davis *et al.*, 2003) and affects translocation of sugar, starches, nitrogen and phosphorus, synthesis of amino acids and proteins. Iron ([Fe](#)) is indispensable for chlorophyll synthesis. It acts as an oxygen carrier and is a constituent of certain enzymes and proteins. It has an important role in carotenoid synthesis in brinjal indirectly improving the quality of [brinjal](#).

Comment [H11]: Add more references in this paragraph.

MATERIALS AND METHODS

The experiment was laid out at Horticulture Farm, S.K.N. College of Agriculture, Jobner during *Kharif* season 2018 under Agro-Climatic Zone-III A (Semi-Arid Eastern Plains). Geographically, Jobner is situated in Jaipur district of Rajasthan at 26° 05' North latitude, 75° 58' East longitudes and an altitude of 427 metres above mean sea level. The experiment was carried out in randomized block design with 8 treatments and each replicated thrice.

Table 1: Details of treatments used in the study

Treatments	Symbols
A. Organic Manure	
(i) Control	M ₀
(ii) Fym (40 t/ ha)	M ₁
(iii) Vermicompost (6.6 t /ha)	M ₂
(iv) Poultry (6.6 t /ha)	M ₃
B. Micronutrient	
(i) Control	N ₀
(ii) Zn (0.5%) foliar spray	N ₁
(iii) Fe (0.5%) foliar spray	N ₂
(iv) B (0.3%) foliar spray	N ₃

Formatted Table

The plot size was 2.25 m × 2.40 m and spacing followed was 60 cm × 45 cm. The treatments were organic manures [control, FYM (40 t/ha), vermicompost (6.6 t/ha), poultry manure (6.6 t/ha)] and micronutrients [control, Zn (0.5%), Fe (0.5%) and B (0.3%) foliar spray]. Four weeks old healthy seedlings were transplanted on 21 July, 2018. Light irrigation was given after transplanting. All cultural practices were followed regularly during crop growth. The observations were recorded on TSS, ascorbic acid, protein content, fruit yield per plot and fruit yield per hectare. The data on the growth and yield were statistically analysed according to the method suggested by Fisher and Yates (1959).

Table 2: Physico-chemical characteristics of the experimental soil

S. No.	Parameters	Value	Method adopted
A. Mechanical analysis			
1.	Coarse sand (%)	26.5	International pipette method (Piper, 1950)
2.	Fine sand (%)	58.4	-do-
3.	Silt (%)	9.4	-do-
4.	Clay (%)	8.3	-do-
5.	Textural class	Loamy sand	USDA, Triangle (Soil survey Piper, 1950)

Formatted Table

B.	Physical analysis (%)		
1.	Field capacity (%)	10.8	Method No. 33, USDA Hand Book No. 60 (Richards, 1954)
2.	Particle density (Mg /m ³)	2.65	Method No. 39, USDA Hand Book No. 60 (Richards, 1954)
3.	Permanent wilting point (%)	3.4	Method No. 31, USDA Hand Book No. 60 (Richards, 1954)
4.	Bulk density (Mg /m ³)	1.50	Method No.31, USDA Hand Book No. 60 (Richards, 1954)
5.	Porosity (%)	42.59	Method No.31, USDA Hand Book No. 60 (Richards, 1954)
C.	Chemical analysis		
1.	Organic carbon (%)	0.16	Walkely and Black rapid titration method (Piper,1950)
2.	Available Nitrogen (Kg /ha)	146.64	Alkaline permagnate method (Subhiah and Asija, 1956)
3.	Available Phosphorus (Kg /ha)	18.52	Olsen's method (Olsen's <i>et al.</i> , 1954)
4.	Available Potassium (Kg /ha)	155.54	Flame photometer method (Metson, 1956)
5.	Available Sulphur (Mg /kg)	8.42	Turbidometric method (Chesnin and Yien,1951)
6.	E _{Ce} of saturation extract of soil at 25 ⁰ C (ds /m)	0.90	Method No. 4, USDA Hand Book No. 60 (Richards, 1954)
7.	pH (1:2 soil water suspension)	8.2	Method No. 21(b), USDA Hand Book No. 60 (Richards, 1954)

Results and discussion

Comment [H12]: All results require further discussion with recent previous studies, whether confirmatory or contradictory with them.

The results of the present investigation revealed that the response of organic manures and micronutrients were non significantly affected the TSS content of fruit. The maximum ascorbic acid (7.18 mg /100 g) and protein content (1.31%) were found under treatment vermicompost @ 6.6 t /ha (M₂), which was statistically at par with the treatment poultry manure @ 6.6 t /ha (M₃). This might be due to improved nutrient availability environment in the root zone and soil plant system. These results are in accordance with the results reported by Patil *et al.* (2004), Choudhary *et al.* (2007), Kashyap *et al.* (2014), and Kumar, (2016).

Micronutrients also significantly affected the ascorbic acid content and protein content. The maximum ascorbic acid content (7.27 mg /100 g) and protein content (1.33%) were recorded under treatment Zn @ 0.5 per cent (N₁). This might be due to accelerated activity of ascorbic acid oxidase enzyme in presence of these micronutrients (Fe, Zn). The role of Zinc in enhancement of ascorbic acid content may be due the ability of Cu uptake, which forms part of the enzyme ascorbic acid oxidase. These results are in agreement with the results obtained by Selvi *et al.* (2004), Savitha *et al.* (2010) and Singh *et al.* (2014a).

The maximum fruit yield per plot (17.13 kg /plot) and fruit yield (317.20 q /ha) were recorded under the treatment vermicompost @ 6.6 t /ha (M₂), which was statistically at par with poultry manure @ 6.6 t /ha (M₃), while, minimum fruit yield per plot (11.52 kg /plot) and fruit yield (213.40 q /ha) were observed in treatment M₀ (control). Increase in yield with application of organic manures was due to accelerated mobility of photosynthates from the source to the sink, released or synthesized due to the organic sources of fertilizers. These results are in accordance with the results reported by Ullal *et al.* (2008), Chumei *et al.* (2014) and Kashyap *et al.* (2014).

Table 3: Effect of organic manures and micronutrients on yield and quality of brinjal.

Treatments	TSS (%)	Ascorbic acid (mg /100g)	Protein content (%)	Fruit yield per plot (kg/plot)	Fruit yield (q/ha)
Organic Manures					
M ₀ -Control	4.29	6.08	1.12	11.52	213.40
M ₁ - FYM (40 t /ha)	4.65	6.57	1.22	14.36	265.87
M ₂ - Vermicompost @ 6.6 t /ha	5.19	7.18	1.31	17.13	317.20
M ₃ - Poultry Manure @ 6.6 t /ha	5.02	6.97	1.29	16.05	297.21
SEm _±	0.10	0.19	0.03	0.38	7.09

CD (P= 0.05)	0.30	0.55	0.08	1.11	20.47
Micronutrients					
N ₀ - Control	4.37	6.19	1.15	12.02	222.59
N ₁ - Zn (0.5%) foliar spray	5.24	7.27	1.33	17.39	321.99
N ₂ - Fe (0.5%) foliar spray	4.85	6.71	1.24	15.00	277.84
N ₃ - B (0.3%) foliar spray	4.69	6.62	1.22	14.65	271.25
SEm±	0.10	0.19	0.03	0.38	7.09
CD (P= 0.05)	0.30	0.55	0.08	1.11	20.47

Application of micronutrients significantly increased fruit yield and yield attributes and the maximum fruit yield/plot (17.39 kg /plot), fruit yield (321.99 q/ha) recorded under the treatment Zn @ 0.5 per cent (N₁), which was significantly superior to other treatments.




Figure 1: fruit yield per hectare for different treatments in brijal

While, the minimum fruit yield/plot (12.02 kg /plot) fruit yield (222.59 q /ha) were observed in treatment (N₀) control. This might be attributed due to enhanced photosynthetic activity, resulting in increased production and accumulation of carbohydrates and favorable effect on vegetative growth and retention of flowers and fruits, which might have increased number and weight of fruits ultimately increased fruit yield per hectare Tawab *et al.* (2015), Kadari, *et al.* (2015), Pandav *et al.* (2016) and Swetha *et al.* (2018).

CONCLUSION

Comment [H13]: Rewrite this section, mention the best obtained results.....

From the above findings it is concluded that the use of vermicompost @ 6.6 t/ha and micronutrient Zn @ 0.5 % resulted in increased yield and quality of brinjal as compared to the control in brinjal under Agro-Climatic Zone-III A (Semi-Arid Eastern Plains).

REFERENCES

- Abdou, A.S., Al-Darwish, F.H., Saleh, M.E., El-Tarabily, K.A., Azirun, M.S. and Rahman, M.M. 2011. Effects of elemental sulphur, phosphorus, micronutrients and Paracoccusversutus on nutrients availability of calcareous soils. *Australian Journal of Crop Science*, 5 (5): 554-561.
- Alam, M.N., Abedin, M.J. and Azad, M.A.K. 2010. Effect of micronutrients on growth and yield of onion under calcareous soil environment. *International Research Journal of Plant Science*, 1 (3): 56-61
- Aliyu, L. 2000. The effect of organic and mineral fertilizer on growth, yield and composition of pepper (*Capsicum annum* L). *Biological Agriculture and Horticulture*, 18: 29–36.
- Bhat, K.L. 2011. *Brinjal*. Daya Publishing House, New Delhi.
- Haque, M.E., Paul, A.K. and Sarker, J.R. 2011. Effect of nitrogen and boron on the growth and yield of tomato (*Lycopersicon esculentum* Mill.). *International Journal of Bio-resource and Stress Management*, 2: 277-282.
- Kashyap, S., Kumar, S., Maji, S. and Kumar, D. 2014. Effect of Organic manures and inorganic fertilizers on growth, yield and quality of brinjal (*Solanum melongena* L.) cv. Pant Rituraj. *International Journal Agriculture Science*, 10 (1): 305-308.
- Choudhary, M., Soni A.K. and Jat R.G. 2007. Effect of organic and Inorganic Sources of Nutrient on Quality of brinjal (*Solanum melongena* L.) cv. Pusa Uttam. *Haryana Journal Horticulture Science*, 36 (1 & 2): 118-119.
- Chumei, S.P. Kanaujia and Singh V.B. 2014. Integrated Nutrient Management in Brinjal. *Progressive Agriculture*, 13 (1): 106-113.
- Dauda, S.N., Aliyu, L. and Chiezey, U.F. 2005. Effect of seedling age at transplanting and poultry manure on fruit yield and nutrients of garden egg (*Solanum melongena* L.). *Journal of Tropical life Science*, 5: 38–41.

Davis, J.M., Sanders, D.C., Nelson, P.V., Lengnick, L. and Sperry, W.J. 2003. Boron improves the growth, yield, quality and nutrient content of tomato. *Journal of the American Society for Horticultural Science*, 128: 441-446.

Fisher, R.A. 1959. Statistical methods for research workers. Oliver and Boyd Edinburgh, London.

Comment [H14]: Rearrange

Kadari, I.A., Dheware, E.M., Waghmare, B.D., Mingire, S.S. and Dhawale, K.N. 2015. Influence of micronutrients and bio fertilizers on growth, yield and quality attributes of tomato (*Lycopersicon esculentum* Mill.). *Green Farming*, 6 (1): 74-76.

Kumar V. 2016. Use of Integrated nutrient management to enhance soil fertility and crop yield of hybrid cultivar of brinjal (*Solanum melongena* L.) under field conditions. *Advances in Plant and Agriculture Research*, 4 (2): 1-9.

Mousavi, S.R. 2011. Zinc in crop production and interaction with phosphorus. *Australian Journal of Basic and Applied Sciences*, 5 (9): 1503-1509.

Nyangani, E.T. 2010. Effect of combined application of organic manure and chemical fertilizers on soil properties and crop yields. Department of Agricultural Education. Taraba State, Nigeria. *Journal Science Technology & Environment Education*, 3: 28-30.

Pandav, A.K., Manoj Kumar Nalla, Aslam, T., Rana, M. K. and Bommesh, J. C. 2016. Effect of foliar application of micronutrients on growth and yield parameters in eggplant cv. HLB 12. *Environment and Ecology*, 35 (3): 1745-1748.

Patil, M.B., Mohammed, R.G. and Ghadge, P.M. 2004. Effect of organic and inorganic growth, yield and quality of tomato. *Journal Maharashtra Agriculture University*, 29 (2): 124-127.

Pramanik, P., Ghosh, G.K., Ghosal, P.K. and Banik, P. 2007. Changes in organic-C, N, P and K and enzyme activities in vermicompost of biodegradable organic wastes under liming and microbial inoculants. *Journal Biores Technilogy*, 98 : 2485-2494.

Saravaiya, S.N., Patel, N.B., Ahir, M.P., Patel, N.M. and Desai, K.D. 2010. Integrated Nutrient Management (INM) approach for brinjal (*Solanum melongena* L.) and other solanaceous vegetables-a review. *Agriculture Review*, 31(2): 79-92.

- Savitha, H.R., Bidari, B.I., Shashidhara, G.C., Kalyana Murthy, K.N., Poornima, D.S. and Mohan, K.S. 2010. Effect of Fe-EDTA on growth, yield and quality of red chilli (*Capsicum annuum* L.). *International Journal of Agricultural Sciences*, 6 (2): 531-533.
- Selvi, D., Thiageswari, S., Santhy, P. and Kannan, B.R. 2004. Fruit yield and nutrient uptake by brinjal due to integrated nutrient management in an inceptisol. *Journal of Maharashtra Agriculture University*, 29 (2): 220-223.
- Singh, N.K., Sharma, T.R., Bisen, N.K. and Deshmukh, K.K. 2014a. Optimization of quantity of foliar spray of boron and zinc in chilli for Kymore plateau and Satpura hills of Madhya Pradesh. *Vegetable Science*, 41 (1): 66-67.
- Swetha K., Saravanan S. and Prasad V.M. 2018. Effect of micronutrients on plant growth, flowering and yield of Tomato (*Solanum lycopersicum* L.) cv. "PKM-1". *International Journal of Agriculture Sciences*, 10 (7): 5734-5736.
- Tawab, Saleha, Gohar, Ayub, Tawab, Faiza, Khan, Owais, Nadia Bostan, Ghazala Ruby, Shawana Ahmad and Ume Kalsoom Afridi. 2015. Response of brinjal (*Solanum melongena* L.) cultivars to zinc levels. *ARPJN Journal of Agricultural and Biological Science*, 10 (5): 172-178.
- Ullah, M.S., Islam, M.S., Islam, M.A. and Haque, T. 2008. Effects of organic manures and chemical fertilizers on the yield of brinjal and soil properties. *Journal Bangladesh Agriculture University*, 6 (2): 271-276.