

Original Research Article

Pre- Harvest Foliar Spray of Zinc, Calcium, Chitosan and Salicylic Acid on Quality Standards of Indian Ber (*Zizyphus mauritiana* Lamk.) cv. Gola

ABSTRACT

Aim:The present experiment was designed to find out the effect of foliar spray of zinc, calcium, chitosan and salicylic acid on fruit quality of ber.

Place and Duration of Study:This experiment was conducted at Horticulture farm, SKN College of Agriculture, Jobner in the newly established orchard of Indian jujube cv. Gola during the two consecutive year 2022 and 2023.

Study design and Methodology:The present experiment laid out in factorial randomized block design with three replications. Factor A consists of 12 level of zinc, calcium, chitosan and salicylic acid treatments along with the control and factor B consists three foliar sprays i.e., single foliar spray at pea size stage, colour turning stage of fruits and double foliar spray at pea size and colour turning stage of fruits (green to yellow). Various quality parameters namely TSS, acidity, ascorbic acid, total, reducing and non-reducing sugars are observed as per standard procedures.

Results:Plants sprayed with $ZnSO_4 @ 4g/L + CaCl_2 @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L$ recorded highest TSS (19.05°Brix), minimum acidity (0.415%) TSS: acid ratio (45.93%), ascorbic acid (89.08mg/100g), total sugars (10.25%), reducing sugars (6.68%) and non-reducing sugars (3.39%) were significantly more in the treated fruits in comparison to untreated fruits. These quality parameters were observed maximum with double foliar spray of nutrients and agrochemicals at pea size stage and colour turning stage (S_3).

Conclusion:It is concluded that $ZnSO_4 @ 4g/L + CaCl_2 @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L$ is effective to improve the quality of ber fruits.

Keywords: Indian jujube; Zinc; Calcium; Chitosan; Salicylic acid; Quality

1. INTRODUCTION

Ber (*Zizyphus mauritiana* Lamk.) is one of the most ancient and common fruits of India and it is cultivated all over the country. Being a hardy fruit, it can be grown even on inferior and marginal lands. Ber is indigenous to an area stretching from India to China and Malaysia. In view of good returns from ber, its cultivation is becoming increasingly popular in Gujarat and other states of India, especially in Punjab, Haryana, and Rajasthan. Gola is the most common cultivar being farmed (Surani *et al.*, 2018). The Gola variety is a well-liked early type. The fruit has a medium size and an oblong to spherical shape. Its color changes from green to golden yellow as it ripens. Fruit typically weighs between 16 and 20 g on average (Chattopadhyay, 2012). A huge loss of horticultural produce takes place during storage. Inadequate means of losses during storage due to poor handling and poor post-harvest management practices. Ber can be cultivated even under constraints of irrigation and fertilization. The essential element of fruit production are bioregulators. These regulators

can be applied to fruits pre-harvest to improve their physical and biochemical characteristics and preserving quality.

Zinc is an essential micro element for plants, being involved in many enzymatic reactions and is necessary for their good growth and development. It is also involved in regulating the protein and carbohydrate metabolism. Its availability to plants is reduced in high pH soils. Fruits generally have a considerable amount of Calcium in their tissues. The Ca application makes the middle lamella of fruit cell wall thicker by increased deposition of calcium pectate and so, maintains the firmness of fruit (Sudha *et al.* 2007). Calcium has long been associated with regulation of the ripening and post-harvest storage life of fruits. Specially, maintenance of relatively high Ca concentrates in fruit tissue result in a slower rate of ripening, as seen in lower respiration rate, reduced ethylene production and slower softening of fruit flesh. Combined with pectin, the “glue” that holds plant cells together, calcium forms a pectate salt that helps to keep the cell wall sturdy and rigid. Chitosan applied topically at various phases of fruit development had a major impact on the postharvest life of jujube fruits during storage. Treatments with chitosan function as barriers, preventing the transfer of water, shielding fruit skin from mechanical damage, and sealing minor wounds to postpone dehydration (Ribeiro *et al.*, 2007). Salicylic acid is an additional phenolic endogenous growth regulator that helps control a number of physiological functions in plants, including transpiration, ion absorption, stomata closure, and the suppression of ethylene production (Shafiee *et al.*, 2010). Keeping in mind the importance of nutrients and agrochemicals current study was planned to improve the quality of ber fruit by pre-harvest exogenous spray of different concentrations of zinc, calcium, chitosan and salicylic acid at various growth stages under semi-arid condition of Rajasthan.

2. MATERIAL AND METHODS

2.1 Experimental site and treatments detail

This experiment was conducted at Horticulture farm, SKN College of Agriculture, Jobner in the newly established orchard of Indian jujube (*Zizyphus mauritiana* Lamk.) cv. Gola during 2022 and 2023. Factor A consists of 12 level of zinc, calcium, chitosan and salicylic acid treatments along with the control and factor B consists three foliar sprays i.e., single foliar spray at pea size stage, colour turning stage of fruits and double foliar spray at pea size and colour turning stage of fruits (green to yellow). The treatment details of zinc, calcium, chitosan and salicylic acid along with the control i.e., T₀ (Control), T₁ (Zinc sulphate @ 4 g / L), T₂ (Calcium chloride @ 2 g / L), T₃ (Chitosan @ 1.5 g / L), T₄ Salicylic acid @ 1 g / L), T₅ (Zinc sulphate @ 4 g / L + Calcium chloride @ 2 g / L), T₆ (Zinc sulphate @ 4 g / L + Chitosan @ 1.5 g / L), T₇ (Zinc sulphate @ 4 g / L + Salicylic acid @ 1 g / L), T₈ (Calcium chloride @ 2 g / L + Chitosan @ 1.5 g / L), T₉ (Calcium chloride @ 2 g / L + Salicylic acid @ 1 g / L), T₁₀ (Chitosan @ 1.5 g / L + Salicylic acid @ 1 g / L) and T₁₁ (Zinc sulphate @ 4 g / L + Calcium chloride @ 2 g / L + Chitosan @ 1.5 g / L + Salicylic acid @ 1 g / L) laid out in factorial randomized block design with three replications. The treatments were applied single foliar spray at pea size stage of fruits during first week of November and double foliar spray at pea size and colour turning stage of fruits (green to yellow) applied last week of December, 2022-23 and 2023-24 after recording initial (base) quality and parameters of ber fruits.

2.2 Quality attributes of ber fruit

For determination of total soluble solids were measured by placing a drop of juice on the prism of hand refractometer. The reading against the light was noted and presented in degree brix. For estimation of ascorbic acid, well homogenized 2 g of sample was taken in a beaker and volume was made up to 20 ml and 5 ml of HPO₃ is added and titrated with the

standard dye solution to a pink end-point which should persist for 15 seconds. The same procedure was repeated in random samples of fruits in both the year of experiment to draw an average value of ascorbic acid content. The ascorbic acid content of the sample was calculated by using the following formula (Ranganna 2000).

$$\text{Ascorbic acid (mg/100g)} = \frac{\text{Titre value} \times \text{dye factor} \times \text{volume made up}}{\text{Aliquot extract taken for estimation} \times \text{weight of sample taken for Estimation}} \times 100$$

Total and reducing sugars were estimated by lane and Eynon copper titration method and express in percentage (Ranganna, 1986). The percentage of non-reducing sugar was determined by subtracting the per cent reducing sugar from the per cent total sugar and multiplying the same with 0.95 as given below (Somogyi, 1952).

$$\text{Non-reducing sugars (\%)} = (\text{Per cent total sugar} - \text{per cent reducing sugar}) \times 0.95$$

2.3 Statistical analysis

The data were statistically analyzed as per analysis of variance technique as suggested by Panse *et al.* (1995). The significance of the treatments was tested through F test at 5 per cent level of significance. The critical difference CD was calculated to assess the significance of difference among the different treatments.

3. RESULTS AND DISCUSSION

Pre- harvest foliar spray of zinc, calcium, chitosan and salicylic acid on quality standards viz., TSS, titratable acidity, TSS: acid ratio and ascorbic acid of ber (*Zizyphus mauritiana* Lamk.) cv. Gola is presented in (Table 1).

The maximum TSS content in fruits (19.05%) was recorded with the application of treatment (ZnSO₄ @ 4g/L + CaCl₂ @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L) followed by T₉ (CaCl₂ @ 2g/L + Salicylic acid @ 1g/L 18.90%), T₇ (ZnSO₄ @ 4g/L + Salicylic acid @ 1g/L 18.83%), T₅ (ZnSO₄ @ 4g/L + CaCl₂ @ 2g/L 18.69%) and minimum TSS content in fruits (16.80) was noted under control. The maximum total soluble solid content (18.05 °Brix) was observed with double foliar spray of nutrients and agrochemicals at pea size stage and colour turning stage (S₃) and it was found at par with foliar spray at S₁ stage. The foliar spray of nutrients and agrochemicals at colour turning stage (S₂) registered lowest total soluble solid (17.38 °Brix). A significant decrease in titratable acidity was observed in ZnSO₄ @ 4g/L + CaCl₂ @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L (0.16%) followed by the T₉, T₇ and T₅ and highest titratable acidity was observed in fruits of controlled treatment (0.469%). The application of nutrients and agrochemicals at different fruit development stages had not shown any significant effect on the titratable acidity in ber fruits.

The maximum TSS: acid ratio (45.93) was recorded with combined application of ZnSO₄ @ 4 g/l + CaCl₂ @ 2 g/l + Chitosan @ 1.5 g/l + Salicylic acid @ 1 g/l (T₁₁) followed by this treatment T₉, T₇ and T₅. However, minimum TSS: acid ratio of (36.32) was recorded under control. Application of double foliar spray of nutrients and agrochemicals exhibited maximum TSS: acid ratio (41.69) in fruits. However, this treatment (S₃) was found statistically at par with treatment S₁. The foliar spray at S₂ recorded lowest TSS: acid ratio (38.90) during the experimentation.

ZnSO₄ @ 4g/L + CaCl₂ @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L significantly increases the ascorbic acid content of ber fruits. Highest ascorbic acid content was observed in fruits treated with ZnSO₄ @ 4g/L + CaCl₂ @ 2g/L + Chitosan @ 1.5g/L + Salicylic acid @ 1g/L that's 89.08 mg/100g followed by T₉ (85.77 mg/100g), T₇ (85.63

mg/100g) and T₅ (85.31 mg/100g). Whereas, the lowest contents were observed in control (74.63 mg/100g). However, the maximum ascorbic acid content (83.42 mg/100g) was observed with double foliar spray of nutrients and agrochemicals at pea size stage and colour turning stage (S₃) and it was found statistically at par with S₁(82.75 mg/100g). The treatment combination S₂ recorded minimum ascorbic acid content (81.96 mg/100g) the investigation.

Table 1 Pre- Harvest Foliar Spray of Zinc, Calcium, Chitosan and Salicylic Acid on TSS, titratable acidity, TSS: acid ratio and ascorbic acid of ber fruit.

Treatments	TSS (^o Brix)	Titratable acidity (%)	TSS: acid ratio	Ascorbic acid (mg/100g)
Nutrients and agrochemicals				
T ₀ Control	16.80	0.469	36.32	74.63
T ₁ ZnSO ₄	16.90	0.457	37.03	80.38
T ₂ CaCl ₂	16.94	0.457	37.15	80.72
T ₃ Chitosan	16.83	0.465	36.26	76.44
T ₄ Salicylic acid	16.94	0.447	37.96	81.15
T ₅ ZnSO ₄ + CaCl ₂	18.69	0.421	44.47	88.07
T ₆ ZnSO ₄ + Chitosan	17.43	0.448	38.97	81.53
T ₇ ZnSO ₄ + Salicylic acid	18.83	0.418	45.05	88.32
T ₈ CaCl ₂ + Chitosan	17.77	0.450	39.57	81.65
T ₉ CaCl ₂ + Salicylic acid	18.90	0.417	45.36	88.80
T ₁₀ Chitosan+ Salicylic acid	17.94	0.434	41.55	81.77
T ₁₁ ZnSO ₄ + CaCl ₂ + Chitosan + SA	19.05	0.415	45.93	89.08
SEm_±	0.21	0.00	0.61	0.48
CD (P = 0.05)	0.59	NS	1.70	1.35
Different Stage of Foliar Spray				
S ₁	17.83	0.440	40.81	82.75
S ₂	17.38	0.449	38.90	81.96
S ₃	18.05	0.435	41.69	83.42
SEm_±	0.11	0.00	0.30	0.24
CD (P = 0.05)	0.29	NS	0.85	0.68

It is evident from data presented in Table 2 related to application of nutrients and agrochemicals showed significantly effect on total, reducing and non-reducing sugars in fruits. Significantly higher total sugars was observed in ZnSO₄ @ 4g/L + CaCl₂ @ 2g/L + Chitosan @ 1.5g/L +Salicylic acid @ 1g/L treatment (10.25%). Controlled treatment shows minimum total sugars among other treatments (8.16%). However, the highest total sugars of (9.54 %) was observed with double foliar spray of nutrients and agrochemicals at pea size stage and colour turning stage (S₃) and it was found statistically at par with treatment S₁(9.32 %). While, the treatment combination S₂ recorded lowest total sugars content (8.96 %) during the experimentation.

ZnSO₄ @ 4g/L + CaCl₂ @ 2g/L + Chitosan @ 1.5g/L +Salicylic acid @ 1g/L significantly increases reducing and non-reducing sugars (6.52%, 4.23% respectively) followed by the fruits treated with T₉, T₇ and T₅ and minimum values were observed in control treatment. However, the maximum reducing (6.46%) and non-reducing (3.01%)

sugars was observed with double foliar spray of nutrients and agrochemicals at pea size stage and colour turning stage (S₃) and this treatment (S₃) was found statistically at par with treatment S₁. Whereas, minimum reducing (6.17%) and non-reducing (2.56%) sugars was recorded with foliar spray of nutrients and agrochemicals done at colour turning stage (S₂).

Table 2Pre- Harvest Foliar Spray of Zinc, Calcium, Chitosan and Salicylic Acid on total, reducing and non-reducing sugars offer fruit.

Treatments	Total sugars (%)	Reducing sugars (%)	Non-reducing sugars (%)
Nutrients and agrochemicals			
T ₀ Control	8.16	5.92	2.13
T ₁ ZnSO ₄	8.84	6.15	2.56
T ₂ CaCl ₂	8.92	6.23	2.55
T ₃ Chitosan	8.64	6.07	2.44
T ₄ Salicylic acid	9.01	6.19	2.68
T ₅ ZnSO ₄ + CaCl ₂	9.93	6.56	3.21
T ₆ ZnSO ₄ + Chitosan	9.01	6.24	2.64
T ₇ ZnSO ₄ + Salicylic acid	10.05	6.58	3.30
T ₈ CaCl ₂ + Chitosan	9.10	6.28	2.68
T ₉ CaCl ₂ + Salicylic acid	10.16	6.63	3.35
T ₁₀ Chitosan+ Salicylic acid	9.17	6.31	2.71
T ₁₁ ZnSO ₄ + CaCl ₂ + Chitosan + SA	10.25	6.68	3.39
SEm±	0.17	0.10	0.17
CD (P = 0.05)	0.47	0.27	0.48
Different Stage of Foliar Spray			
S ₁	9.32	6.33	2.84
S ₂	8.96	6.17	2.56
S ₃	9.54	6.46	3.01
SEm±	0.08	0.05	0.09
CD (P = 0.05)	0.23	0.13	0.24

The application of zinc, CaCl₂, chitosan and SA increased TSS content in the fruits, which might be due to the enhancement of auxin synthesis. Auxin, in turn, enhances the synthesis of metabolites and their rapid translocation from one part of the plant to developing fruits (Alrawi *et al.*, 2012 and Badway *et al.*, 2019). The improvement in fruit quality might also be due to zinc's role in enzymatic reactions, such as hexokinase, and its involvement in carbohydrate formation and protein synthesis (Pamila *et al.*, 1992). The reason for decreased TA is due to breakup of fermentation of acids into sugars during respiration process. An increase in the TSS: acid ratio is likely due to an increase in TSS content and a decrease in acidity (Abdel rhman *et al.*, 2017). Zinc, CaCl₂, chitosan and SA application also increased the ascorbic acid content, which appears to be due to increased growth and the availability of more metabolites for ascorbic acid synthesis. The possible reason for the increase in total sugar and reducing sugars in fruits from foliar sprays of minerals and agrochemicals might be the conversion of starch and acid into sugars, along with the continuous mobilization of sugars from leaves to fruits (Martinez *et al.*, 2022). Treatments that delayed the start of fruit senescence and the conversion of complex carbohydrates into simple sugars may have contributed to the greater non-reducing sugar levels in treated samples (Sajid *et al.* 2020).

4. CONCLUSION

The findings of this study demonstrate that as compared with individual applications of zinc, calcium, chitosan or salicylic acid, a combination of these elicitors had a much superior impact on fruit quality aspects of ber fruits. The present study is noteworthy because it offers evidence that a combination of these elicitors may enhance fruit quality both during the postharvest storage period and at the time of harvest. Future research should concentrate on optimizing the concentrations and combinations that might be used commercially on ber fruit from different crops or cultivars in order to increase the fruit's marketability.

REFERENCES

- Abdel Rhman, I.E, Attia M.F, Elhady, E.S., Haggag, L.F. Effect of foliar spraying of some antioxidants and micronutrients on yield, fruit quality and leaf mineral content of manfaloty pomegranate trees (*Punica granatum* L.) grown in a calcareous soil. *Middle East Journal of Applied Science*, 2017; 7(4): 713-725.
- Alrawi, W., Jassim, N.A., Al-Hadethi, M.E. Effect of iron and zinc spray on yield, growth and content of the seeds of apricot amygdalin. *Egyptian Journal of Applied Sciences*, 2012; 27(4): 173-187.
- Badawy, I.F., Abouzaid, E.A. and Hussein, E.M. Cracking and fruit quality of "Manfalouty" pomegranate as affected by pre-harvest of chitosan, calcium chloride and gibberellic acid spraying. *Middle East Journal*, 2019; 8(3): 873-882.
- Surani, BR., Patel, AP. and Yadav, L. Effect of post-harvest application of chemicals on biochemical parameters of different Ber (*Ziziphus mauritiana* Lamk.) varieties during storage. *International Journal of Chemical Studies*, 2018; 6(4): 2804-280
- Chattopadhyay TK. *A Text Book on Pomology (Subtropical Fruits)*. Kalyani Publishers, New Delhi, 2012.
- Martínez-Camacho, J. E., Guevara-González, R. G., Rico-García, E., Tovar-Pérez, E. G., and Torres-Pacheco, I. Delayed senescence and marketability index preservation of blackberry fruit by pre-harvest application of chitosan and salicylic acid. *Frontiers in Plant Science*, 2022; 13(04): 796393.
- Pamila, S., Chatterjee, S.R. and Deb, D.L. Seed yield, harvest index, protein content and amino acid composition of chickpea as affected by sulphur and micronutrients. *Annals of Agriculture Research*, 1992; 3(1): 7-11.
- Panse V G and Sukhatme P V. *Statistical methods for agricultural workers*, 4th edition. ICAR, New Delhi, 1995; 58-92.
- Ranganna, K., Yatsu, F.M., Hayes, B.E., Milton, S.G. and Jayakumar, A. Butyrate inhibits proliferation-induced proliferating cell nuclear antigen expression (PCNA) in rat vascular smooth muscle cells. *Molecular and Cellular Biochemistry*, 2000; 205:149-161.

Ranganna, S. Handbook of analysis and quality control for fruit and vegetable products. Tata McGraw-Hill Education, 1986; 670.

Ribeiro, C., Vicente, A.A., Teixeira, J.A. and Miranda, C. Optimization of edible coating composition to retard strawberry fruit senescence. *Post-Harvest Biology and Technology*, 2007; 44(1): 63-70.

Sajid, M., Basit, A., Ullah, Z., Shah, S.T., Ullah, I., Mohamed, H.I. and Ullah, I. Chitosan-based foliar application modulated the yield and biochemical attributes of peach (*Prunus persica* L.) cv. Early Grand. *Bulletin of the National Research Centre*, 2020; 44(1):1-11.

Shafiee, M., Taghavi, T.S. and Babalar, M. Addition of salicylic acid to nutrient solution combined with post-harvest treatments (hot water, salicylic acid, and calcium dipping) improved post-harvest fruit quality of strawjuberly. *Scientia Horticulturae*, 2010; 124(1): 40-45.

Somogyi, M. Notes on sugar determination. *Journal of Biological Chemistry*, 1952; 195: 19-23.

UNDER PEER REVIEW