

# Zebrafish: a versatile experimental model for application in toxicological research

## ABSTRACT

The field of scientific research has advanced exponentially in recent years, with more sophisticated techniques emerging and the diversity of experimental models also growing. One species that has stood out in animal experimentation is the Zebrafish (*Danio rerio*), an aquatic vertebrate that has a fully sequenced genome, anatomical similarities to mammals, high fecundity, rapid embryonic development outside the body and with transparent embryos, as well as low maintenance costs. This experimental model is a favorable alternative for neurobehavioral and genetic studies, but especially for the toxicological field, where it has been widely applied due to all its characteristics that allow toxicity to be analyzed efficiently, while also meeting the 3R's criteria. Since it is a convenient alternative model, its use in different methodologies and areas of study is growing.

*Keywords: Biomodel; Danio rerio; Embryonic development; Ecotoxicology; Toxicokinetics*

## 1. INTRODUCTION

Toxicology is a science dedicated to the study of various chemical substances and their generated effects, addressing and evaluating their harmful impacts on the environment and living organisms. This science is of great importance as it has a multidisciplinary nature, encompassing themes such as biology, chemistry, pharmacology, ecology, public health, among others [1]. In recent years, the field of toxicology has advanced exponentially, with the number of publications on research and reviews in the area following the same pattern [2].

This behavior has been observed for many decades, with toxicological studies, even if simple, being recorded since before the Middle Ages concerning poisons and toxic plants. This trajectory has evolved through discoveries of therapeutic agents and has matured scientifically into modern times, where numerous researchers are dedicated to understanding complex chemicals, ensuring food safety, producing pharmaceuticals, and herbal medicines, as well as gaining a deeper understanding of the effects of toxic compounds [3].

This progression is marked by technological advancements, where new tools and more refined techniques provide not only greater ease and speed in conducting existing processes but also the emergence of new approaches. Thus, technological development has begun to assist researchers beyond traditional systems, with the introduction of new procedures such as in vitro testing, nanotechnology, and even nanotoxicology [4]. Computational tools such as simulations (In silico), epigenetics, and toxicogenomics [5].

In relation to this, another methodology that has evolved is linked to experimental models. Ferreira et al. [6] state that an experimental model is the materialization of a part of reality through a simple representation that must present precision and similarity to what is intended to be analyzed. In line with this, laboratory animals are the most commonly used. This has been the case for many years, as they provide essential mechanisms for the development of scientific work, such as the opportunity to perform *in vivo* procedures and represent real and natural conditions. Additionally, physiological and genetic similarities, ease of manipulation, the ability to induce diseases, and well-established protocols for the use of animal models contribute to their relevance [7].

However, despite the notable laboratory rats and mice, other species have emerged as alternatives to replace them. These include fish, which have become significant allies in science as alternative models that demonstrate equal or even superior efficiency compared to traditional systems. Certainly, these animals respond to the requirements of the Animal Ethics Committee (CEUA) concerning the principle of the 3Rs.

The concepts of reduction, refinement, and replacement are indispensable elements in scientific research. They aim to ensure that advancements are linked to ethics, with the objective of minimizing suffering and ensuring the proper management of animals in experiments [8]. In this regard, the Zebrafish, emerges as one of the most commonly used samples when the methodology refers to the use of aquatic vertebrates.

In this context, *Danio rerio* is not new to the scientific community. Researchers have focused on this species since the 1950s, when biologist George Streisinger conducted genetic analyses on the animal. By 1990, a group of scientists in Oregon hosted a meeting regarding the characteristics of this model for researchers from around the world. In Brazil, the country has been one of the largest producers of research involving Zebrafish, and the trend is towards growth, as many topics related to toxicology and behavioral neuroscience are extensively addressed here, making this fish exceptionally well-suited for such studies [9].

In light of this, the present work aims to conduct an in-depth literature review on Zebrafish as an experimental model, its characteristics, effectiveness, and limitations, as well as its applications in the study of toxicology, providing a comprehensive perspective on the subject.

## **2. METHODOLOGY**

Searches were conducted in the databases PubMed, SciELO, and ScienceDirect using the following keywords: "Experimental fish model applied to toxicology," "Use of fish as an experimental model for toxicological studies," and "Zebrafish and *in vitro* experimental model." A total of 1,047 articles were found on the PubMed platform, 0 on SciELO, and 39,605 on ScienceDirect. Additionally, filters were applied in the ScienceDirect database to exclude encyclopedias, editorials, and reviews, selecting only research articles, review articles, and book chapters. Subsequently, the process of selecting and excluding articles was based on a review of titles and abstracts, eliminating those that did not address or deviated from the central theme of the article, namely, the use of fish in toxicological studies. In the end, 46 articles were included. Furthermore, the selected articles span a period of ten years, from 2014 to 2024.

## **3. RESULTS AND DISCUSSION**

### **3.1 CHARACTERIZATION OF ZEBRAFISH**

*Danio rerio*, commonly known as Zebrafish due to the striking stripes that run along the sides of its body, is a notable teleost fish, specifically belonging to the family Cyprinidae, with origins in South Asia [10]. This species has a small size of approximately 2 centimeters and can reach up to about 4 centimeters [11]. In terms of its habitat, these animals are freshwater fish and typically inhabit shallow, slow, or even stagnant waters [12].

Following this, Zebrafish can exist in three stages: larval, juvenile, and adult, and can live up to five years. Its development is notably rapid, with these periods measured in hours post-fertilization (hpf) or days post-fertilization (dpf). Total embryonic development occurs at 72 hpf, and it reaches adulthood by the end of 90 dpf [13]. Furthermore, this aquatic vertebrate exhibits diurnal behavior, and the spawning of its eggs typically occurs during the early hours in the presence of light; thus, the photoperiod is an important factor in the environments where they are kept, allowing for a stable environment and a regular circadian cycle [14].

Additionally, despite its natural habitat, this tropical fish is commonly sold commercially for use in personal aquariums but is primarily utilized in scientific research, living in breeding facilities.

### **3.2 ADVANTAGES VS. LIMITATIONS**

Regarding the use of Zebrafish as an experimental model, this fish presents various advantages that make it an excellent asset for research. Firstly, in experiments aimed at evaluating toxicity, the effects of a metabolite, or even those intending to generate new medications for human use, it is essential that the sample possesses a certain degree of compatibility so that the generated effects do not deviate from reality. In this regard, Zebrafish has its genome completely sequenced, as reported by Howe et al. [15] in 2013, with 70% homology to that of humans, allowing for the modeling of many human diseases in these animals [16].

Additionally, there are anatomical similarities with mammals, as this fish has structures such as the heart, kidneys, liver, pancreas, bones, muscles, and central nervous system (CNS). Furthermore, there is correspondence regarding the immune system, sensory system, and signaling profiles. In other words, the similarities are not limited to a single factor but are evident at physiological, molecular, anatomical, and genetic levels [17].

Regarding reproduction, females exhibit high fecundity [18] and are capable of producing approximately 200 to 300 eggs per week, with embryogenesis completed after 72 hpf and reaching new reproductive maturity by the end of 3 months [19]. This characteristic of rapid development is highly valuable for experimentation, as it allows researchers to avoid relying on extended growth periods, enabling them to observe all stages of the fish and quickly focus on tests and analyses.

In this regard, the process of embryonic maturation occurs *ex vivo*, meaning outside the organism, and the embryo is transparent during the early stages of development. This factor facilitates the visualization of responses, enabling the monitoring of inflammatory processes, imaging-based tracking of treatment impacts or the effects of a chemical compound, as well as the visibility of fluorescent markers and morphological changes [20, 21, 22].

Moreover, due to their small size as vertebrates, they can be stored together in larger quantities compared to other biomodels such as primates, while still avoiding stressful conditions. For this reason, their breeding and maintenance within laboratory aquariums become less costly [23]. Additionally, swimming profiles, social behavior, light-dark

preference, anxiety, and aggressiveness are observable in *Danio rerio* and can be utilized as assessment tools in neurotoxicological testing [24].

Despite this significant number of advantages, there are still some limitations regarding the use of this animal. The lack of standardization and firm establishment of Zebrafish strains is a concern, as this heterogeneity can lead to variations in experimental studies utilizing them. Their size also necessitates more refined and sensitive techniques. Since they develop externally and are protected by a chorion until approximately 72 hours post-fertilization (hpf), the exposure of the fish to the absorption of a compound may be indirect and relatively minimal. Lastly, the absence of organs such as lungs and the lack of proper limbs, joints, and mammary glands disqualify them from research that may involve these targets [25, 26].

### **3.3 TRADITIONAL MODEL AND ALTERNATIVE MODEL**

When discussing scientific research involving animal experimentation, the first species that come to mind are the well-known rats (*Rattus norvegicus*) and mice (*Mus musculus domesticus*). However, in contemporary science, questions have arisen regarding animal pain and suffering, leading to an increased need for the application of alternative models.

As a result, despite being proven effective, the need arose to establish a model that required low maintenance while maintaining efficacy. Zebrafish have been used in research for quite some time, with George Streisinger pioneering the field of Zebrafish research in the 1970s [27]. He sought a model that could be used in genetic studies and, as an aquarist, recognized the advantages of Zebrafish. Its rapid development, small size, and ease of maintenance were some of the characteristics that led to Zebrafish being considered a model organism [28].

It was around the 1930s that Zebrafish were introduced as a model organism for developmental and embryological studies. Consequently, it was applied as a genetic model following advances in genetic methodologies, including cloning, mutagenesis, and transgenesis [29]. Thus, Zebrafish were proposed as a model system for understanding and studying vertebrate development. Shortly thereafter, the Zebrafish genome was sequenced, revealing that 70% of human genes are homologous to those in Zebrafish [28].

From that point on, Zebrafish established themselves as a model for understanding human diseases.

### **3.4 APPLICATIONS IN TOXICOLOGY STUDIES**

Zebrafish as an experimental model represents a viable alternative to the mammalian models currently used in toxicity testing and other biological research. Since Zebrafish are easier and less expensive to house and care for than popular rodent models [21]. Additionally, the rapid development of Zebrafish larvae allows for the evaluation of both acute and chronic effects of substances on their organism [30]. Features such as their small size, transparent embryos, rapid external development, and high reproductive capacity [31] have contributed to making Zebrafish an attractive model for developmental studies.

### **3.5 CARDIOTOXICITY**

Although Zebrafish and humans have distinct structures for their cardiovascular systems: such as the two-chambered heart with one atrium and one ventricle in Zebrafish, compared to the four-chambered heart in humans-Zebrafish are still used as an alternative model for studying the cardiovascular system. This is due to their advantages, including low cost,

external development, transparent embryos, the possibility of genetic manipulation, and genetic homology with humans [27].

An example is the study by Arman and İşisağÜçüncü (2020), which evaluated the cardiotoxicity of Zebrafish embryos exposed to acrolein, a pollutant found in the environment that can be derived from the incomplete combustion of organic substances, industrial discharges, power plant emissions, among others. Another example is the work by Ming Li et al. [33], which analyzed muscone, one of the most physiologically active compounds of natural musk, frequently used to treat heart diseases. The results showed that the compound is toxic to Zebrafish embryos, causing mortality, reduced hatching rates, pericardial edema, and decreased heart rate.

Anthracyclines are an effective chemotherapy for antineoplastic treatment; however, cumulative cardiotoxicity is the primary side effect with a poor prognosis, and currently, there is no therapy that reverses anthracycline-induced cardiotoxicity. The article by Lu et al. [34] aimed to demonstrate the cardioprotective effect of Calycosin against anthracycline using Zebrafish as an embryonic model.

### **3.6 NEUROTOXICITY**

Neurotoxicity occurs when exposure to toxic substances alters the normal activities of the nervous system, including neural transmission, connectivity, and survival [35]. In the work of Faria et al. [36], it was demonstrated that fenitrothion, an organophosphate insecticide, caused neurotoxicity in fish after 24 hours of exposure. The toxicological effects were analyzed through behavior and the expression of *gap43a*, *gfap*, *atp2b1a*, *mbp*, and *syt1a*.

Another example is Azevedo et al. [37], whose study investigated the metabolic and synaptic deficiencies triggered by pyriproxyfen, a pesticide with various applications, including antiparasitic medications for pets and domestic/agricultural pest control.

Additionally, the study by Prabha et al. [38] evaluated the antioxidant and neuroprotective capacity of the RW20 peptide. This peptide is derived from histone acetyltransferases and was analyzed with the aim of being a protective agent against neurodegenerative diseases caused by oxidative stress.

### **3.7 GENOTOXICITY**

Genotoxicity generally refers to DNA damage, which includes single-strand breaks, double-strand breaks, and the formation of adducts, as well as alterations in the genes/proteins involved in repair. Zebrafish have been suggested as a suitable model for evaluating these effects [27]. Derikvandy et al. [39] used Zebrafish to analyze the toxicological effects of untreated wastewater from the ethyl alcohol industry. Their results showed that after 21 days of exposure to 2% wastewater, the expressions of the genes *Sod1*, *Gstp-1a*, and *Gpx1a* were increased in hepatocytes. Thus, the authors concluded that the analyzed water caused significant increases in the expression of antioxidant and detoxification-related genes in the liver of Zebrafish.

Chowdhury et al. [40] investigated the concentration of amoxicillin residue (AMXR) in raw and pasteurized milk samples from cows with mastitis. The study was conducted using the standard HPLC method compared to pure amoxicillin, examining the effect of the residue on developmental toxicity and genotoxicity in Zebrafish. The results showed DNA damage caused by AMXR.

### 3.8 HEPATOTOXICITY

The liver is the largest gland in the bodies of vertebrates. It plays an important role in the metabolism of carbohydrates, fats, proteins, vitamins, and hormones, as well as in detoxification regulation. Consequently, the liver is the primary target organ for toxic reactions. Hepatic injury due to toxic substances, or hepatotoxicity, is primarily caused by drugs and poisons that induce liver damage, including oxidative stress and apoptosis [41].

An example is the work of Hu et al. [42], which investigates Iprodione, a broad-spectrum fungicide used for the early control of fruit trees and vegetables. Due to rainfall, it may pose a risk of toxicity to non-target individuals, and the authors specifically focus on liver development. They concluded that Iprodione can result in hepatotoxicity in Zebrafish and may potentially represent toxicity risks for other aquatic fish, organisms, and even humans.

Shaw et al. [43] analyzed the effect of shikimic acid, a mycosporine-like amino acid, against stress induced by hexavalent chromium, a toxic heavy metal, in the hepatocytes of Zebrafish. The work of Guidony et al. [44] studies the toxicological effect of the bactericide Triclosan on the hepatocyte cells of Zebrafish.

### 3.9 FUTURE PERSPECTIVES

In light of all the above, Zebrafish is poised to emerge as an appropriate experimental model with many applications yet to be tested. Regarding toxicology, this animal could be incorporated into nanoscale approaches for the distribution and absorption of drugs, tested for cancer treatment using inhibitor medications, and there is also the possibility of generating transgenic models with fluorescent organs for tracking and drug screening [45].

Moreover, beyond the field of toxicology, this biomodel is expected to expand in applications in genetics, where it has already proven convenient for genome editing using CRISPR/Cas to study functional aspects of pathologies [46], as well as for generating transgenic animals and their application in in vivo studies [18]. In the realm of neurobehavioral studies, they can be assessed for cause and effect of neurotoxins generated by environmental pollutants [47], which is a relevant theme considering contemporary conditions. Additionally, studies on chronic stress, anxiety control, and other complex brain disorders [48, 49, 50] are also possible.

Thus, Zebrafish is expected to see its applications increasingly expanded across various and relevant fields of scientific knowledge, with technological advancements serving as significant contributors to the rapidity of this occurrence, leading to the development of ever more sophisticated and sensitive techniques.

## 4. CONCLUSION

This study presented a positive overview of the use of fish, with an emphasis on Zebrafish, as alternative models for scientific experimentation. *Danio rerio* has proven to be one of the most suitable samples when the objective of the research is to understand the toxicology of a compound, a new drug, and how humans may be affected. Thus, by demonstrating its suitability for the application of the fundamental principle of the 3Rs in scientific research, along with representing low cost and high efficiency, it is expected that this animal may emerge as a substitute experimental model of equal or even superior utility compared to laboratory rats.

## REFERENCES

1. Bachur TPR. Toxicologia: uma abordagem multidisciplinar – Volume I. São Paulo: Editora Ampolla; 2021.
2. Alwafi H, Khayat R, Banjabi A, Salawati E, Alotaibi BJ, Nassir R, Naser AY. A methodological review of drug-related toxicological studies in Saudi Arabia. *Cureus*. 2023;15(3).
3. Loregian B. Produção e colaboração científica brasileira em Toxicologia nos anos de 2006-2012 representada na Web of Science. 2014.
4. Oberdörster G, Oberdörster E, Oberdörster J. Nanotoxicology: an emerging discipline evolving from studies of ultrafine particles. *Environ Health Perspect*. 2005;113(7):823-39.
5. Rouquié D, Heneweer M, Botham J, Ketelslegers H, Markell L, Pfister T, Hennes C. Contribution of new technologies to characterization and prediction of adverse effects. *Crit Rev Toxicol*. 2015;45(2):172-83.
6. Ferreira LM, Hochman B, Barbosa MVJ. Modelo experimental em pesquisa. *Acta Cir Bras*. 2005;20:28-34.
7. Fagundes DJ, Taha MO. Modelo animal de doença: critérios de escolha e espécies de animais de uso corrente. *Acta Cir Bras*. 2004;19:59-65.
8. Freires IA, Sardi JDC, de Castro RD, Rosalen PL. Alternative animal and non-animal models for drug discovery and development: bonus or burden? *Pharm Res*. 2017;34:681-6.
9. Costa TLC D. História do Zebrafish na pesquisa científica: uma breve revisão. 2022.
10. Ota S, Kawahara A. Zebrafish: a model vertebrate suitable for the analysis of human genetic disorders. *Congenit Anom*. 2014;54(1):8-11.
11. Santana ÁLAD. Caracterização fenotípica do zebrafish (*Danio rerio*) portador de knockout no gene *prop1* gerado pela técnica de edição genômica CRISPR/Cas9 na fase adulta [tese de doutorado]. São Paulo: Universidade de São Paulo; 2023.
12. Engeszer RE, Patterson LB, Rao AA, Parichy DM. Zebrafish in the wild: a review of natural history and new notes from the field. *Zebrafish*. 2007;4(1):21-40.
13. van Wijk RC, Krekels EH, Kantae V, Harms AC, Hankemeier T, van der Graaf PH, Spaik HP. Impact of post-hatching maturation on the pharmacokinetics of paracetamol in zebrafish larvae. *Sci Rep*. 2019;9(1):2149.
14. Kütter MT, Barcellos LJG, Boyle RT, Marins LF, Silveira T. Boas práticas na criação e manutenção de zebrafish (*Danio rerio*) em laboratório no Brasil. *Ciênc Anim Bras*. 2023;24.
15. Howe K, Clark MD, Torroja CF, Torrance J, Berthelot C, Muffato M, et al. The zebrafish reference genome sequence and its relationship to the human genome. *Nature*. 2013;496(7446):498-503.

16. Fukushima H, Bailone RL, Baumgartner I, Borra RC, Correa T, de Aguiar L, et al. Potenciais usos do modelo animal Zebrafish *Danio rerio* em pesquisas na Medicina Veterinária. *Rev Educ Contin Med Vet Zoot.* 2020;18(1).
17. He JH, Gao JM, Huang CJ, Li CQ. Zebrafish models for assessing developmental and reproductive toxicity. *Neurotoxicol Teratol.* 2014;42:35-42.
18. Chen L, Xu M, Gong Z, Zonyane S, Xu S, Makunga NP. Comparative cardio and developmental toxicity induced by the popular medicinal extract of *Sutherlandia frutescens* (L.) R. Br. detected using a zebrafish Tuebingen embryo model. *BMC Complement Altern Med.* 2018;18:1-11.
19. Escaleira, R. (2017). O Zebrafish (*Danio rerio*). *Arquivos Brasileiros de Medicina Naval,* 78(1), 43-48.
20. Mwaffo V, Butail S, Porfiri M. In-silico experiments of zebrafish behaviour: modeling swimming in three dimensions. *Sci Rep.* 2017;7(1):39877.
21. Garcia GR, Noyes PD, Tanguay RL. Advancements in zebrafish applications for 21st century toxicology. *Pharmacol Ther.* 2016;161:11-21.
22. Pereira TCB, Campos MM, Bogo MR. Copper toxicology, oxidative stress and inflammation using zebrafish as experimental model. *J Appl Toxicol.* 2016;36(7):876-85.
23. Rivero-Wendt CLG, Matias R, do Nascimento GCZ, Heredia-Vieira SC, Santos MRM, Júnior OBP. Zebrafish: Recomendações de Manutenção em Laboratório. *Uniciências.* 2020;24(2):141-5.
24. Horzmann KA, Freeman JL. Making waves: New developments in toxicology with the zebrafish. *Toxicol Sci.* 2018;163(1):5-12.
25. Ribeiro OM, Félix L, Monteiro SM, Ribeiro C, Fontainhas-Fernandes A, Carrola JS. Embriões de peixe-zebra (*Danio rerio*). *Rev Ciênc Elementar.* 2024;12(1).
26. Simonetti RB. Zebrafish (*Danio rerio*): futuro do modelo animal em pesquisas biomédicas. 2014.
27. Zhao W, Chen Y, Hu N, Long D, Cao Y. The uses of zebrafish (*Danio rerio*) as an in vivo model for toxicological studies: A review based on bibliometrics. *Ecotoxicol Environ Saf.* 2024;272:116023.
28. Meyers JR. Zebrafish: development of a vertebrate model organism. *Curr Protoc Essent Lab Tech.* 2018;16(1)
29. Liu J, Yuan X, Fan C, Ma G. Application of the zebrafish model in human viral research. *Virus Res.* 2024;341:199327.
30. Kurach Ł, Chłopaś-Konowalek A, Budzyńska B, Zawadzki M, Szpot P, Boguszevska-Czubara A. Etazene induces developmental toxicity in vivo *Danio rerio* and in silico studies of new synthetic opioid derivative. *Sci Rep.* 2021;11(1):24269.

31. Andrade ALC D. Avaliação dos parâmetrotoxicológicos e comportamentais do Zebrafish exposto à Rotenonacomodelo animal da doença de Parkinson-Like e desenvolvimento de alternativananotecnológica para terapêutica. 2022.
32. Arman S, İşisağÜçüncü S. Cardiac toxicity of acrolein exposure in embryonic zebrafish (Danio rerio). *Environ Sci Pollut Res Int.* 2020;27(18):22423-33.
33. Li M, Yao L, Chen H, Ni X, Xu Y, Dong W, et al. Chiral toxicity of muscone to embryonic zebrafish heart. *AquatToxicol.* 2020;222:105451.
34. Lu X, Lu L, Gao L, Wang Y, Wang W. Calycosin attenuates doxorubicin-induced cardiotoxicity via autophagy regulation in zebrafish models. *Biomed Pharmacother.* 2021;137:111375.
35. Ton C, Lin Y, Willett C. Zebrafish as a model for developmental neurotoxicity testing. *Birth Defects Res A Clin Mol Teratol.* 2006;76(7):553-67.
36. Faria M, Prats E, Ramírez JRR, Bellot M, Bedrossiantz J, Pagano M, et al. Androgenic activation, impairment of the monoaminergic system and altered behavior in zebrafish larvae exposed to environmental concentrations of fenitrothion. *Sci Total Environ.* 2021;775:145671.
37. Azevedo, R. D., Falcão, K. V., Assis, C. R., Martins, R. M., Araújo, M. C., Yogui, G. T., ... & Bezerra, R. S. (2021). Effects of pyriproxyfen on zebrafish brain mitochondria and acetylcholinesterase. *Chemosphere*, 263, 128029.
38. Prabha N, Guru A, Harikrishnan R, Gatashah MK, Hatamleh AA, Juliet A, Arockiaraj J. Neuroprotective and antioxidant capability of RW20 peptide from histone acetyltransferases caused by oxidative stress-induced neurotoxicity in in vivo zebrafish larval model. *J King Saud Univ Sci.* 2022;34(3):101861.
39. Derikvandy A, Pourkhabbaz HR, Banaee M, Sureda A, Haghi N, Pourkhabbaz AR. Genotoxicity and oxidative damage in zebrafish (Danio rerio) after exposure to effluent from ethyl alcohol industry. *Chemosphere.* 2020;251:126609.
40. Chowdhury J, Mandal TK, Mondal S. Genotoxic impact of emerging contaminant amoxicillin residue on zebrafish (Danio rerio) embryos. *Heliyon.* 2020;6(11).
41. Qiu L, Jia K, Huang L, Liao X, Guo X, Lu H. Hepatotoxicity of tricyclazole in zebrafish (Danio rerio). *Chemosphere.* 2019;232:171-9.
42. Hu W, Chen G, Yuan W, Guo C, Liu F, Zhang S, Cao Z. Iprodione induces hepatotoxicity in zebrafish by mediating ROS generation and upregulating p53 signalling pathway. *Ecotoxicol Environ Saf.* 2024;270:115911.
43. Shaw P, Sen A, Mondal P, Bhowmik AD, Rath J, Chattopadhyay A. Shinorine ameliorates chromium induced toxicity in zebrafish hepatocytes through the facultative activation of Nrf2-Keap1-ARE pathway. *AquatToxicol.* 2020;228:105622.
44. Guidony NS, Scaini JLR, Oliveira MWB, Machado KS, Bastos C, Escarrone AL, Souza MM. ABC proteins activity and cytotoxicity in zebrafish hepatocytes exposed to triclosan. *Environ Pollut.* 2021;271:116368.

45. Lee KY, Jang GH, Byun CH, Jeun M, Searson PC, Lee KH. Zebrafish models for functional and toxicological screening of nanoscale drug delivery systems: promoting preclinical applications. *Biosci Rep.* 2017;37(3).
46. Sharma P, Sharma BS, Verma RJ. CRISPR-based genome editing of zebrafish. *Prog Mol Biol Transl Sci.* 2021;180:69-84.
47. Zhang B, Yang X, Zhao J, Xu T, Yin D. Studying neurobehavioral effects of environmental pollutants on zebrafish larvae. *J Vis Exp.* 2020;(156).
48. Demin KA, Taranov AS, Ilyin NP, Lakstygai AM, Volgin AD, de Abreu MS, et al. Understanding neurobehavioral effects of acute and chronic stress in zebrafish. *Stress.* 2021;24(1):1-18.
49. Chahardehi AM, Hosseini Y, Mahdavi SM, Naseh I. Zebrafish, a biological model for pharmaceutical research for the management of anxiety. *Mol Biol Rep.* 2023;50(4):3863-72.
50. Kalueff AV, Stewart AM, Gerlai R. Zebrafish as an emerging model for studying complex brain disorders. *Trends Pharmacol Sci.* 2014;35(2):63-75.