

# Review Article

## A Review on Conversion of Agricultural Waste to Bioenergy: Processes and Environmental Impacts

### ABSTRACT

Agricultural waste is a substantial and underutilized resource with the potential to contribute enormously to sustainable bioenergy production. The conversion technologies for these wastes is a significant step in the bioenergy process, therefore analyzing their impacts on the environment is considered valuable. A review of current literature, with an emphasis on studies that examines the relationship between agricultural wastes, different conversion technologies of wastes to bioenergy, and their environmental impacts was carried out. Databases like PubMed, Google scholar and Web of Science were searched using keywords like "Lignocellulosic biomass hydrolysis", "bioenergy", "Agricultural food wastes", and "environmental impacts". This review demonstrates how different conversion methods may alleviate climate change, enhance energy security while fostering sustainable development. We examined biochemical conversion methods such as anaerobic digestion and fermentation to show their advantages over the thermochemical approaches. It was found that these conversion processes are not only circular but also environmentally friendly with respect to biofuel production. Anaerobic digestion was identified as one of the most promising options because of its high energy yields and decentralized character. However, there are still issues in optimizing bioenergy conversion processes and conducting a comprehensive life cycle assessment that explains some of these challenges is required. This review highlights the current and emerging technologies for transformation of agricultural wastes into bioenergy and its environmental impacts, as it is necessary to transition into a bioenergy-driven economy to foster a healthy and robust future.

*Keywords: Bioenergy, Bioethanol, Agricultural wastes, Environmental impacts, Biomass hydrolysis*

### 1. INTRODUCTION

The global energy landscape is transforming due to urbanization which has led to increasing energy demands and environmental concerns [1,2]. Traditional energy sources, predominantly fossil fuels such as natural gas, coal and oil, have a limited nature and are contributing to the continuous emission of greenhouse gases into the atmosphere, which results in climate change and other environmental damages [2,3,4]. The urgency to mitigate these impacts has pushed global focus toward renewable energy sources which provide a cleaner and environmentally friendly option. Renewable energy sources including solar power, wind power, hydro-energy, and bioenergy, present opportunities to reduce carbon emissions and improve energy security [1,4,5].

Bioenergy produced from agricultural waste, is a prospective option among the various renewable energy sources available [6,7]. Agricultural wastes including residues from crop production, processing by-products, and other organic materials form a vast and often underutilized resource [7,8]. Transforming these wastes into bioenergy offers several advantages, such as decreasing the volume of waste released into landfills, curbing methane release from decomposition, and creating an energy source that supports

sustainability in the agricultural sector [9,10,11]. By tackling waste management issues, this method of converting agricultural wastes to bioenergy promotes energy variety and boosts rural economies.

Although agricultural waste holds promise as a bioenergy source, additional research is necessary to determine the best conversion methods for agricultural waste-to-energy and understand the environmental consequences of these processes. This review seeks to present a comprehensive summary of the processes involved in converting agricultural waste to bioenergy and to evaluate their associated environmental impacts.

## 2. OVERVIEW OF AGRICULTURAL WASTE, ITS GENERATION AND DISPOSAL

Various farming and processing activities generate agricultural wastes, which can be divided into organic and inorganic types [12]. Organic wastes originate from biological sources and are biodegradable [13]. Key types of organic agricultural wastes include residues from harvested crops, livestock wastes, and agro-industrial secondary products [14]. After harvesting, plants leave behind materials like wheat straw, barley straw, rice husks, and corn husks, which often amount to a considerable volume, constituting what is known as crop residues [15,16]. Livestock wastes include manure from cattle, pigs, and poultry, and bedding materials [17]. These are rich in organic matter and nutrients, making them a valuable resource if managed properly. The by-products of agro-industries, such as sugarcane bagasse (the fibrous material remaining after juice extraction), rice husks, and olive pomace (the residue from olive oil extraction), are generated during the processing of agricultural products and are accumulated in large quantities [18].

Traditionally, agricultural waste has been managed through techniques such as burning, landfilling, anaerobic digestion, composting or utilized in the feed of animals [13,14,19,20]. Burning which releases pollutants into the atmosphere and contributes to air pollution is still prevalent in many regions [21,22]. While landfilling reduces waste accumulation, it often leads to methane emissions, a potent greenhouse gas, and land degradation [23]. An environmentally friendly alternative is composting which involves the aerobic decomposition of organic waste to produce nutrient-filled compost which enhances the richness of the soil [13,24,25]. Anaerobic digestion involves decomposition of organic waste without the presence of oxygen, resulting in the production of biogas, which consists of a combination of methane and carbon dioxide, along with digestate, a by-product that is rich in nutrients [26,27]. Anaerobic digestion is mostly used for managing livestock manure and organic residues [28]. Certain agricultural by-products, like rice husks and some agro-industrial residues, are also repurposed as animal feed or feed supplements [29,30]. These conventional methods have several limitations, including environmental pollution, loss of valuable soil nutrients, and inefficient resource utilization [31]. Organic agricultural wastes are thus majorly used for bioenergy conversion, offering an eco-friendly and renewable energy source while addressing waste disposal challenges [7].

## 3. OVERVIEW OF BIOENERGY

Bioenergy is a form of renewable energy that is obtained from organic substances known as biomass, which includes plant and animal matter, agricultural by-products, forestry residues [32]. The World Economic Forum states that bioenergy serves as a sustainable energy alternative, reducing dependence on fossil fuels and alleviating the negative impacts of climate change [2,33,34]. Production-related manufacturing methods for bioenergy can be roughly divided into two groups. Biological and thermochemical technologies. Biomass is converted into energy via thermochemical procedures, which include liquefaction, combustion, gasification, and pyrolysis. Liquefaction is the process of turning biomass into liquefied products, using a synthesis of both physical and chemical processes.

Bioenergy can be produced through the oxidation of biomass substrates, which can be divided into three categories: first-generation (edible food sources), second-generation (non-edible sources), and third generation (biomass derived from algae). Algal substrates have advantages over other materials due to their quick growth, abundance, capacity to be grown in arable terrain, and low energy consumption. However, further research and development is still needed for their large-scale conversion procedures. Many different mechanisms, including biological, chemical, and thermal, can convert biomass into bioenergy. The International Energy Agency (IEA) reported that approximately 55% of the world's

**Commented [H1]:** In this section some technique mentioned but only defined Liquefaction, you may add about others.

renewable energy comes from bioenergy. It also contributes significantly to the overall energy supply, representing over 6% of the global energy consumption [35].

### **3.1. Types of Biofuels**

#### **3.1.1. Bioethanol**

Bioethanol, a widely utilized biofuel, is a type of alcohol produced by fermenting plant-based sugars. While traditionally made from food crops like corn or sugar cane (first-generation bioethanol), technological advancements have enabled the use of woody plants and grasses (second-generation bioethanol). Algae are being used as a potential source to produce bioethanol (third-generation bioethanol) [36,37]. Bioethanol has around 34% less energy per unit volume than gasoline.

#### **3.1.2. Biogas**

Biogas is a renewable gas produced from organic wastes like agricultural byproducts, manure, food scraps, and sewage [11]. It is created through anaerobic digestion, where methanogens or anaerobic microorganisms degrade organic matter in the absence of oxygen in a bioreactor, biodigester, or anaerobic digester [26,38,39]. The resulting gas from this procedure is primarily a mixture of carbon dioxide and methane, with other gases like moisture, siloxanes, and hydrogen sulfide (H<sub>2</sub>S) in trace amount [40,41]. Biogas can be incinerated directly for heat or used in engines to generate electricity. Its production efficiency and the composition are affected by various factors, including the type of waste utilized and the prevailing conditions (temperature, pH, and substrate concentration) within the digestion process [11,42,43].

#### **3.1.3. Biodiesel**

Biodiesel, also known as Fatty Acid Methyl Ester (FAME), is a renewable diesel fuel from biological materials such as vegetable oils, animal fats, or recycled greases. It is a clear to dark brown liquid with a high flash point compared to regular diesel. Compared to traditional diesel, biodiesel is slightly denser and has a lower energy content [44]. Despite these differences, biodiesel provides numerous environmental advantages such as a decrease in greenhouse gas emissions and enhanced air quality when compared to conventional diesel fuel. It is biodegradable and can be mixed with petroleum diesel to produce a less polluting fuel [45,46].

### **3.2 Advantages of Bioenergy**

Bioenergy's renewable nature makes it a valuable resource. It is made from organic components that may be renewed in a short amount of time, such as plant and animal waste. Sustainable forestry and agriculture methods provide bioenergy supplies, which are easily accessible in contrast to fossil fuels that require millions of years to develop. Bioenergy can be used as a long-term, renewable energy source because of its constant availability, strengthening the foundation of an increasingly resilient and sustainable energy system [47].

The possibility for bioenergy to be carbon neutral is another significant benefit. When biomass is burned to produce energy, the carbon dioxide (CO<sub>2</sub>) emitted during combustion is offset by the CO<sub>2</sub> taken up by the plants as they grow, thereby establishing a closed carbon cycle. This lessens the effect of climate change by preventing input of new CO<sub>2</sub> to the atmosphere [48]. Bioenergy is an environmentally benign energy source since it may drastically reduce greenhouse gas emissions when utilized as a substitute for fossil fuels within well-managed systems.

By diversifying energy sources and lowering dependency on fossil fuels, bioenergy also improves energy security. Its local production can be achieved, thereby decreasing the reliance on energy imports and protecting economies from changes in the price of energy globally. Furthermore, the production of bioenergy boosts regional economies by generating employment in the forestry, agriculture, and bioenergy sectors [49]. This promotes stability and economic progress in the neighborhood while also bolstering energy security [50].

## 4. PROCESSES FOR CONVERSION OF AGRICULTURAL WASTE TO BIOENERGY

### 4.1. Biochemical Processes

#### 4.1.1. Anaerobic Digestion

The series of procedures known as anaerobic digestion is how microbes decompose biodegradable materials when there is no oxygen present. There are four main phases of anaerobic digestion which include acidogenesis, acetogenesis, methanogenesis, and hydrolysis [51,52]. The overall procedure can be explained by the chemical reaction in figure 1, wherein anaerobic microorganisms biochemically decompose organic material such as glucose resulting in the production of carbon dioxide (CO<sub>2</sub>) and methane (CH<sub>4</sub>).




Figure 1: The anaerobic digestion process simplified

Biodegradable waste products including crop residues, grass clippings, food wastes, wastewater, and animal waste can all be used as feedstocks [53]. Woody wastes are an exception, as they are mostly untouched by digestion since most anaerobic microorganisms are unable to break down lignin [54,55]. Lignin can be decomposed by xylophagous anaerobes, or through high-temperature pretreatments such as pyrolysis. [54,55] For the purpose of producing biogas, anaerobic digesters can be supplied with energy crops that are grown specifically for them, such as silage.

Anaerobic digesters (methanogen species) are identified by their two standard operating temperature levels [56]. The optimal temperature range for mesophilic digestion lies between 30 and 38°C, with ambient temperatures ranging from 20 to 45 °C, where mesophilic bacteria are the dominant microorganisms. In contrast, thermophilic digestion involving thermophiles as the key microorganisms occur best at temperatures between 49 and 57°C or higher, up to 70°C [56,57]. The temperature at which a digester operates influences the types of bacteria present and how quickly waste breaks down. Higher temperatures can speed up the process but may need extra energy to maintain [57].

The retention time, which is the amount of time feedstock remains in the digester, is a crucial factor in digestion efficiency [58]. It depends on factors such as feedstock composition, digester design, and operating conditions [58,59]. Retention times for two-stage mesophilic digestions typically range from 15 to 40 days [60], whereas retention lengths for single-stage thermophilic digestions are often faster, taking about 14 days [61]. Because some of these systems are plug-flow, it is possible that the material has not fully degraded in this amount of time, leading to incomplete digestion. In this case, the digestate that leaves the system will smell stronger and have a darker color [62].

#### 4.1.2. Fermentation

Microbial fermentation of sugar yields bioethanol [63]. Starch and cellulose, two major components of plant matter, are both composed of sugars. While theoretically both can be fermented into sugars, currently only those containing significant amounts of sugar (like sugarcane) or starch (like corn) are economically feasible for biofuel production [63].

Ethanol and carbon dioxide are produced during the fermentation of glucose and other sugars found in agricultural wastes [64]. The fermentation process for ethanol is anaerobic when it involves yeast, but it can also be aerobic when it involves certain other microbes and enzymes [63,64,65]. Ethanol is a fuel that burns efficiently and produces fewer pollutants compared to other types of fuels [66]. It is a renewable energy source as it is derived from plant materials.




Figure 2: Overview of the fermentation process

While starch-based feedstocks are more common, ethanol can also be produced from cellulosic materials like wood, grasses, and agricultural residues [67]. However, cellulose must first be pretreated to break down its complex structure and release the sugars it contains [68,69]. This pretreatment step often involves the use of enzymes or chemicals [69,70]

## 4.2. Thermochemical Processes

### 4.2.1. Combustion

Biomass is converted into heat via biomass heating systems [71]. The systems can generate heat through gasification, anaerobic digestion, combined heat and power (CHP), direct combustion, or aerobic digestion [72,73]. Biomass heating systems can be either fully or partially automated, or they may function as combined heat and power systems [73].

As biomass uses waste from urban, rural, and industrial areas as well as agricultural wastes to create heat with less of an effect on the environment compared to fossil fuels, it is advantageous to use biomass in heating systems [8,74,75]. In the past, biomass utilized as wood fuel supplied most of the human warmth until fossil fuels were used in large numbers [8,76]. Also, the carbon contained in biomass is an integrated natural carbon cycle, in contrast to the carbon found in fossil fuels which is not part of this cycle [76,77]. The combustion of fossil fuels results in a permanent release of carbon into the atmosphere, thereby indicating that this method of energy production has a restricted long-term environmental impact or carbon footprint [77].

### 4.2.2. Pyrolysis

The process of thermally breaking down substances at high temperatures, frequently in an inert environment without oxygen availability is known as pyrolysis [78]. In most cases, pyrolysis involves heating the material above the point at which it breaks down chemically to release the bonds between its components [79,80]. Usually, the pieces unite to form residues with a greater molecular mass, even amorphous covalent solids, but they can also become smaller molecules [78]. In many circumstances, there may be some oxygen, water, or other substances present, allowing for the possibility of hydrolysis, combustion, or other chemical processes in addition to the actual pyrolysis. In other cases, such as when burning firewood, making charcoal the old-fashioned way, or steam cracking crude oil, those chemicals are introduced on purpose.

Many common organic compounds decompose between 100 and 500 °C [81]. Sugars degrade at temperatures between 160 and 180°C, while cellulose, found in wood, paper, and cotton, breaks down at approximately 350°C [82,83]. Another important component of wood, lignin, begins to decompose at 350 °C and can continue to release volatile compounds until 500 °C [84,85]. Water, carbon monoxide (CO), carbon dioxide (CO<sub>2</sub>) and a variety of organic compounds are typically among the breakdown products [85].

### 4.2.3. Gasification

The process of gasification involves the transformation of carbon-rich materials sourced from biomass or fossil fuels into gaseous products, consisting of nitrogen (N<sub>2</sub>), carbon monoxide (CO), hydrogen (H<sub>2</sub>), and carbon dioxide (CO<sub>2</sub>) [86,87]. To accomplish this, the feedstock material undergoes a reaction at high temperatures typically exceeding 700 °C without burning, by regulating the level of oxygen and/or steam present in the reaction [88,89,90]. As the gas is mostly made of hydrogen (H<sub>2</sub>) and carbon monoxide (CO)

which are flammable, the resulting gas mixture is known as syngas or producer gas, and it can be used as fuel on its own [91]. The process of gasification, which produces power when the resulting gas is burned, is referred to as a renewable energy source provided that the feedstock utilized to create the gasified compounds was biomass [86,87,90].

### **4.3. Emerging and Innovative Technologies**

#### **4.3.1. Hydrothermal Liquefaction**

A possible technique for turning wet biomass into bio-crude that resembles the natural production of fossil fuels but happens considerably more quickly is hydrothermal liquefaction (HTL). HTL operates under subcritical or supercritical conditions in a water-rich environment, in contrast to traditional thermal processes like pyrolysis. The procedure usually entails heating wet biomass at pressures between 4 and 22 MPa and temperatures between 250 and 374 degrees Celsius while it is wet. These circumstances make it easier for the biomass's complex organic molecules to break down into smaller, more energy-dense compounds, which is how bio-crude is formed [92].

The capacity of HTL to process high-moisture feedstocks without the need for previous drying, which is energy-expensive, is one of its main advantages. Because of this, a variety of wet biomasses, including algae, sewage sludge, and agricultural leftovers, are especially well-suited for conversion via HTL [93]. High-pressure water serves as a medium for several reactions, including hydrolysis and depolymerization, which generate bio-crude, gases, solid and aqueous byproducts. With typical refining procedures, the bio-crude produced through hydrothermal liquefaction can be enhanced to yield transportation fuels including gasoline, diesel, and jet fuel. It has a similar composition to that of petroleum crude oil [94]. In addition, the gaseous phase—which is predominantly made up of carbon dioxide, methane, and hydrogen—may be used for energy recovery, while the aqueous phase contains important nutrients that can be recycled [92,93].

For biomass valorization, HTL provides a sustainable path with a negligible environmental impact. Its ability to convert a variety of wet biomass feedstocks into high-energy-density bio-crude. This makes theis technology desirable for the production of sustainable energy, contributing to a decrease in dependence on fossil fuels and the emission of greenhouse gases [94].

#### **4.3.2. Algal Bioenergy Production**

Using agricultural wastewater to grow algae offers a sustainable way to produce biofuel while resolving environmental issues related to wastewater treatment. Nutrients like phosphorus and nitrogen, which are necessary for algae growth, are abundant in agricultural effluent. By using this wastewater, extra nutrients that would otherwise contribute to eutrophication and water pollution in natural water bodies are lessened for the environment, while also offering a cost-effective supply of nutrients [95].

Through photosynthesis, algae can effectively absorb these nutrients from the wastewater and transform them into biomass. Following processing, this biomass has the potential to be transformed into various types of biofuels, such as biodiesel, bioethanol, and biogas [96]. Algal farming combined with agricultural wastewater treatment can improve water quality before release or irrigation by means of bioremediation. This technique still needs to be optimized for large-scale production. Important research topics include the algal strains selection with high fatty content, the variety in wastewater composition, and economical algal biomass collection and processing [97]. However, the combined advantages of producing renewable energy and treating wastewater make this a desirable path for the development of sustainable biofuels [96].

## **5. ENVIRONMENTAL IMPACTS OF AGRICULTURAL WASTE-TO-BIOENERGY CONVERSION**

### **5.1 Positive Environmental Impacts**

### **5.1.1 Reduction of Greenhouse Gas Emissions**

One of the most significant advantages of converting agricultural waste into bioenergy is the reduction of greenhouse gas emissions. When these wastes are burned outdoors or allowed to break down naturally, they can emit gases into the environment, including methane (CH<sub>4</sub>) and carbon dioxide (CO<sub>2</sub>) [98,99]. Both are strong greenhouse gases and have a major role in climate change and global warming.

Technologies like anaerobic digestion, gasification, or combustion, which provide a controlled environment to harness energy while limiting harmful emissions, are frequently used in the production of bioenergy, notably biofuels generated from agricultural waste. For example, methane emissions are captured during anaerobic digestion, and this degrades organic waste in the absence of oxygen and generates biogas that can be used for heating, electricity or transportation fuel [100].

By using this procedure, [101,102] reported that methane which would have otherwise leaked into the atmosphere via landfills or open burning of garbage is prevented. Moreover, because bioenergy is regarded as carbon-neutral, a closed carbon cycle is established when the CO<sub>2</sub> emissions from burning it are offset by the CO<sub>2</sub> that plants absorb through their growth. Additionally, the use of bioenergy lessens reliance on fossil fuels, which account for around 75% of greenhouse gas emissions worldwide [99,101,102]. In comparison to traditional fossil fuels, bioenergy can cut greenhouse gas emissions by up to 90%. Thus, turning agricultural waste into bioenergy is essential to reducing global warming and helping nations fulfill their obligations to cut their carbon emissions under frameworks like the Paris Agreement [98].

### **5.1.2 Mitigation of Agricultural Waste Disposal Problems**

Agricultural activities produce vast amounts of waste, including crop residues (e.g., straw, husks) and animal manure, estimated globally at 5 billion tons annually [103]. If not managed properly, these wastes can lead to significant environmental problems, such as contamination of air and water resources. Open burning of residues releases harmful pollutants such as particulate matter, carbon monoxide, and volatile organic compounds, which exacerbate air pollution and can lead to respiratory issues. Additionally, landfilling organic waste leads to the formation of leachate, which contaminates groundwater [104].

Converting agricultural waste into bioenergy addresses these challenges by using waste as feedstock for energy production, thereby reducing pollution. [105] reported that bioenergy reduces waste volume by up to 90%, minimizing its environmental footprint. Decentralized bioenergy systems, especially in rural areas, provide a sustainable alternative to traditional waste disposal methods. By setting up local facilities, agricultural residues can be continuously recycled into energy, reducing the need for harmful disposal practices and creating a circular, sustainable waste management system [103,104].

### **5.1.3 Improvement of Soil Health Through Biochar Application**

Biochar is a byproduct of pyrolysis used to produce bioenergy and has several advantages for the sustainability of the environment and soil health. It is created by thermal decomposition of organic material in an oxygen-free environment, improving the composition of soil structure, increasing its ability to retain nutrients, and water-holding capacity. In the presence of degraded soils, biochar has been reported to increase crop yields by up to 30% [106], facilitating plant development by its porous nature, which traps moisture and nutrients.

Additionally, biochar raises soil organic carbon, bringing fertility back to areas that have lost nutrients and been degraded. According to [107], biochar may trap carbon for hundreds to thousands of years, capturing 1-3 gigatons of CO<sub>2</sub> yearly and assisting in the mitigation of climate change. Furthermore, biochar limits pollution from agricultural runoff and promotes sustainable farming methods by lowering nutrient leaching into groundwater.

**Commented [H2]:** You may cite this paper at here  
10.1080/03067319.2021.1974013

## 5.2 Adverse Environmental Impacts

### 5.2.1 Air pollution from combustion processes.

Air pollution from the combustion of biomass is one of the major environmental concerns connected to the conversion of agricultural waste into bioenergy. In long-term carbon cycles, bioenergy is thought to be carbon-neutral; nevertheless, burning biomass can emit a considerable quantities of air pollutants, including sulfur dioxide (SO<sub>2</sub>), particulate matter (PM), nitrogen oxides (NO<sub>x</sub>), and volatile organic compounds (VOCs) [108,109]. The neighboring populations' respiratory health problems, acid rain, and poor air quality are all caused by these pollutants.

[110] reported that depending on the feedstock and combustion method employed, burning biomass can release particulate matter up to amounts that are on par with or even greater than those from burning conventional fossil fuels. Specifically, it is recognized that agricultural waste products like straw and husks can generate significant levels of fine particulate matter (PM<sub>2.5</sub>) when burned inefficiently, posing considerable dangers to both human health and the environment. These emissions have the potential to worsen the regional air quality in rural areas, as bioenergy plants are frequently situated close to agricultural sources. Moreover, inefficient combustion of agricultural waste can result in the potent greenhouse gasses carbon monoxide (CO) and methane (CH<sub>3</sub>) being released. Even with the development of more effective combustion systems and emissions control mechanisms, biomass energy plants continue to be a source of air pollution despite technological breakthroughs [109,111]. As an illustration of the continued difficulties with air quality, a 2022 assessment found that even the most advanced biomass power facilities in Europe released 10–20% more PM<sub>2.5</sub> than natural gas power plants [109,110].

### 5.2.2 Potential water contamination from leachate and runoff

Leachate, a liquid byproduct of organic waste breakdown during anaerobic digestion or storage, can be produced during the production of bioenergy. When leachate is not controlled, it can seep into the soil and contaminate both surface and groundwater [112]. It can also carry organic contaminants, heavy metals, and minerals like phosphorus and nitrogen. When agricultural waste is converted to bioenergy, one major environmental risk is water contamination. Aquatic habitats may be harmed by eutrophication, which is caused by algae blooms that reduce oxygen levels. [113] reported that nutrient-rich leachate from bioenergy plants is a major cause of North American algal blooms.

Furthermore, the growth of bioenergy feedstocks, mainly switch grass and maize, can produce runoff that contaminates surrounding waterways with fertilizers and pesticides. Water contamination from bioenergy is still a serious problem despite regulatory measures, particularly in locations where the sector is expanding rapidly [104],[114].

### 5.2.3 Land use and biodiversity considerations

The growth in the generation of bioenergy from agricultural waste has significant implications on biodiversity and land use. While using agricultural wastes reduces the strain on resources, increased demand for bioenergy feedstocks, such as energy crops, frequently results in habitat degradation and land change [115]. Large-scale bioenergy projects need specialized land, which means they must replace natural ecosystems and agricultural areas that are often utilized to produce food. The International Union for Conservation of Nature (IUCN) reported in 2022 that the biodiversity in Europe is declining as a result of wetlands and grasslands being converted into monoculture energy crop plantations, which only marginally support wildlife [116].

Additionally, there is growing rivalry for land between the production of food and bioenergy. Growing concerns regarding the loss of arable land for food arise as the demand for bioenergy rises, especially in areas where there is food insecurity. According to [117], the World Resources Institute revealed that 12%

**Commented [H3]:** May use reference  
<https://www.tandfonline.com/doi/abs/10.1080/03067319.2022.2100256>

of global arable land could be diverted for bioenergy feedstocks by 2030, exacerbating food shortages and raising food prices.

Greenhouse gas emissions can also be attributed to land-use changes related to bioenergy. The benefits of bioenergy for the environment are offset by deforestation and the deterioration of ecosystems rich in carbon, which releases stored carbon. While crop rotation and agroforestry can lessen some negative effects, continuous bioenergy expansion runs the danger of causing more environmental degradation if not accompanied by effective, sustainable land-use planning and initiatives for biodiversity protection [115,118].

### 5.3 Impacts Comparison of Conversion Processes for Bioenergy Production

Thermochemical and biochemical approaches for turning agricultural waste into bioenergy influence the environment that is different from the other. Biochemical conversion methods, such as fermentation and anaerobic digestion, strongly rely on microbial action to degrade organic material. For instance, agricultural wastes are converted into bioenergy in Germany, one of the leading countries in the generation of biogas with over 9,000 plants. This biochemical method lowers emissions of nitrogen oxides (NOx) and particulate matter (PM) in comparison to fossil fuels [119]. However, methane leakage remains a critical concern. According to a study by [120], biogas facilities may lose 3-5% of their methane, negating the climatic advantages they might otherwise provide. Additionally, anaerobic digestion generates nutrient-rich effluents that need a lot of water, which might pollute water in places like rural China where small-scale biogas facilities don't have access to adequate wastewater treatment [119,120].

Conversely, thermochemical processes use high temperatures to decompose biomass through chemical reactions. For example, in the United States, gasification is frequently used to turn agricultural residues like corn stover into syngas for electricity. A 2022 Midwest case study found that while gasification emits fewer greenhouse gases (GHGs) than coal, it produces more sulfur dioxide (SO<sub>2</sub>) and particulate matter (PM<sub>2.5</sub>). In Brazil, large-scale combustion of sugarcane bagasse for energy is efficient, but it produces high emissions of PM and NOx, which adversely affect air quality [121,122].

In Kenya, agricultural waste is converted by pyrolysis into biochar and bio-oil. The carbon sequestered by the biochar improves soil fertility [123]. However, pyrolysis has faced criticism for potentially harming the environment due to emission of carbon monoxide (CO) and volatile organic compounds (VOCs). Thermochemical processes are more effective and adaptable than biological ones, although producing greater pollutants. Though they produce beneficial byproducts like organic fertilizers and emit less emissions, biochemical processes like anaerobic digestion are ultimately environmentally favorable [124]. Strict water management regulations and enhanced methane capture technology are needed to propel anaerobic digestion approach forward. Scaling up such innovations, particularly in regions focused on air quality and sustainability, can help expand the adoption of anaerobic digestion as a cleaner energy solution worldwide [124,125].

## 6. CONCLUSION

Transformation of agricultural waste into bioenergy offers a promising solution for tackling pressing global issues such as climate change, waste management, and energy security. This review has explored agricultural waste sources, the diverse technologies utilized for converting waste into energy, and their associated environmental impacts. While significant progress has been made, there is still considerable potential to optimize bioenergy production processes, enhance their economic viability, and minimize negative externalities.

## CONSENT

Not applicable

**Commented [H4]:** Expand the conclusion section by adding some important points.

## ETHICAL APPROVAL

Not applicable

## REFERENCES

1. Chen X.H., Tee K., Marwa Elnahass M., and Ahmed R. Assessing The Environmental Impacts of Renewable Energy Sources: A Case Study on Air Pollution and Carbon Emissions in China. *Journal of Environmental Management*. 2023, Volume 345. 118525, ISSN 0301-4797. <https://doi.org/10.1016/j.jenvman.2023.118525>.
2. Yongjun Lv. Transitioning to Sustainable Energy: Opportunities, Challenges, and the Potential of Blockchain Technology. *Front. Energy Res.*, 2023, Volume 11. <https://doi.org/10.3389/fenrg.2023.1258044>
3. Foster S. and Elzinga D. The Role of Fossil Fuels in a Sustainable Energy System. *Sustainable Energy*. 2015, No. 3 Vol. I.I. Accessed 14 August 2024. Available: <https://www.un.org/en/chronicle/article/role-fossil-fuels-sustainable-energy-system>
4. Kumar J.C.R., And Majid, M.A. Renewable Energy for Sustainable Development in India: Current Status, Future Prospects, Challenges, Employment, and Investment Opportunities. *Energ Sustain Soc*. 2020, 10 (2). <https://doi.org/10.1186/s13705-019-0232-1>
5. Chen S., Zhang C., and Lu X. Energy Conversion from Fossil Fuel to Renewable Energy. In: Akimoto, H., Tanimoto, H. (eds) *Handbook of Air Quality and Climate Change*. 2023. Springer, Singapore. [https://doi.org/10.1007/978-981-15-2760-9\\_42](https://doi.org/10.1007/978-981-15-2760-9_42)
6. Muhammad S. Possibility of Utilizing Agriculture Biomass as a Renewable and Sustainable Future Energy Source, *Heliyon*. 2022, Volume 8, Issue 2. E08905, ISSN 2405-8440. <https://doi.org/10.1016/j.heliyon.2022.e08905>
7. Kumar J.A., Sathish S., Prabu D., Renita A.A., Saravanan A., Deivayanai V.C., et al. Agricultural Waste Biomass for Sustainable Bioenergy Production: Feedstock, Characterization and Pre-Treatment Methodologies. *Chemosphere*. 2023, Volume 331, 138680, ISSN 0045-6535. <https://doi.org/10.1016/j.chemosphere.2023.138680>
8. Kalak T. Potential Use of Industrial Biomass Waste as a Sustainable Energy Source in the Future. *Energies*. 2023; 16(4):1783. <https://doi.org/10.3390/en16041783>
9. Xu M., Yang M., Sun H., Gao M., Wang Q., and Wu C. Bioconversion of Biowaste into Renewable Energy and Resources: A Sustainable Strategy. *Environmental Research*. 2022, Volume 214, Part 2, 113929, ISSN 0013-9351. <https://doi.org/10.1016/j.envres.2022.113929>
10. Un C. A Sustainable Approach to the Conversion of Waste into Energy: Landfill Gas-to-Fuel Technology. *Sustainability*. 2023; 15(20):14782. <https://doi.org/10.3390/su152014782>
11. Kabeyi M.J.B., and Olanrewaju O.A. Biogas Production and Applications in the Sustainable Energy Transition. *Journal of Energy*. 2022 <https://doi.org/10.1155/2022/8750221>
12. Obi F.O., Ugwuishiwu B.O., and Nwakaire J.N. Agricultural Waste Concept, Generation, Utilization And Management. *Nigerian Journal of Technology (NIJOTECH)*, 2016, Vol. 35, No. 4, pp. 957 – 964. <http://dx.doi.org/10.4314/njt.v35i4.34>
13. Sayara T., Basheer-Salimia R., Hawamde F., and Sánchez A.. Recycling of Organic Wastes through Composting: Process Performance and Compost Application in Agriculture. *Agronomy*. 2020; 10(11):1838. <https://doi.org/10.3390/agronomy10111838>
14. Ogbu C.C. and Okey N.S. Agro-Industrial Waste Management: The Circular and Bioeconomic Perspective. *Agricultural Waste - New Insights*. IntechOpen; 2023. Accessed 14 August 2024. Available: <http://dx.doi.org/10.5772/intechopen.109181>
15. Sarkar S., Skalicky M., Hossain A., Brestic M., Saha S., Garai S., et al. Management of Crop Residues for Improving Input Use Efficiency and Agricultural Sustainability. *Sustainability*. 2020; 12(23):9808. <https://doi.org/10.3390/su12239808>
16. Awogbemi O., Vandi Von Kallon D. Application of Biochar Derived from Crops Residues for Biofuel Production. *Fuel Communications*. 2023, Volume 15, .100088, ISSN 2666-0520. <https://doi.org/10.1016/j.jfueco.2023.100088>
17. Parihar S.S., Saini K.P.S., Lakhani G.P., Jain A., Roy B., Ghosh S. and Aharwal B. Livestock Waste Management: A Review. *Journal of Entomology and Zoology Studies*. 2019; 7(3): 384-393. Accessed

- 14 August 2024. Available: <https://www.entomoljournal.com/archives/2019/vol7issue3/PartG/6-6-95-692.pdf>
18. Šelo G., Planinić M., Tišma M., Tomas S., Komlenić D.K., and Bucić-Kojić A. A Comprehensive Review on Valorization of Agro-Food Industrial Residues by Solid-State Fermentation. *Foods*. 2021; 10(5):927. <https://doi.org/10.3390/foods10050927>
  19. Koul B., Yakoob M., and Shah M.P. Agricultural Waste Management Strategies For Environmental Sustainability. *Environmental Research*. 2022, Volume 206. 112285, ISSN 0013-9351, <https://doi.org/10.1016/j.envres.2021.112285>
  20. Kolawole I.D., Kolawole, G.O., Sanni-manuel, B.A. Kolawole S.K., Ewansiha J.U., Kolawole V.A., et al. Economic Impact Of Waste From Food, Water, And Agriculture In Nigeria: Challenges, Implications, And Applications—A Review. *Discov Environ* 2024, 2 (51). <https://doi.org/10.1007/s44274-024-00086-6>
  21. Usmani M., Kondal A., Wang J., and Jutla A. Environmental Association of Burning Agricultural Biomass in the Indus River Basin. *Geohealth*. 2020 Nov 1; 4(11):e2020GH000281. Doi: 10.1029/2020GH000281. PMID: 33163827; PMCID: PMC7597142.
  22. Lan R., Eastham S.D., Liu, T. et al. Air Quality Impacts Of Crop Residue Burning In India And Mitigation Alternatives. *Nat Commun*. 2022, 13, 6537. <https://doi.org/10.1038/s41467-022-34093-z>
  23. Abubakar I.R., Maniruzzaman K.M., Dano U.L., AlShihri F.S., AlShammari M.S., Ahmed S.M.S., et al. Environmental Sustainability Impacts of Solid Waste Management Practices in the Global South. *Int J Environ Res Public Health*. 2022 Oct 5;19(19):12717. DOI: 10.3390/ijerph191912717. PMID: 36232017; PMCID: PMC9566108.
  24. Ho T.T.K., Tra V.T., Le T.H., Nguyen N., Tran C., Nguyen P., et al. Compost To Improve Sustainable Soil Cultivation And Crop Productivity. Case Studies in Chemical and Environmental Engineering. 2022, Volume 6, 100211, ISSN 2666-0164. <https://doi.org/10.1016/j.cscee.2022.100211>
  25. Ayilara M.S., Olanrewaju O.S., Babalola O.O., and Odeyemi O. Waste Management through Composting: Challenges and Potentials. *Sustainability*. 2020; 12(11):4456. <https://doi.org/10.3390/su12114456>
  26. Aworanti O.A., Agbede O.O., Agarry S.E., Ajani A.O., Ogunkunle O., Laseinde O.T., et al. Decoding Anaerobic Digestion: A Holistic Analysis of Biomass Waste Technology, Process Kinetics, and Operational Variables. *Energies*. 2023; 16(8):3378. <https://doi.org/10.3390/en16083378>
  27. Haririchi S., Wainaina S., Sar T., Nojumi S.A., Parchami M., Parchami M., et al. Microbiological Insights into Anaerobic Digestion for Biogas, Hydrogen or Volatile Fatty Acids (VFAs): A Review. *Bioengineered*. 2022 Mar;13(3):6521-6557. doi: 10.1080/21655979.2022.2035986. PMID: 35212604; PMCID: PMC8973982.
  28. Kadam R., Jo S., Lee J., Khanthong K., Jang H., and Park J. A Review on the Anaerobic Co-Digestion of Livestock Manures in the Context of Sustainable Waste Management. *Energies*. 2024; 17(3):546. <https://doi.org/10.3390/en17030546>
  29. Yafetto L., Odamtten G.T., and Wiafe-Kwagyan M. Valorization Of Agro-Industrial Wastes Into Animal Feed Through Microbial Fermentation: A Review Of The Global And Ghanaian Case. *Heliyon*. 2023 Mar 22;9(4):e14814. Doi: 10.1016/j.heliyon.2023.e14814. PMID: 37025888; PMCID: PMC10070663.
  30. Vastolo A., Calabrò S., and Cutrignelli M.I. A review on the Use of Agro-industrial CO-products in Animals' Diets. *Italian Journal of Animal Science*. 2022, 21(1), pp. 577–594. DOI: 10.1080/1828051X.2022.2039562.
  31. Hajam Y.A., Kumar R., and Kumar A. Environmental Waste Management Strategies And Vermi Transformation For Sustainable Development. *Environmental Challenges*. 2023, Volume 13, 100747, ISSN 2667-0100. <https://doi.org/10.1016/j.envc.2023.100747>
  32. U.S Energy Information Administration. Biomass Explained. Last updated: July 30, 2024. Accessed 14 August 2024. Available: <https://www.eia.gov/energyexplained/biomass/>
  33. Philip Siu. How Turning Biomass Waste Into Sustainable Fuels Can Help Restore The Carbon Balance. *Energy Transition*. 2023. Accessed 14 August 2024. Available: <https://www.weforum.org/agenda/2023/12/biomass-waste-sustainable-fuels-carbon-climate-change/>
  34. Rial R.C. Biofuels Versus Climate Change: Exploring Potentials And Challenges In The Energy Transition. *Renewable and Sustainable Energy Reviews*. 2024, Volume 196, 114369, ISSN 1364-0321. <https://doi.org/10.1016/j.rser.2024.114369>

35. The International Energy Agency (IEA). Assessing Critical Energy Technologies For Global Clean Energy Transitions. Tracking Clean Energy Progress Report 2023. Accessed 14 August 2024. Available: <https://www.iea.org/reports/tracking-clean-energy-progress-2023>
36. Cavalius P., Engelhart-Straub S., Mehlmer N., Lercher J., Awad D., and Brück T. The Potential Of Biofuels From First To Fourth Generation. PLoS Biol. 2023 Mar 30;21(3):e3002063. Doi: 10.1371/journal.pbio.3002063. PMID: 36996247; PMCID: PMC10063169.
37. Tse T.J., Wiens D.J., and Reaney M.J.T. Production of Bioethanol—A Review of Factors Affecting Ethanol Yield. Fermentation. 2021; 7(4):268. <https://doi.org/10.3390/fermentation7040268>
38. Ngan, N.V.C., et al. Anaerobic Digestion of Rice Straw for Biogas Production. In: Gummert, M., Hung, N., Chivenge, P., Douthwaite, B. (eds) Sustainable Rice Straw Management. 2020. Springer, Cham. [https://doi.org/10.1007/978-3-030-32373-8\\_5](https://doi.org/10.1007/978-3-030-32373-8_5)
39. Kumar D.J.P., Mishra R.K., Chinnam S., Binnal P., and Dwivedi N. A Comprehensive Study On Anaerobic Digestion Of Organic Solid Waste: A Review On Configurations, Operating Parameters, Techno-Economic Analysis And Current Trends. Biotechnology Notes. 2024, Volume 5, Pages 33-49, ISSN 2665-9069. <https://doi.org/10.1016/j.biotno.2024.02.001>
40. Werkneh A.A. Biogas Impurities: Environmental And Health Implications, Removal Technologies And Future Perspectives. Heliyon. 2022 Oct 6;8(10):e10929. Doi: 10.1016/j.heliyon.2022.e10929. PMID: 36299513; PMCID: PMC9589174.
41. Jameel M.K., Mustafa M.A., Ahmed H.S., Mohammed A.J., Ghazy H., Shakir M.N., et al. Biogas: Production, Properties, Applications, Economic And Challenges: A Review. Results In Chemistry. 2024, Volume 7, 101549, ISSN 2211-7156. <https://doi.org/10.1016/j.rechem.2024.101549>
42. Wang S., Ma F., Ma W., Wang P., Zhao G., and Lu X. Influence of Temperature on Biogas Production Efficiency and Microbial Community in a Two-Phase Anaerobic Digestion System. Water. 2019; 11(1):133. <https://doi.org/10.3390/w11010133>
43. Abanades S., Abbaspour H., Ahmadi A., Das B., Ehyaei M.A., Esmaeilion F., et al. A Critical Review Of Biogas Production And Usage With Legislations Framework Across The Globe. Int J Environ Sci Technol (Tehran). 2022;19(4):3377-3400. doi: 10.1007/s13762-021-03301-6. Epub 2021 May 16. PMID: 34025745; PMCID: PMC8124099.
44. Neupane D. Biofuels from Renewable Sources, a Potential Option for Biodiesel Production. Bioengineering (Basel). 2022 Dec 25;10(1):29. doi: 10.3390/bioengineering10010029. PMID: 36671601; PMCID: PMC9855116.
45. Aljaafari A., Fattah I.M.R., Jahiril M.I., Gu Y., Mahlia T.M.I., Islam M.A., and Islam M.S.. Biodiesel Emissions: A State-of-the-Art Review on Health and Environmental Impacts. Energies. 2022; 15(18):6854. <https://doi.org/10.3390/en15186854>
46. Jeswani H.K., Chilvers A., and Azapagic A. Environmental Sustainability Of Biofuels: A Review. Proc Math Phys Eng Sci. 2020 Nov;476(2243):20200351. doi: 10.1098/rspa.2020.0351. Epub 2020 Nov 25. PMID: 33363439; PMCID: PMC7735313.
47. Panwar NL, Kaushik SC, and Kothari S. Role of Renewable Energy Sources in Environmental Protection: A Review. *Renewable and Sustainable Energy Reviews*. 2011;15(3):1513-1524. <https://doi.org/10.1016/j.rser.2010.11.037>
48. United Nations Food and Agriculture Organization (FAO). The Role of Bioenergy in Energy Security. 2017. Accessed 14 August 2024. Available: <https://www.fao.org/energy/bioenergy/en/>
49. International Renewable Energy Agency (IRENA). Bioenergy from Boreal Forests: Swedish Approach to Sustainable Wood Use. 2021. Accessed 14 August 2024. Available: <https://www.irena.org/publications/2021/Apr/Bioenergy-from-boreal-forests>
50. Kalt G, & Kranzl L. Assessing the economic performance of bioenergy systems: A Review of Current Practices and Methods. *Renewable and Sustainable Energy Reviews*. 2011; 15(6): 3662-3671. <https://doi.org/10.1016/j.rser.2011.07.069>
51. United States Environmental Protection Agency. How Does Anaerobic Digestion Work? Last Updated on January 20, 2024. Accessed 14 August 2024. Available: <https://www.epa.gov/agstar/how-does-anaerobic-digestion-work>
52. Department of Environment and Conservation, State of Tennessee. Anaerobic Digestion: What is it? Accessed 14 August 2024. Available: <https://www.tn.gov/environment/program-areas/sw-mm-organics/anaerobic-digestion.html>
53. Odejebi O.J., Ajala O.O., and Osulolafe F.N. Review on Potential of Using Agricultural, Municipal Solid and Industrial Wastes as Substrates for Biogas Production in Nigeria. *Biomass Conv. Bioref*. 2024, 14,

- 1567–1579. <https://doi.org/10.1007/s13399-022-02613-y>
54. Kucharska K., Rybarczyk P., Hołowacz I., Łukajtis R., Glinka M., and Kamiński M. Pretreatment of Lignocellulosic Materials as Substrates for Fermentation Processes. *Molecules*. 2018; 23(11):2937. <https://doi.org/10.3390/molecules23112937>
  55. Manyi-Loh C.E., and Lues R. Anaerobic Digestion of Lignocellulosic Biomass: Substrate Characteristics (Challenge) and Innovation. *Fermentation*. 2023; 9(8):755. <https://doi.org/10.3390/fermentation9080755>
  56. Vítěz T., Novák D., Lochman J., and Vítězová M. Methanogens Diversity during Anaerobic Sewage Sludge Stabilization and the Effect of Temperature. *Processes*. 2020; 8(7):822. <https://doi.org/10.3390/pr8070822>
  57. Nie E., He P., Zhang H., Hao L., Shao L., and Lü F., How Does Temperature Regulate Anaerobic Digestion? *Renewable and Sustainable Energy Reviews*. 2021, Volume 150, 111453. ISSN 1364-0321. <https://doi.org/10.1016/j.rser.2021.111453>
  58. Meegoda JN, Li B, Patel K, Wang LB. A Review of the Processes, Parameters, and Optimization of Anaerobic Digestion. *International Journal of Environmental Research and Public Health*. 2018; 15(10):2224. <https://doi.org/10.3390/ijerph15102224>
  59. Anukam A, Mohammadi A, Naqvi M, Granström K. A Review of the Chemistry of Anaerobic Digestion: Methods of Accelerating and Optimizing Process Efficiency. *Processes*. 2019; 7(8):504. <https://doi.org/10.3390/pr7080504>
  60. Parajuli A, Khadka A, Sapkota L, Ghimire A. Effect of Hydraulic Retention Time and Organic-Loading Rate on Two-Stage, Semi-Continuous Mesophilic Anaerobic Digestion of Food Waste during Start-Up. *Fermentation*. 2022; 8(11):620. <https://doi.org/10.3390/fermentation8110620>
  61. Paranjpe A., Saxena S., and Jain P. Biogas Yield Using Single And Two Stage Anaerobic Digestion: An Experimental Approach. *Energy for Sustainable Development*. 2023, Volume 74, Pages 6-19. ISSN 0973-0826. <https://doi.org/10.1016/j.esd.2023.03.005>
  62. Perman E., Westerholm M., Liu T., and Schnürer A. Comparative Study of High-solid Anaerobic Digestion at Laboratory and Industrial Scale – Process Performance and Microbial Community Structure. *Energy Conversion and Management*. 2024, 300, 117978. <https://doi.org/10.1016/j.enconman.2023.117978>
  63. Tse TJ, Wiens DJ, Reaney MJT. Production of Bioethanol—A Review of Factors Affecting Ethanol Yield. *Fermentation*. 2021; 7(4):268. <https://doi.org/10.3390/fermentation7040268>
  64. Waseem W., Noor R.S. and Umair M. The Anaerobic Transformation Of Agricultural Waste For Bioethanol Production. *Biomass Conv. Bioref.* 14, 14163–14174 (2024). <https://doi.org/10.1007/s13399-023-04143-7>
  65. Maicas S. The Role of Yeasts in Fermentation Processes. *Microorganisms*. 2020; 8(8):1142. <https://doi.org/10.3390/microorganisms8081142>
  66. Iliev S. A Comparison of Ethanol, Methanol, and Butanol Blending with Gasoline and Its Effect on Engine Performance and Emissions Using Engine Simulation. *Processes*. 2021; 9(8):1322. <https://doi.org/10.3390/pr9081322>
  67. Rosales-Calderon O., and Arantes V. A Review On Commercial-Scale High-Value Products That Can Be Produced Alongside Cellulosic Ethanol. *Biotechnol Biofuels* 12, 240 (2019). <https://doi.org/10.1186/s13068-019-1529-1>
  68. Vasić K, Knez Ž, and Leitgeb M. Bioethanol Production by Enzymatic Hydrolysis from Different Lignocellulosic Sources. *Molecules*. 2021; 26(3):753. <https://doi.org/10.3390/molecules26030753>
  69. Zborowska M, Waliszewska H, Waliszewska B, Borysiak S, Brozdowski J, Stachowiak-Wencek A. Conversion of Carbohydrates in Lignocellulosic Biomass after Chemical Pretreatment. *Energies*. 2022; 15(1):254. <https://doi.org/10.3390/en15010254>
  70. Yi T, Zhao H, Mo Q, Pan D, Liu Y, Huang L, Xu H, Hu B, Song H. From Cellulose to Cellulose Nanofibrils—A Comprehensive Review of the Preparation and Modification of Cellulose Nanofibrils. *Materials*. 2020; 13(22):5062. <https://doi.org/10.3390/ma13225062>
  71. Osman, A.I., Mehta, N., Elgarahy, A.M. et al. Conversion Of Biomass To Biofuels And Life Cycle Assessment: A Review. *Environ Chem Lett* 19, 4075–4118 (2021). <https://doi.org/10.1007/s10311-021-01273-0>
  72. Obileke K, Makaka G, Nwokolo N. Recent Advancements in Anaerobic Digestion and Gasification Technology. *Applied Sciences*. 2023; 13(9):5597. <https://doi.org/10.3390/app13095597>
  73. De Souza R, Casisi M, Micheli D, Reini M. A Review of Small–Medium Combined Heat and Power

- (CHP) Technologies and Their Role within the 100% Renewable Energy Systems Scenario. *Energies*. 2021; 14(17):5338. <https://doi.org/10.3390/en14175338>
74. Sevillano CA, Pesantes AA, Peña Carpio E, Martínez EJ, Gómez X. Anaerobic Digestion for Producing Renewable Energy—The Evolution of This Technology in a New Uncertain Scenario. *Entropy*. 2021; 23(2):145. <https://doi.org/10.3390/e23020145>
  75. Nazari, L., Xu, C.(., Ray, M.B. (2021). Advanced Technologies (Biological and Thermochemical) for Waste-to-Energy Conversion. In: *Advanced and Emerging Technologies for Resource Recovery from Wastes. Green Chemistry and Sustainable Technology*. Springer, Singapore. [https://doi.org/10.1007/978-981-15-9267-6\\_3](https://doi.org/10.1007/978-981-15-9267-6_3)
  76. Variny M, Varga A, Rimár M, Janošovský J, Kizek J, Lukáč L, Jablonský G, Mierka O. Advances in Biomass Co-Combustion with Fossil Fuels in the European Context: A Review. *Processes*. 2021; 9(1):100. <https://doi.org/10.3390/pr9010100>
  77. Nunes LJR, Meireles CIR, Pinto Gomes CJ, Almeida Ribeiro NMC. Forest Management and Climate Change Mitigation: A Review on Carbon Cycle Flow Models for the Sustainability of Resources. *Sustainability*. 2019; 11(19):5276. <https://doi.org/10.3390/su11195276>
  78. Lewandowski WM, Ryms M, Kosakowski W. Thermal Biomass Conversion: A Review. *Processes*. 2020; 8(5):516. <https://doi.org/10.3390/pr8050516>
  79. Galko G, Sajdak M. Trends for the Thermal Degradation of Polymeric Materials: Analysis of Available Techniques, Issues, and Opportunities. *Applied Sciences*. 2022; 12(18):9138. <https://doi.org/10.3390/app12189138>
  80. Ore O.T., and Adebisi F.M. A Review On Current Trends And Prospects In The Pyrolysis Of Heavy Oils. *J Petrol Explor Prod Technol* 11, 1521–1530 (2021). <https://doi.org/10.1007/s13202-021-01099-0>
  81. Fahmy, T.Y.A., Fahmy, Y., Mobarak, F. et al. Biomass Pyrolysis: Past, Present, And Future. *Environ Dev Sustain* 22, 17–32 (2020). <https://doi.org/10.1007/s10668-018-0200-5>
  82. Teh JS, Teoh YH, How HG, Sher F. Thermal Analysis Technologies for Biomass Feedstocks: A State-of-the-Art Review. *Processes*. 2021; 9(9):1610. <https://doi.org/10.3390/pr9091610>
  83. Raza M, Inayat A, Ahmed A, Jamil F, Ghenai C, Naqvi SR, et al. Progress of the Pyrolyzer Reactors and Advanced Technologies for Biomass Pyrolysis Processing. *Sustainability*. 2021; 13(19):11061. <https://doi.org/10.3390/su131911061>
  84. Mensah, R.A., Jiang, L., Renner, J.S. et al. Characterisation Of The Fire Behaviour Of Wood: From Pyrolysis To Fire Retardant Mechanisms. *J Therm Anal Calorim* 148, 1407–1422 (2023). <https://doi.org/10.1007/s10973-022-11442-0>
  85. Apaydin Varol E, Mutlu Ü. TGA-FTIR Analysis of Biomass Samples Based on the Thermal Decomposition Behavior of Hemicellulose, Cellulose, and Lignin. *Energies*. 2023; 16(9):3674. <https://doi.org/10.3390/en16093674>
  86. Glushkov D, Nyashina G, Shvets A, Pereira A, Ramanathan A. Current Status of the Pyrolysis and Gasification Mechanism of Biomass. *Energies*. 2021; 14(22):7541. <https://doi.org/10.3390/en14227541>
  87. Sieradzka M, Mlonka-Mędrala A, Kalemba-Rec I, Reinmöller M, Küster F, Kalawa W, Magdziarz A. Evaluation of Physical and Chemical Properties of Residue from Gasification of Biomass Wastes. *Energies*. 2022; 15(10):3539. <https://doi.org/10.3390/en15103539>
  88. Trubetskaya A. Reactivity Effects of Inorganic Content in Biomass Gasification: A Review. *Energies*. 2022; 15(9):3137. <https://doi.org/10.3390/en15093137>
  89. Block, C., Ephraim, A., Weiss-Hortala, E. et al. Co-pyrogasification of Plastics and Biomass, A Review. *Waste Biomass Valor* 10, 483–509 (2019). <https://doi.org/10.1007/s12649-018-0219-8>
  90. Jha S, Nanda S, Acharya B, Dalai AK. A Review of Thermochemical Conversion of Waste Biomass to Biofuels. *Energies*. 2022; 15(17):6352. <https://doi.org/10.3390/en15176352>
  91. Monteiro E, Ferreira S. Some Perspectives for the Gasification Process in the Energy Transition World Scenario. *Energies*. 2023; 16(14):5543. <https://doi.org/10.3390/en16145543>
  92. Laredo GC, Reza J, Meneses Ruiz E. Hydrothermal Liquefaction Processes For Plastics Recycling: A Review. *Cleaner Chemical Engineering*. 2023;5:100094. DOI:10.1016/j.clce.2023.100094
  93. Mahima J, Sundaresh RK, Gopinath KP, et al. Effect Of Algae (*Scenedesmus Obliquus*) Biomass Pre-Treatment On Bio-Oil Production In Hydrothermal Liquefaction (Htl): Biochar And Aqueous Phase Utilization Studies. *Sci Total Environ*. 2021;778:146262. DOI:10.1016/j.scitotenv.2021.146262
  94. López Barreiro D, Prins W, Ronsse F, Brilman W. Hydrothermal Liquefaction (Htl) Of Microalgae For

- Biofuel Production: State Of The Art Review And Future Prospects. *Biomass and Bioenergy*. 2013;53:113-127. DOI:10.1016/j.biombioe.2012.12.029
95. Al-Jabri H, Das P, Khan S, Thaher M, Abdulquadir M. Treatment Of Wastewaters By Microalgae And The Potential Applications Of The Produced Biomass—A Review. *Water (Basel)*. 2020;13(1):27. DOI:10.3390/w13010027
  96. Ahmad A, Banat F, Alsafar H, Hasan SW. Algae Biotechnology For Industrial Wastewater Treatment, Bioenergy Production, And High-Value Bioproducts. *Sci Total Environ*. 2022;806(Pt 2):150585. DOI:10.1016/j.scitotenv.2021.150585
  97. Chen H, Xia A, Zhu X, Huang Y, Zhu X, Liao Q. Hydrothermal Hydrolysis Of Algal Biomass For Biofuels Production: A Review. *Bioresour Technol*. 2022;344(Pt B):126213. DOI:10.1016/j.biortech.2021.126213
  98. Galford GL, Peña O, Sullivan AK, et al. Agricultural Development Addresses Food Loss And Waste While Reducing Greenhouse Gas Emissions. *Sci Total Environ*. 2020;699:134318. DOI:10.1016/j.scitotenv.2019.134318
  99. Yaman C, Anil I, Jaunich MK, et al. Investigation And Modelling Of Greenhouse Gas Emissions Resulting From Waste Collection And Transport Activities. *Waste Manag Res*. 2019;37(12):1282-1290. DOI:10.1177/0734242X19882482
  100. Ambaye TG, Vaccari M, Bonilla-Petriciolet A, Prasad S, van Hullebusch ED, Rtimi S. Emerging technologies for biofuel production: A critical review on recent progress, challenges and perspectives. *J Environ Manage*. 2021;290:112627. DOI:10.1016/j.jenvman.2021.112627
  101. Tshemese Z, Deenadayalu N, Liganiso LZ, Chetty M. An Overview of Biogas Production from Anaerobic Digestion and the Possibility of Using Sugarcane Wastewater and Municipal Solid Waste in a South African Context. *ASL*. 2023;6(1):13. DOI:10.3390/asi6010013
  102. Piadeh F, Offie I, Behzadian K, et al. A critical review for the impact of anaerobic digestion on the sustainable development goals. *J Environ Manage*. 2024;349:119458. DOI:10.1016/j.jenvman.2023.119458
  103. Reetsch A, Schwärzel K, Kapp G, et al. Data set of smallholder farm households in banana-coffee-based farming systems containing data on farm households, agricultural production and use of organic farm waste. *Data Brief*. 2021;35:106833. DOI:10.1016/j.dib.2021.106833
  104. Kannankai MP, Devipriya SP. Air quality impacts of landfill fires: A case study from the Brahmapuram Municipal Solid Waste Treatment Plant in Kochi, India. *Sci Total Environ*. 2024;916:170289. DOI:10.1016/j.scitotenv.2024.170289
  105. Traven L. Sustainable energy generation from municipal solid waste: A brief overview of existing technologies. *Case Studies in Chemical and Environmental Engineering*. 2023;8:100491. DOI:10.1016/j.csee.2023.100491
  106. Kabir E, Kim K-H, Kwon EE. Biochar as a tool for the improvement of soil and environment. *Front Environ Sci*. 2023;11. DOI:10.3389/fenvs.2023.1324533
  107. Haider G, Joseph S, Steffens D, et al. Mineral nitrogen captured in field-aged biochar is plant-available. *Sci Rep*. 2020;10(1):13816. DOI:10.1038/s41598-020-70586-x
  108. Obeng GM, Aram SA, Agyei D, Saalidong BM. Exposure to particulate matter (PM2.5) and volatile organic compounds (VOCs), and self-reported health symptoms among fish smokers: A case study in the Western Region of Ghana. *PLoS ONE*. 2023;18(3):e0283438. DOI:10.1371/journal.pone.0283438
  109. Yao W, Zhao Y, Chen R, Wang M, Song W, Yu D. Emissions of Toxic Substances from Biomass Burning: A Review of Methods and Technical Influencing Factors. *Processes*. 2023;11(3):853. DOI:10.3390/pr11030853
  110. Tamire M, Kumie A, Addissie A, et al. High Levels of Fine Particulate Matter (PM2.5) Concentrations from Burning Solid Fuels in Rural Households of Butajira, Ethiopia. *Int J Environ Res Public Health*. 2021;18(13). DOI:10.3390/ijerph18136942
  111. Bari MA, Kindziński WB. Ambient fine particulate matter (PM2.5) in Canadian oil sands communities: Levels, sources and potential human health risk. *Sci Total Environ*. 2017;595:828-838. DOI:10.1016/j.scitotenv.2017.04.023
  112. He D, Xiao J, Wang D, et al. Digestion liquid based alkaline pretreatment of waste activated sludge

- promotes methane production from anaerobic digestion. *Water Res.* 2021;199:117198. DOI:10.1016/j.watres.2021.117198
113. Lavery AM, Backer LC, Roberts VA, DeVies J, Daniel J, DHSc1. Evaluation of Syndromic Surveillance Data for Studying Harmful Algal Bloom-Associated Illnesses - United States, 2017-2019. *MMWR Morb Mortal Wkly Rep.* 2021;70(35):1191-1194. DOI:10.15585/mmwr.mm7035a2
  114. Water Use, Water Pollution, and Biofuels - The Nexus of Biofuels, Climate Change, and Human Health - NCBI Bookshelf. Accessed August 16, 2024. Available: <https://www.ncbi.nlm.nih.gov/books/NBK196445/>
  115. Vera I, Wicke B, Lamers P, et al. Land use for bioenergy: Synergies and trade-offs between sustainable development goals. *Renew Sustain Energy Rev.* 2022;161:112409. DOI:10.1016/j.rser.2022.112409
  116. New IUCN report: To fulfill its promise, circular economy must take biodiversity into account - News | IUCN. Accessed August 16, 2024. Available: <https://iucn.org/news/202212/new-iucn-report-fulfil-its-promise-circular-economy-must-take-biodiversity-account>
  117. Nordin I. Cost- effective use of abandoned agricultural land for biofuel production. *Glob Change Biol Bioenergy.* 2024;16(7). DOI:10.1111/gcbb.13165
  118. Núñez-Regueiro MM, Siddiqui SF, Fletcher RJ. Effects of bioenergy on biodiversity arising from land-use change and crop type. *Conserv Biol.* 2021;35(1):77-87. DOI:10.1111/cobi.13452
  119. Breuer JL, Samsun RC, Peters R, Stolten D. The impact of diesel vehicles on NOx and PM10 emissions from road transport in urban morphological zones: A case study in North Rhine-Westphalia, Germany. *Sci Total Environ.* 2020;727:138583. DOI:10.1016/j.scitotenv.2020.138583
  120. Scheutz C, Fredenslund AM. Total methane emission rates and losses from 23 biogas plants. *Waste Manag.* 2019;97:38-46. DOI:10.1016/j.wasman.2019.07.029
  121. Gao Y, Wang M, Raheem A, et al. Syngas Production from Biomass Gasification: Influences of Feedstock Properties, Reactor Type, and Reaction Parameters. *ACS Omega.* 2023;8(35):31620-31631. DOI:10.1021/acsomega.3c03050
  122. Hofsetz K, Silva MA. Brazilian sugarcane bagasse: Energy and non-energy consumption. *Biomass and Bioenergy.* 2012;46:564-573. DOI:10.1016/j.biombioe.2012.06.038
  123. Gebrezgabher SA, Taron A, Odero J, et al. Circular bioeconomy business models - energy recovery from agricultural waste: cases from Kenya and Burkina Faso. Published online December 1, 2022.
  124. Uddin MM, Wright MM. Anaerobic digestion fundamentals, challenges, and technological advances. *Physical Sciences Reviews.* 2022;0(0). DOI:10.1515/psr-2021-0068
  125. Adekunle KF, Okolie JA. A review of biochemical process of anaerobic digestion. *ABB.* 2015;06(03):205-212. DOI:10.4236/abb.2015.63020

## ABBREVIATIONS

HTL - Hydrothermal Liquefaction  
 CHP - Combined Heat and Power