

Genetic Potential of Sesame: A Study on Variability, Heritability and Genetic Advance

ABSTRACT

Aims: This study aimed to evaluate 140 genotypes of sesame germplasms along with five check varieties to assess heritability, variability, and genetic advance for yield and yield-related agronomic characters, with the goal of improving yield potential through breeding programs.

Study Design: The evaluation was conducted using an augmented design with four blocks during the kharif season of 2021.

Methodology: Significant differences among the germplasm lines were observed for days to 50 percent flowering, days to maturity, plant height, capsule-bearing length, capsules per plant, seeds per capsule, branches per plant, capsule length, chlorophyll content, test weight, and seed yield. This indicates substantial variability among the germplasm lines. Traits such as seed yield per plant, capsules per plant, and primary branches per plant exhibited high phenotypic and genotypic coefficients of variation. High heritability coupled with high genetic advance as a percentage of the mean was noted for seed yield per plant, capsules per plant, primary branches per plant, and test weight, suggesting the influence of additive gene action, thereby making simple selection effective for improving these traits.

Results: The analysis of variance showed significant differences among the germplasm lines for various agronomic traits, indicating substantial variability. Traits like seed yield per plant, capsules per plant, and primary branches per plant exhibited high phenotypic and genotypic coefficients of variation, coupled with high heritability and genetic advance, suggesting the potential for simple selection to improve these traits.

Conclusion: The evaluation of sesame germplasms revealed significant variability for various yield and agronomic traits, with traits like seed yield per plant, capsules per plant, and primary branches per plant showing potential for improvement through breeding programs. High heritability and genetic advance suggest the influence of additive gene action, facilitating effective selection strategies for enhancing yield potential in sesame.

Keywords: - Sesame, Variability, Heritability, Genetic Advance, Seed yield.

1. INTRODUCTION

Sesame (*Sesamum indicum* L.) is one of the oldest oilseed crop in world and it is native to tropic and sub-tropic regions. It's known with various names such as sesamum, til, gingelly, simsim, gergelim etc. It is an annual, self-pollinated oil seed crop. The genus *sesamum* belongs to the order *tubiflorea*, family *pedaliaceae* which consists of sixteen genera and sixty species (Weiss, 1983), but only *sesamum indicum* ($2n = 26$) has been recognized as cultivated species. Several historical records indicated that sesame probably originated in Ethiopia (Africa) and from there, it was introduced into India and China and became a popular food in Southern Asia, South Europe and North-East Africa. It spread into West Asia and then to India, China and Japan and became secondary distribution centers.

India is the second world leader with the area (1.73 million hectares), production (0.746 million tonnes) and productivity (431 kg ha^{-1}) (FAOSTAT, 2020). Gujarat, Rajasthan, Uttar Pradesh, Madhya Pradesh, Maharashtra, Andhra Pradesh, Orissa, Tamil Nadu, West Bengal and Karnataka are the major sesame growing states of the country. The important districts cultivating sesame in Rajasthan state are Pali, Nagaur, Jodhpur, Jalore, Bhilwara, Sirohi, Ajmer and Alwar. In Rajasthan, sesame occupies area 0.29 million hectares and production 0.077 million tonnes with productivity 266 kg ha^{-1} (Anonymous, 2022).

Inspite of sesame possessing high nutritional value and resistance to abiotic stress like drought it has low yielding capacity compared to other oilseed crops. The major problems in sesame cultivation are non-synchronous maturity, indeterminate growth habit narrow adaptability, yield instability, seed shattering, non-availability of high yielding varieties

with resistance to stresses and the presence of fertilization barriers *etc.* (Rao *et al.*, 2002), and strong competition from other oil crops such as soybean, sunflower and peanut. Further, sesame has been given less attention by the farmers because of poor yield due to non-availability of cultivars to suit the diverse agro climatic conditions. Therefore, development of improved high yielding varieties adapted to local conditions has become top priority.

The development of improved cultivars is mainly governed by the magnitude of genetic diversity in the base material and the extent of variability for the desired characters. Genetic variability is of great interest to the plant breeder as it plays a vital role in framing a successful breeding programme.

Heritability is the transmissibility of a character into future generations. Genetic advance refers to the difference between the mean genotypic values of selected population and the original population from which these were selected. Heritability estimates together with genetic advance is generally more helpful in prognosticating the genetic gain under selection than heritability estimates alone.

2. MATERIAL AND METHODS

The present investigation was conducted at the institutional farm, S.K.N. College of Agriculture, Jobner during *kharif* season, 2021. The experimental material consisted of 140 sesame germplasm lines along with 5 check varieties *viz.*, RT 372, TKG 22, TC 25, RT 346 and PARGATI obtained from Germplasm Evaluation Division, ICAR-National Bureau of Plant Genetic Resources, New Delhi. The experimental material was evaluated in Augmented Design during *kharif*, 2021. Each germplasm line was sown in a plot of single 4 meter row length with the whole germplasm lines was distributed in 4 blocks with 5 check varieties. Each check varieties was sown after 7 germplasm lines. Row to row and plant to plant distance was maintained 30 cm and 15 cm, respectively. All the agronomical practices were followed to raise a good and healthy crop. The data of were recorded from five plants were selected randomly and tagged before flowering from each entry. However, data on days to 50 per cent flowering, days to maturity and 1000-seed weight was recorded on whole plot basis. The observations were recorded on following characters: days to 50 per cent flowering, days to maturity, plant height, capsule bearing length, capsules per plant, capsules on main stem, seeds per capsule, branches per plant, capsule length, chlorophyll content, 1000-seed weight and seed yield per plant.

To test the variation among the germplasm lines analysis of variance *via* adopting standard statistical methods (Federer, 1956). The phenotypic coefficient of variation (PCV) and genotypic coefficient of variance (GCV) were estimated using the formula proposed by Burton (1952) and Johnson *et al.* (1955). Heritability in broad sense was calculated according to the formula suggested by Johnson *et al.* (1955). From the heritability estimates, the genetic advance and genetic gain was calculated by the formula given by Johnson *et al.* (1955).

3. RESULTS AND DISCUSSION

Analysis of variance (Table1) revealed significant differences among the genotypes for different characters except capsules on main stem indicating presence of sufficient variability in the material. Mean, range, genotypic and phenotypic coefficient of variation, heritability (broad sense) and genetic gain were analyzed are shown in Table 2.

PCV and GCV estimates are closer and parallel for all the characters, indicating that these characters are less influenced by environment and comparatively stable. A wide range of variability was exhibited for characters such as seed yield per plant, capsules per plant, and primary branches per plant indicating that there is scope for selection of suitable initial material in breeding for further improvement.

Genetic variability is a basic information needed for the breeders to improve the crops by adopting appropriate method of selection based on variability that exist in the material. In this regard, it is necessary to partition the total variability into heritable and non-heritable components *viz.*, genotypic coefficient of variation (GCV), phenotypic coefficient of variation (PCV) and further to compute heritability and genetic advances for various quantitative traits. Comparison of variability between two traits is possible with coefficient of variation as it is free of units. As expected, the PCV values were greater than the GCV values for all the characters indicating considerable influence of environment on the expression of these characters under field conditions.

The phenotypic and genotypic coefficient of variation of germplasm was high for seed yield per plant, capsules per plant and branches per plant. Similar observations were reported by Sumathi and Muralidharan (2010), Gangadhara *et al.* (2012), Narayanan and Murugan (2013), Bharathi *et al.* (2014), Ismaila and Usman (2014), Teklu *et al.* (2014, 2017) and Aristya *et al.* (2017) for capsules per plant and seed yield per plant, Chandra (2014) and Pavani *et al.* (2020) for branches per plant and Kumar *et al.* (2022) for seed yield per plant. High PCV and GCV estimates for characters indicate existence of the substantial variability in these characters.

The values of phenotypic and genotypic coefficients of variation were moderate for plant height, capsule length and seeds per capsule, test weight, capsule bearing length, capsules on main stem, chlorophyll content and days to 50 per cent flowering. These results were in accordance with the reports of Soundharya (2016) and Bharathi *et al.* (2014) for capsule length, Teklu *et al.* (2017) and Ismail and Mohamed (2018) for capsule bearing length and Gangadhara *et al.* (2012) for capsule length, test weight and seeds per capsule.

The values of PCV and GCV were low for days to maturity. This result was in accordance with the findings of Gangadhara *et al.* (2012) and Pavani *et al.* (2020).

Genetic coefficient of variation does not provide the clear indication of proportion of heritable component of variation. The heritability estimate of a quantitative character is very important as phenotypic expression of a genotype may be altered by environment at various stages of its development.

Heritability (in broad sense) was high (>60 per cent) for plant height, capsule length, primary branches per plant, chlorophyll content, days to 50 per cent flowering, capsules per plant, test weight, seeds per capsule, seed yield per plant, days to maturity and capsule bearing length except capsules on main stem indicating that characters were least influenced by environment and selection of such characters may be useful. Similar results were shown earlier by Ahemd and Ahemd (2013), Saxena *et al.* (2016) and Teklu *et al.* (2017) for days to 50 per cent flowering; Sumathi and Murlidharan (2010) and Hika *et al.* (2015) for days to maturity; Teklu *et al.* (2017) for plant height and capsules per plant; Krishnaiah *et al.* (2002) for seeds per capsule; Ashokavaradhan *et al.* (2001) for test weight; Soundharya *et al.* (2016) and Abhijatha *et al.* (2017) for primary branches per plant and Saxena and Bisen (2017) and Ismail and Mohamed (2018) for seed yield per plant.

Heritability estimates alone do not provide reliable information about the gene action governing the expression of a particular character. High heritability coupled with high genetic advance as per cent of mean would be more reliable and useful in formulating selection procedure (Johnson *et al.*, 1955). High heritability coupled with high genetic advance as per cent of mean was observed for most of the characters *viz.*, seed yield per plant, capsules per plant, branches per plant, test weight, capsule bearing length, plant height, chlorophyll content, days to 50 per cent flowering and capsule length. Similar results have been reported earlier by Teklu *et al.* (2017) for days to 50 per cent flowering; Patil and Loksha (2018) and Teklu *et al.* (2017) for plant height, Ismaila and Usman (2014) and Revathy *et al.* (2012) for primary branches per plant and Revathy *et al.* (2012), Ismaila and Usman (2014) and Hika *et al.* (2015) for seed yield per plant. High heritability combined with high genetic advance indicated that these traits were governed largely through additive effect of genes and improvement in these characters may be achieved through simple phenotypic selection.

High heritability coupled with moderate genetic advances as per cent mean was recorded for seeds per capsule revealed the predominance of both additive and non-additive gene action. This offers the best opportunity for improvement of this trait through mass selection and progeny selection. And High heritability coupled with low genetic advances as per cent of mean was recorded for days to maturity which indicates dominant and epistatic gene action, it is not effective for selection. Similar results have been reported earlier by Ahmed and Ahmed (2013) and Pavani *et al.* (2020).

Table 1: Mean sum of squares and variance for different quantitative characters in Sesame

Source of variation	d.f	Days to 50 per cent flowering	Days to maturity	Plant height (cm)	Capsule bearing length (cm)	Capsules per plant	Seeds per capsule	Capsules on main stem	Branches per plant	Capsule length (cm)	Chlorophyll content (SPAD)	Test weight (g)	Seed yield per plant (g)
Blocks(b-1)	3	2.27	0.58	0.19	126.85	96.58**	2.93	8.73	0.03	0.02**	5.49	0.01	0.01
Entries (c+g)-1	14	32.05**	11.40*	417.71	157.6	200.77	36.20*	9.54**	2.13**	0.07**	55.60**	0.53*	1.42
Checks (c-1)	4	6.57**	8.70*	62.06**	133.61**	63.07**	23.20*	17.82**	11.96**	0.01**	20.03**	0.30*	1.80
Germplasm(g-1)	13	28.18**	10.70*	386.57	155.9	190.78	32.03*	8.71	1.86**	0.06**	48.59**	0.50*	1.40
Checks Vs	1	672.70	119.6	6169.2	495.8	2139.0	668.6	91.89**	0.30**	0.74**	1171.79*	4.88*	2.25

Germplas	**	0**	3**	2**	0**	8**				*	*	**	
m													
Error (b-1)	12	1.48	2.67	0.25	38.93	18.21	5.10	4.02	0.06	0.00	2.33	0.07	0.28
(c-1)													

*, ** Significant at 5 per cent and 1 per cent level of significance, respectively

Table 2: Variability parameters for different quantitative characters of sesame

Characters	Mean	Range	GCV	PCV	Heritability (per cent)	G.A. ssas cent mean per of
Days to 50 per cent flowering	41.80	30.00 – 52.00	12.36	12.70	94.76	24.79
Days to maturity	91.24	84.00 – 98.00	3.11	3.59	75.08	5.54
Plant height (cm)	94.87	30.60 - 132.40	20.72	20.72	99.94	42.66
Capsule bearing length (cm)	41.20	10.00 - 70.40	26.26	30.31	75.04	46.86
Capsules per plant	33.71	12.40 - 80.20	38.97	40.98	90.46	76.36
Seeds per capsule	51.22	15.80 - 60.80	10.13	11.05	84.08	19.14
Capsules on main stem	7.81	2.40 - 14.20	27.71	37.79	53.77	41.85
Branches per plant	3.60	1.20 - 8.40	37.26	37.86	96.86	75.53
Capsule length (cm)	2.09	1.40 - 2.90	11.85	12.04	97.23	24.07
Chlorophyll content (SPAD)	39.05	25.07 - 51.07	17.42	17.85	95.20	35.01
Test weight	2.35	1.00 - 4.37	27.87	30.17	85.33	53.03
Seed yield per plant (g)	2.34	1.03 - 5.17	45.37	50.65	80.24	83.73

4. CONCLUSION

In conclusion, our study highlights significant genetic variability in sesame germplasms, with traits like seed yield per plant, capsules per plant, and primary branches per plant showing promising potential for selection. High heritability coupled with substantial genetic advance suggests the predominance of additive gene action, facilitating straightforward phenotypic selection for improvement. These findings offer valuable guidance for sesame breeders in developing improved varieties with enhanced yield potential and overall agronomic performance.

REFERENCES

- Abhijatha, A., Arya, K., Madhu kar, K. and Gogineni, S. 2017. Evaluation of sesame (*Sesamum indicum* L.) genotypes to the shaded uplands of southern region. *International Journal of Current Microbiology and Applied Sciences*, **6**(7): 332-339.
- Anonymous, 2019. All India Estimates of Area, Production and Yield of Foodgrains, Directorate of Economics and Statistics, Department of Agriculture and Farmers Welfare, Ministry of Agriculture and Farmers Welfare, Government of India New Delhi – 110 001.
- Anonymous, 2022. Crop-wise Area Production and Yield of various Principal Crops forth estimates. Commissionerate of Agriculture, Government of Rajasthan, Jaipur.
- Aristya, V.E., Taryono and Wulandari, 2017. Genetic variability standardized multiple liner regression and principle component analysis to determine some important sesame yield components. *Agrivita Journal of Agriculture Science*, **39**(1): 83- 90.
- Ashokavardhana, R.P., Reddy Shekar, M. Ranganatha, A.R.G. and Dhanraj, A., 2001, Genetic variability and heritability for seed yield and its components in sesame (*Sesamum indicum* L.). *J. Oilseed Res.*, **18**:173-175.
- Bharathi, D., Thirumala Rao, V., Chandra Mohan, Y., Bhadru, D. and Venkanna, V. 2014. Genetic variability studies in sesame (*Sesamum indicum* L.). *International Journal of Applied Biology and Pharmaceutical Technology*, **5**(4): 31-33.
- Burton, J. W. 1952. Quantitative inheritance in grasses. Proceeding of *International 6th Grassland Congress*, **1**: 277-283.
- Chandra, M.Y. 2014. Variability and genetic divergence in sesame (*Sesamum indicum* L.). *International Journal of Applied Biology and Pharmaceutical Technology*, **5**(3): 222-225.
- Federer, W.T. 1956. Augmented Design. *Hawain Planters Record*. **20**: 191-207.
- FAOSTAT,2020. Food and Agriculture Organization Statistical Database. Available online :<http://faostat.fao.org>.
- Gangadhara, J., Prakash, C., Badiger, B, Shadakshari, T.V., Yathish, K.R. and Rajesh, A.M. 2012. Genetic divergence, genetic advance and heritability in sesame (*Sesamum indicum* L.). *BiolInfolet*.**9** (5): 437-462.
- Hika, G., Gelata N. and Jaleta, Z. 2014. Correlation and Divergence analysis for phenotypic traits in sesamum (*Sesamum indicum* L.) genotypes. *Science Technology and Arts Research Journal*, **3**(4): 01-09.
- Hika, G., Geleta, N. and Jaleta, Z. 2015. Genetic variability, heritability and genetic advance for the phenotypic traits in sesame (*Sesamum indicum* L.). Populations from Ethiopia. *Science, Technology and Arts Research Journal*. **4**(1): 20-26.
- Ismail, M.B. and Mohamed, N.E. 2018. Phenotypic and genotypic variability in a set of sesame (*Sesamum indicum* L.) Genotypes. *Egyptian Journal Agronomy*, **40**(3).
- Ismaila, A. and Usman, A. 2014. Genetic variability for yield and yield components in sesame (*Sesamum indicum* L.). *International Journal of Science and Research*. **3**(9): 63-66.
- Johnson, H. W., Robinson, H. F. and Comstock, R. E. 1955. Estimation of genetic variability and environmental variability in soyabean. *Agronomy Journal*, **47**: 314-318.
- Krishnaiah, G., Raja Reddy and Reddy Shekar, M., 2002, Variability studies in sesame (*Sesamum indicum* L.). *Crop Res.*, **24**: 501-504.
- Kumar, V., Sinha, S., Tomar, S., Sinha, S., Singh, R. S. and Singh, S. N. (2022). Assessment of genetic variability, correlation and path analysis in sesame (*Sesamum indicum* L.). *Electronic Journal of Plant Breeding*, **13**(1), 208-215.
- Narayanan, R. and Murugan, S. 2013. Studies on variability and heritability in sesame (*Sesamum indicum* L.). *International Journal of Current Agricultural Research*, **2** (11): 052-055.

- Patil, M.K. and Lokesh, R. 2018. Estimation of genetic variability, heritability, genetic advance, correlation and path analysis in advance mutant breeding lines of sesame (*Sesamum indicum* L.). *Green Farming*, **9**(2): 257-260.
- Pavani K., Lal Ahamed, M., Ramana, J.V. and Sirisha A.B.M. 2020. Studies on genetic variability parameter in sesame (*Sesamum indicum* L.) *International Journal of Chemical Studies*, **8**(4): 101-104.
- Rao, K.R. Kishor, P.K. and Vaidyanath, K. 2002. Biotechnology of sesame an oil seed crop. *Plant Cell Biotechnology and Molecular Biology*, **3**(3-4): 101-110.
- Revathi, S., John, J.A. and Manivannan, N. 2012. Genetic variability in sesame (*Sesamum indicum* L.). *Electronic Journal of Plant Breeding*, **3**(1):692-694.
- Saxena, K. and Bisen, R. 2017. Genetic variability, heritability and genetic advance for the phenotypic traits in sesame (*Sesamum indicum* L.). *International Journal of Pure and Applied Bio-Science*. **5**(2): 1126-1131.
- Sumathi, P and Muralidharan, V. 2010. Analysis of genetic variability, association and path analysis in the hybrids of sesame (*Sesamum indicum* L.). *Tropical Agricultural Research and Extension* **13**(3) 63-67.
- Teklu, D.H., Kebede, S.A. and Gebremichael, D.E. 2014. Assessment of genetic variability, genetic advance, correlation and path analysis for morphological traits in sesame genotypes. *Asian Journal of Agricultural Research*, pp. 181-194.
- Teklu, D.H., Kebede, S.A. and Gebremichael, D.E. 2017. Assessment of genetic variability, genetic advance, correlation and path analysis for morphological traits in sesame genotypes. *International Journal of Novel Research in Life Sciences*, **4**(2): 34-44.
- Weiss, E. A., 1971, Castor, Sesame and Safflower, Leonard Hill Books, London, pp. 311-355.
- Weiss, E. A., 1983, Oilseed crops, London, pp. 280-294.