

Case report

Clinical Management of Postpartum Cervico-Vaginal Prolapse In Graded Murrah Buffalo: A Case Report

Comment [H1]: give a gap

ABSTRACT

The present case report describes the postpartum cervico-vaginal prolapse and its clinical management in a graded Murrah buffalo with the history of parturition a day ago, presented to the Veterinary Clinical Complex, C.V.Sc., Korutla. The prolapsed mass was having lacerations, mild straining and bleeding. After thorough cleaning and reduction of edema, the prolapsed mass was properly repositioned. A rope truss was applied to prevent reoccurrence of prolapse. The animal showed a good response to manual correction and no complications were observed. Early identification and intimate veterinary intervention may help in effective management of postpartum cervico-vaginal prolapse in a buffalo.

Comment [H2]: give a gap

Comment [H3]: college name is not needed as you will mention in the body of the article

Comment [H4]: Early diagnosis

Comment [H5]: timely intervention will help

Keywords: Graded murrah buffalo, Post partum, Cervico-vaginal prolapse, Rope truss method.

1. INTRODUCTION

Buffaloes are very important in Indian dairy industry as they contribute 44.81% of total milk production [1]. Various health and reproductive issues will show direct effect on its milk yield. Prolapse of genitalia is one of the major maternal reproductive problem commonly seen in advanced gestation or postpartum conditions [2]. In cervico-vaginal condition cervix and vagina protrudes out of the vulvar lips. Showing the rede, throwing the rede, throwing the rose, bearing the trouble and pushing out the button are some other names known [3].

Higher estrogen levels in third trimester will increase relaxation of pelvic structures and decreased levels of calcium may result in reduced tonicity of vaginal and uterine muscles which predisposes the vaginal prolapse in animals [4]. Early finding and proper treatment are requirements in cervico-vaginal prolapse in buffaloes to save its health

and breeding life [5]. The present case report describes the condition of cervico-vaginal prolapse in Murrah buffalo and its management.

2. CASE PRESENTATION

A 3 years old graded Murrah buffalo was presented to Veterinary Clinical Complex, C.V.Sc. Korutla with the history of cervico-vaginal prolapse 6 hours after the normal calving (fig.1). Initial attempts to replace the prolapse mass was performed by owners at the farm level. Feed and water intake were reduced with ceased urination. Animal was restless with rectal temperature of 101.3°F and pale conjunctival mucous membrane. On clinical observation edematous, swollen prolapsed mass noticed in between vulvar lips with severe straining.

Comment [H10]: case history and observation

Comment [H6]: reproductive disorder, seen

Comment [H7]: give space

Comment [H11]: write complete name

Comment [H8]: check once again

Comment [H12]: remove this

Comment [H9]: rewrite the sentence




Fig 2. Lifting the prolapsed mass and relieving the retained urine



Fig 1. post-partum cervico-vaginal prolapse in buffalo

2.1 Treatment

Animal was secured in trevis for proper restraining. Fresh cool water used to clean and remove dung, blood clots and soiled material from the prolapsed mass followed by potassium permanganate solution (1:1000) to prevent further infection. Injection Tranexaleed 8ml (Tranexamic Acid 800mg) I/M given intramuscularly and epidural anaesthesia was performed by injecting 5ml of 2% lignocaine hydrochloride solution into the sacrococcygeal space. Both topical sugar application and popin spray used to reduce the edema.

Prolapsed mass was lifted in upward direction to remove retained urine (fig.2). By lubricating enough with liquid paraffin, the prolapsed mass slowly pushed into the pelvic cavity by using fist. Uterine horns, body, cervix and vagina were repositioned into their correct positions as explained by Kumbhar et al. 2009 [6] and retained by rope truss (fig.3,4). The animal was treated with Inj. D25 1000 ml (25% dextrose) I/V, Inj. RL 1000 ml I/V, Inj. Melonex 12 ml (Meloxicam, 60mg) I/M, Inj. Moxel 3gm (Amoxicillin: 2 gm & Cloxacillin: 1 gm) I/M, Inj. Dexona 5 ml (Dexamethasone sodium: 22mg) I/M, Inj. Zeet 10 ml (Chlorpheniramine maleate: 100mg) I/M.

3. DISCUSSION

Cervico-vaginal prolapse is a common reproductive disorder in the late gestation and post-partum ruminants and can be noticed by protrusion and eversion of vaginal wall and cervix through vulva [7]. There are many factors which leads to the genital prolapse through vulva. The primary cause of prolapse is thought to be the hormonal changes seen in last trimester and parturition, particularly estrogen, which makes relaxation of the pelvic ligaments and surrounding soft tissue structures [8]. Serum macro-mineral deficiency mainly calcium and phosphorus are also considered as predisposing cause of vaginal prolapse [9]. Avoiding higher intra-abdominal pressure including tympany, excessive estrogen content in the feed will decrease the occurrence of vaginal prolapse to greater extent [10].

Genital prolapses are considered as obstetrical emergencies which require early intervention [11]. Delay in treatment lead to dystocia in prepartum conditions [12]. After manual reduction, ropetruss method showed effective reoccurrence prevention [13]. Postoperative treatment for cervicovaginal prolapse can be done by allopathy (antibiotics, non-steroidal anti-inflammatory drugs), homoeopathic formulation (prolapse cure) or in their combination [14].

Comment [H13]: from title it is not matching, so remove

Comment [H14]: cervix and vagina was repositioned as explained by Kumbhar et al 2009 and retained by rope truss

Comment [H15]: treatment may lead to

Comment [H16]: retention method



Fig. 3. Rope truss applied – Lateral view.




Fig. 4. Rope truss applied – rare view.

3. CONCLUSION

Animals with cervico-vaginal prolapse when immediately corrected and properly treated may conceive successfully and deliver live calf in future.

References

1. Department of animal husbandry and dairying. Annual report 2023-2024.2024;6.https://dahd.nic.in/hi/document/annual_report.
2. Markandeya NM. Postpartum complications in buffaloes-A review. *IntasPolivet*. 2014;15(2):376-92.
3. Kumar S, Kumar B, Biswas N, Warghat C, Ghosh SK. Clinical management of pre-partum cervico-vaginal prolapse in an advanced pregnant Murrah buffalo. *Int J Vet Sci Anim Husb*. 2024; 9(5): 89-91.

Comment [H17]: 4. Conclusion, animal with cervico vaginal prolapse, if corrected and properly treated may conceive

Comment [H18]: check the reference style as per journal

4. Roberts SJ. Veterinary obstetrics and genital diseases. 2nd ed. CBS Publishers & Distributors Pvt. Limited; 2004.
5. Sah SK, Nakao T. Some characteristics of vaginal prolapse in Nepali buffaloes. *J Vet Med Sci.* 2003;65(11):1213-1215.
6. Kumbhar UB, Suryawanshi AA, Mulani JB, Raghuwanshi DS. Clinical management of post-partum eversion of uterus in Marathwadi buffalo. *Vet World.* 2009;2(5):202.
7. ArthurGH, England GC, Noakes DE, Parkinson TJ. *Arthur's Veterinary Reproduction and Obstetrics.* 8th ed:(theriogenology). Harcourt (India); 2001.
8. Thota C, Gangula PR, Dong YL, Yallampalli C. Changes in the expression of calcitonin receptor-like receptor, receptor activity-modifying protein (RAMP) 1, RAMP2, and RAMP3 in rat uterus during pregnancy, labor, and by steroid hormone treatments. *BiolReprod.* 2003;69(4):1432-1437.
9. Akhtar MS, Lodhi LA, Ahmad I, Qureshi ZI, Muhammad G. Serum concentrations of calcium, phosphorus and magnesium in pregnant Nili-Ravi buffaloes with or without vaginal prolapse in irrigated and rain fed areas of Punjab, Pakistan. *Pak Vet J.* 2008;28(3).
10. Kumar AS, Yasotha A. Correction and management of total uterine prolapse in a crossbred cow. *Journal of Agriculture and Veterinary Science.* 2015;8(1):14-16.
11. Akambaram S, Gupta C. Genital Prolapse and Its Management in Cows. *Periparturient Diseases of Cattle.* 2024 Aug 6:277-86.
12. Maheswari S, Ravikumar K, Sathishkumar S, Sureshkumar R, Monica G, Umamageswari J, Varma DD. Obstetrical management of dystocia due to prepartum cervico vaginal prolapse in a jersey cross-bred cow by cervicotomY—A case report. *Int J Vet Sci Anim Husb.* 2024; 9(4): 44-46
13. Selvaraju M, Prakash S, Varudharajan V, Periyannan M, Ravikumar K, Palanisamy M. Modified vulval truss technique to prevent recurrence of cervico-vaginal prolapse in water buffaloes. *Buffalo Bulletin.* 2023 Jun 30;42(2):133-41.
14. Patel MK, Parsani HR, Chandel BS. Therapeutic management of pre and post-partum vaginocervical prolapse in bovine by homoeopathic formulation "Prolapse Cure". *Int J Vet Sci Anim Husb.* 2024; SP-9(2): 402-405