

Relationship between weather parameters and agrometeorological indices with pod yield of groundnut

ABSTRACT

Experiments were conducted at [the](#) Department of Agricultural Meteorology, Anand Agricultural University, Anand, during [the](#) Summer of 2015 and 2016 with four varieties of groundnut (GG-2, GG-20, GJG-31 and TG-26) sown on three different dates to study the effect of weather parameters and agrometeorological indices on pod yield of groundnut. The results of [the](#) present investigation indicated that pod yield ~~of groundnut~~ was negatively correlated with maximum, ~~minimum~~ [minimum](#) and mean temperature during most of the phenophases except P₉ (pod maturity to physiological maturity), [where the](#) mean temperature ~~has been~~ varied between 32.7^oC to 35.1^oC during pod maturity to physiological maturity. [In our experiment, we have found that more than 33.9^oC required for proper seed development which results in higher test weight resulting in less than 33.90C is required for adequate seed development, resulting in higher test weight and](#) higher pod yield. Agrometeorological indices ~~has have a~~ significant and positive correlation with pod yield only during P₇ (pod initiation to pod development) phase.

~~Key words~~ [keywords](#): Phenological stages, Agrometeorological indices, correlation, groundnut

INTRODUCTION

Groundnut is ~~essentially a tropical plant and requires~~ [requires](#) a tropical plant and [requires](#) a long and warm growing season. It requires more than 16^oC soil temperature for germination. Low temperatures retard [the](#) germination and growth of plants and ~~lengthens~~ [lengthen](#) flowering. Temperature above 35^oC inhibits the growth of groundnut. ~~Mean~~ [The mean](#) temperature for optimum growth is 30^oC and the growth ceases at [a](#) temperature less than 15^oC (Prasad *et al.*, 2000). The productivity of the crop mainly depends upon the climatic requirement of the particular crop like groundnut which can grow in ~~the a~~ wide range of climatic conditions, but its productivity ~~is largely depends~~ [depends](#) on the prevailing weather conditions throughout the life cycle of the groundnut crop. Further, the different weather parameters affect growth and development of ~~crop~~ [crops](#) differently. Change in temperature, sunshine hours, [and](#) day length influence the phenophasic development of crop. Sowing time, seed germination, physiological and metabolic processes of plant life are controlled by temperature (Aggarwal *et al.* 2016). Therefore, it was contemplated to have an insight on the effect of different weather

parameters and ~~there by~~ agrometeorological indices for the production of groundnut crop and to identify the critical phases at which groundnut crop was most sensitive to the effect of particular weather parameters. This experiment was conducted to study the correlation ~~coefficient between of~~ pod yield ~~and with~~ weather parameters ~~as well as~~ ~~between and~~ the agrometeorological indices.

MATERIALS AND METHODS

~~Field-A field~~ experiment was conducted in ~~the~~ Department of Agricultural Meteorology, Anand Agricultural University, Anand during summer 2015 and 2016 ~~16~~ with treatments ~~consist~~ ~~consisting of~~ four varieties of groundnut viz., GG-2, GG-20, GJG-31 and TG-26 were sown on three different dates viz., D₁ early date (3rd January), D₂ normal date (15th February) and D₃ late date (2nd March). The crop was sown as per ~~the~~ recommended package of practices. ~~The phenological events parameters~~ viz: days to emergence, days to first flower opening, days to 50% flowering, days to peg initiation, days to 100 % flowering and days to pod initiation, days to pod development, days to pod maturity and days to physiological maturity were recorded by visual observation on tagged five plants per replication from each plot. The daily meteorological data during both crop seasons was recorded at ~~the Agrometeorological observatory~~ ~~observatory~~ of Anand Agricultural University, Anand. Correlation between the total pod yield ~~with various weather parameters during different phenophases~~ ~~were was and various weather parameters during different phenophases~~ was carried out using a methodology described by Gomez and Gomez (1984).

RESULTS AND DISCUSSION

Temperature

Correlation analysis showed that maximum temperature (T_{max.}) was significantly and negatively correlated with ~~the~~ P₂ (emergence to first flower initiation) phase. T_{max.} during this phase was between 30.8 and 37.2 °C. Thus the higher T_{max.} (>34°C) during early vegetative period was not suitable for groundnut. The significant positive correlation was observed between pod yield and T_{max.} during maturity period P₉ (pod maturity to physiological maturity) phase, during this period T_{max.} varied between 38.7 and 42.8°C, pod maturity requires T_{max.} more

Comment [P1]: Mention all the phases in materials and methods

than 40.7 °C for proper seed development which resulted in higher test weight resulting in higher pod yield of groundnut.

The pod yield was negatively correlated with minimum temperature (Tmin.) during most of the phenophases except during the maturity period. Higher night temperature Tmin. during P₂ and P₆ phase adversely affected the flowering, which ultimately contributed to lower pod yield. Similarly the pod yield was negatively correlated with mean temperature (Tmean.) during most of the phenophases except during maturity period. However, the insignificant negative correlation was obtained with Tmean during P₂ (emergence to first flower initiation) phase. Tmean during this was between 22.9 and 28.6 °C. Similarly, significant and negative correlation was obtained with Tmean during P₆ (100% flowering to pod initiation) phase. Tmean during P₆ between 23.7 and 30.3 °C. Thus, the higher Tmean. were (>25.7 °C) and (>27 °C) during P₂ and P₆ phase was not suitable for groundnut.

Bright sunshine hours (BSS)

The pod yield was negatively correlated with bright sunshine hours (BSS) during most of the phenophases except during P₃, P₄, P₇ and P₉ phases. However, the insignificant negative correlation was obtained with BSS during P₆ (100% flowering to pod initiation) phase. BSS during this phase were between 8.6 and 10.7 hrs. Similarly significant and negative correlation was obtained with BSS during P₈ (pod development to pod maturity) phase. BSS during P₈ between 9.8 and 11.0 hr. Thus, the higher BSS varied between >9.6 hrs. and >10.4 hrs. 10.4 hrs. during P₆ and P₈ phase was not suitable for groundnut. Higher BSS during P₆ and P₈ phase adversely affected the pod yield.

Relative humidity

The significant and positive correlation between pod yield and afternoon relative humidity RH₂ during P₁(sowing to emergence) phase suggest that during this period RH₂ varied between 28.0 and 48.8 per cent. Afternoon relative humidity requires more than 38.4 per cent for a higher pod yield of groundnut. However, the a significant negative correlation was obtained with RH₂ during P₃ (first flower initiation to 50 % flowering) phase. RH₂ during this was between 17.0 and 41.67 %. Similarly, a significant and negative correlation was obtained with T_{mean} during P₅(peg initiation to 100 % flowering) and P₉ (pod maturity to physiological maturity) phases. RH₂ during P₅ between 12 and 68 % and P₉ between 27 and 53%. Thus, the higher RH₂ were was (>27.8%), (>40%) and (>40%) during P₃, P₅ and P₉ phases was not suitable for groundnut.

Vapour Pressure

The results induced showed that the pod yield was negatively correlated with morning vapour pressure VP₁ during all the phenophases. In case in case afternoon vapour pressure VP₂ during P₁(sowing to emergence) phase was had a positive correlation with pod yield, Pod yield was also negatively correlated with mean vapour pressure VP_{mean} during all the phenophases.

Agrometeorological Indices

The significant and positive correlation between pod yield and growing degree days GDD °C day only during P₇(pod initiation to pod development) phase suggest that during this period GDD varied between 285 and 411 °C day. GDD requires more than 348 °C day for higher pod yield of groundnut during pod initiation to pod development phase. Similarly, the significant and positive correlation between pod yield and heliothermal units HTU during P₃ (first flower initiation to 50% flowering) and P₇(pod initiation to pod development) phase suggest that during this period HTU varied between (382 and 922 °C day hrs) and (2577 and 4175 °C day hrs). HTU

requires more than 652⁰C day hrs. and 3376⁰C day hrs. for higher pod yield of groundnut during P₃ and P₇ phase. Also, the significant and positive correlation between pod yield and photothermal units PTU during P₇ (pod initiation to pod development) phase suggest that during this period PTU varied between 3655 and 5196 ⁰C day hrs. PTU requires more than 4425⁰C day hrs. for higher pod yield of groundnut during pod initiation to pod development phase. Meena and Dahama (2004) also, reported that the pod yield was positively correlated with heliothermal units HTU and photothermal units PTU from flower initiation to maturity.

CONCLUSION

There was a good correlation with Agrometeorological indices like GDD, HTU and PTU was ~~positive~~ positively correlated with pod yield during pod initiation to pod development and negatively correlated with maximum, minimum temperature, mean temperature, BSS during most of the phenophases except during maturity period.

REFERENCES

- Aggarwal, N., Singh, A. and Singh, S. P. (2016) Heat utilization and radiation interception in transplanted rice (*Oryza sativa* L.) in relation to seedling age. *Journal of Agrometeorology*, 18(1): 93-96
- Gomez and Gomez (1984). Statistical procedures for agricultural research. 2nd Ed. An International Rice Research Institute. John Willey and sons, New York.
- Prasad, P. V. V., Craufurd, P. Q. and Summerfield, R. J. (2000). Effect of high air and soil temperature on dry matter production, pod yield and yield components of groundnut. *Plant and Soil*, 222: 231-239.

Parameters	P1	P2	P3	P4	P5	P6	P7	P8	P9
Tmax.	-0.20	-0.50*	-0.21	0.15	-0.32	-0.36	-0.40	-0.2	0.54**
Tmin.	-0.29	-0.48*	0.04	-0.38	-0.38	-0.50*	-0.29	-0.35	0.23
Tmean	-0.26	-0.49*	-0.08	-0.11	-0.28	-0.52**	-0.37	-0.32	0.53**

Table 1: Correlation coefficients between ~~phase-wise~~ phase-wise weather parameters, agrometeorological indices and pod yield of groundnut

Comment [P2]: Please add the table title at top of the table

BSS	-0.27	-0.11	0.19	0.25	-0.32	-0.44*	0.08	-0.41*	0.51**
RH1	-0.05	0.16	-0.08	-0.09	-0.26	0.11	-0.31	-0.07	-0.16
RH2	0.53**	0.28	-0.42*	-0.26	-0.48*	-0.00	-0.17	-0.29	-0.49*
Rhmean	0.27	0.22	-0.17	-0.20	-0.39	0.03	-0.27	-0.24	-0.40*
VP1	-0.35	-0.59**	-0.31	-0.35	-0.46*	-0.30	-0.52**	-0.32	-0.12
VP2	0.41*	-0.12	-0.47*	-0.28	-0.42*	-0.18	-0.43*	-0.27	-0.42*
Vpmean	-0.19	-0.42*	-0.40*	-0.32	-0.44*	-0.24	-0.50*	-0.30	-0.31
Rainfall	.a	0.20	.a	0.14	-0.28	-0.17	-0.14	0.32	-0.41*
GDD	0.33	-0.04	0.33	0.30	-0.11	0.27	0.54**	0.07	-0.14
HTU	0.28	-0.22	0.42*	0.11	-0.17	-0.16	0.52**	0.06	-0.10
PTU	0.33	-0.07	0.29	0.23	-0.15	0.22	0.52**	0.05	-0.15

Where P₁ (from sowing to emergence), P₂ (from emergence to first flower opening), P₃ (from first flower opening to 50 % flowering), P₄ (from 50 % flowering to peg initiation), P₅ (from peg initiation to 100 % flowering), P₆ (from 100 % flowering to pod initiation), P₇ (from pod initiation to pod development), P₈ (from pod development to pod maturity), P₉ (from pod maturity to physiological maturity) phase