

Case report

Poisoning with local anesthetics following circumcision :a case report

Abstract

Introduction: Circumcision or posthectomy is the removal of the foreskin, this act seems commonplace practiced by paramedical personnel or unqualified traditional practitioners who may cause significant complications.

Objective: Our work aimed to highlight one of the complications of this act which must remain a supervised medical act.

Observation: We report the case of a 2-month-old infant who presented with complications related to anesthesia. the clinic was dominated by convulsions and alteration of the state of consciousness 15 minutes after circumcision in this well infant carrying without any particular history. The evolution was favorable thanks to support rapid and urgent about 20 minutes with the cessation of convulsions and a return to consciousness, the patient was monitored for 48 hours then return home was indicated due to the absence other complications.

Conclusion: Circumcision is a widespread act throughout the world either by medical necessity or by religious tradition. its trivialization and its practice by non-experts, explains the multiple complications observed.

Keywords: Circumcision-Poisoning - Xylocaine-Convulsion-Infant-Libreville

1. INTRODUCTION

Circumcision is the surgical removal of the foreskin, also called posthectomy. The intervention proposed to the boy consists of partially or completely removing the foreskin for medical reasons or not. This intervention is better known under the term circumcision, it concerns 30% of the world male population [1-3]. Ritual practice imposed by mosaic law for the israelites, religious rite for muslims, and rite of transition to the adult state for several peoples of africa and the orient [4-7]. However, this act minor surgery is sometimes carried out by paramedical and traditional personnel [5,6]. Patrician in our context. The latter ignore the notions of anatomy, aseptic and non knowledge of dosage related to anesthesia. This lack of knowledge can be harmful for children and affect the sexual and urinary functional prognosis, even the vital prognosis of the child [8-12]. The

fight against pain in a hospital environment is imperative both in terms of physiological and human, adequate anesthesia requires an initial assessment of pain and its mechanisms, appropriate monitoring and reassessment. A knowledge of possible side effects and ability to manage them. Anesthesia local core regional is a technique of choice since it provides level 3 analgesia. Pediatric anesthesia, this principle is abandoned and the vast majority of peri-nerve blocks are performed under general anesthesia. The aim of this work is to highlight one of the complications of this act which must remain a supervised medical act.

2. CASE PRESENTATION

The infant G.J, male, aged 2 months, was hospitalized in the department of pediatric emergencies of the CHUME FJE on 08/21/21 for investigation and the care of a medical condition convulsive evolving in a non-febrile context. The patient had consulted for a convulsion occurring immediately in the context of the use of lidocaine during a circumcision. The practitioner would have used a whole bottle of 2% xylocaine for anesthetic purposes, i.e. 20 ml intramuscularly or Lidocaine is commonly injected around the root of the penis. This infant has no particular medical-surgical history, circumcision was scheduled and carried out at the request of parents. The clinical examination on admission found a Weight: 7100 g, a temperature 36.6 °C a heart rate at 160 bpm, a syndrome of cortical irritation with convulsions not stopping despite 3 doses of diazepam at 0.5 mg/kg at 5 min intervals, a syndrome of deterioration of general condition with a score of Blantyre rated 3/5, reactive isocore pupils, blood sugar level of 5.6 mmol/l and SaPO₂ 94% ambient air. The problem posed was the etiology of a convulsive illness in a infant in apparent good health and having received a high dose of xylocaine injection. We discussed metabolic causes that we eliminated in the face of an assessment. normal metabolic rate (blood ionogram, serum calcium and blood sugar). We therefore retained xylocaine poisoning. The treatment consisted of clearing the airways superior, oxygen therapy at 3 L/min with glasses, hydration at 80ml/kg/d. The evolution was favorable with the improvement of the convulsions after 15 min, the return progressive to normal consciousness after 4 hours, discharge was decided after 48 hours of monitoring with normal EEG.

Discussion

We reported a case of post-circumcision intoxication. Circumcision is an act responded around the world, in 1996, during the 4th international conference organized in Lausanne, it was estimated that 13,300,000 male children were circumcised each year [13]. The incidence of complications varies depending on the studies, whether or not they were carried out in a healthcare establishment, operator qualification and duration of follow-up. During studies controlled, randomized studies aimed at evaluating the protective effect of circumcision medically framed on HIV infection, carried out in Africa, the authors reported complications in 1.5 to 3.8% of cases [14]. The age of our patient is 2 months, this age is found in other studies which encourage circumcision in the neonatal period because they highlight its prophylactic effect against STIs, urinary infections as well as certain cancers of the genital tract [15,16]. According to WHO, circumcision should be

practiced exclusively by competent health professionals and should be carried out in regulatory hygiene conditions, conditions necessary to reduce complications [17,18]. They vary in frequency and severity and each requires an attitude particular therapy. Some are seen early, anesthetic accidents as found in the literature and in our patient. In our case it was carried out by unqualified medical personnel, thus increasing the risk of complications. Our patient presented convulsions 15 minutes after the circumcision and after use of xylocaine, this confirms the literature data on neurological toxicity of local anesthetics at high concentrations. The margin of safety is very low, and anesthesiologists are very attentive to the appearance of signs announcers. Systemic neurological toxicity is by far the most common of toxic manifestations with the frequency of a convulsive accident per 800 to 1,000 loco-regional anesthesia [19]. Xylocaine 20 mg/ml without preservative is a solution Lidocaine Hydrochloride injection, it is a local anesthetic used to anesthetize part of the body during surgical operations and also to relieve pain. There maximum recommended dose is between 2 and 7 mg/kg Local infiltration and anesthesia regional. The occurrence of an adverse effect should raise suspicion of an overdose. The reactions toxic, indicative of an overdose of local anesthetic, can appear in two conditions: either immediately by a relative overdose due to intravenous passage accidental, or later by a true overdose due to the use of too much amount of anesthetic. Signs of toxicity can be; in terms of the nervous system central sensations of dizziness, numbness of the lips and tongue, ringing in the ears, hearing dysfunction and dysarthria, disorientation and occasional feelings of drowsiness. Objective signs of toxicity, include shivering, muscle contractions and tremors initially involving the muscles of the face and distal parts of the extremities and ultimately convulsions appear. If a very large dose is administered, the first signs are monitored quickly by a state of generalized depression of the central nervous system which can lead to respiratory arrest [20]. In children from approximately 10 days to 4-6 months, lidocaine 1%, at a maximum dose of 7 to 10 mg/kg provides short or medium duration anesthesia [21], in our case the patient received a dose of 16 mg/kg causing convulsions after 15 minutes. The poisonings appear when the so-called toxic plasma thresholds are exceeded following an accidental injection or an overdose due to a miscalculation. This toxicity manifests itself first by neurological signs with prodromes which precede the convulsive attack. Several experimental works have shown that the mechanisms involved are multiple. with dysfunction of calcium and mitochondrial metabolism [22].

4. CONCLUSION

Circumcision is a widespread act throughout the world either by medical necessity or by religious tradition. Indeed, its trivialization and its practice by non-experts, explain the multiple complications observed as described in our case, thus being able to put into play the functional, aesthetic and sometimes vital prognosis of patients. It is essential that this act be supervised in order to minimize the frequency of complications.

Disclaimer (Artificial intelligence)

We hereby declare that NO generative AI technologies such as Large Language models (ChatGPT, COPILOT, etc.) and image text generators have been used when writing or editing this manuscript

Ethical Approval:

As per international standards or university standards written ethical approval has been collected and preserved by the author(s).

Consent

As per international standards, parental written consent has been collected and preserved by the author(s).

REFERENCES

1. Beydon L, Ecoffey C, Lienhart A, Puybasset L. Circumcision in children: a simple question of organization of care or an ethical issue? *Ann Fr Anesth Reanim* (2012) <https://doi.org/10.1016/j.annfar.2012.01.038>.
2. Tambo, F. M., Fadi, S., Ditope, J. J., Nyanit, B. D., Mbouche, L., & Andze, O. G. (2014). Aspects anatomocliniques et thérapeutiques des accidents de la circoncision à l'hôpital Gynéco-obstétrique et pédiatrique de Yaoundé (HGOPY): à propos de 15 cas. *Revue Africaine de Chirurgie et Spécialités*, 8(3), 15-18.
3. Tambo FF. Les accidents de la circoncision à Yaoundé, Cameroun : à propos de cinq observations cliniques [Circumcision accidents in Yaoundé, Cameroon: a report on five cases]. *Prog Urol*. 2012 Jan ;22(1):63-6. French. doi: 10.1016/j.purol.2011.06.003. Epub 2011 Jul 26. PMID: 22196008.
4. Circumcision. <http://fr.wikipedia.org/wiki/Circumcision>. Accessed 5-5-2011.
5. Hassan Y, Rasool H, Rather AA, Ahmad Y, Rasool I. Religious circumcision (Khatna) and circumcision mishaps in Kashmiri children. *Afr J Paediatr Surg*. 2022 Oct- Dec;19(4):213-216. doi: 10.4103/ajps.ajps_158_21. PMID: 36018200; PMCID: PMC9615959
6. Appiah KA, Gyasi-Sarpong CK, Azorliade R, Aboah K, Laryea DO, Otu-Boateng K, Baah-Nyamekye K, Maison PO, Arthur D, Antwi IO, Frimpong-Twumasi B, Yenli EM, Togbe SK, Amoah G. Circumcision-related tragedies seen in children at the Komfo Anokye Teaching Hospital, Kumasi, Ghana. *BMC Urol*. 2016 Nov 8;16(1):65. doi: 10.1186/s12894-016-0183-1. PMID: 27825332; PMCID: PMC5101822.
7. Ouattara A, Paré AK, Yé D, Sherazi A, Simporé M, Rouamba M, Kaboré AF, Kambou T. Complications of non-medical assisted circumcision in Burkina Faso. Clinical presentation, management, and outcomes - about 23 cases and literature review. *Arch Ital Urol Androl*. 2023 Jul 25;95(3):11494. doi: 10.4081/aiua.2023.11494. PMID: 37491935

8. Ghidini F, Virgone C, Madoukeng BM, Franchella A, Vason M, Cumba D, Tognon C, Gamba P. Perioperative management of circumcision in children: Is there a difference between African and European hospitals? *Afr J Paediatr Surg.* 2020 Jul-Dec;17(3 & 4):90-94. doi: 10.4103/ajps.AJPS_24_20. PMID: 33342841; PMCID: PMC8051633
9. Shah A, Shah A. Total Glans Amputation in a Child following Ritual Circumcision - Report of a Case and Review of Literature. *J Indian Assoc Pediatr Surg.* 2023 May-Jun;28(3):256-259. doi: 10.4103/jiaps.jiaps_176_22. Epub 2023 Mar 21. PMID: 37389392; PMCID: PMC10305961
10. Soltani S, Mottaghi M, Jafarpisheh A, Tavakkoli M. Penile Glans Amputation following Circumcision: A Case Report of a Rare Complication. *Case Rep Urol.* 2020 Oct 2;2020:5806987. doi: 10.1155/2020/5806987. PMID: 33083087; PMCID: PMC7556101.
11. Nafatalewa DK, Mukakala AK, Mujinga IMW, Lubosha NA, Yumba SN, Banza MI, Kaoma VP, Katambwa PM, Musapudi EM, Misenga JB, Mukaz PM. Amputation totale du gland lors de la circoncision en milieu non hospitalier : rapport de cas [Penile glans amputation during non-hospital circumcision : about two cases]. *Pan Afr Med J.* 2022 Jul 19;42:214. French. Doi : 10.11604/pamj.2022.42.214.28325. PMID: 36258900; PMCID: PMC9569146.
12. Matiben Tayed. of circumcision in Algeria, thesis for the doctorate in medicine presented and publicly defended on December 16, [1950](#). July 21, [2013](#)
13. Circumcision in ancient Egypt: International Network November 25, 2017
14. <https://www.laprocare.com/product/29263/guggenheim-antoine-la-circoncision-circumcision-of-the-flesh-and-circumcision-of-the-heart-a-reading-from-saint-paul?srsltid=AfmBOorQx0FYp29pX-mpa5Jx2Y4kgsQnZe0qsShU0ahG1JGTrOC0HgTR> 2014.
15. https://www.medecinesciences.org/en/articles/medsci/full_html/2008/07/
16. Bruno Halioua, Bernard Lobel. Current controversies over circumcision. *Med Press*; 11:1171 (2014).
17. Christian G Guevara, Justin K Achua Ruben Blachman-Braun Isabella Cabrera-Valencia, George A Ransford Rafael Gosalbez. Neonatal Circumcision: What Are the Factors Affecting Parental Decision? (11): 9-13 (2021).
18. Laurence S Baskin, MD, FAAP. Charles J Lockwood, MD, MHCM. Duncan Wilcox, MD. Kristen Eckler, MD, FACOG. Neonatal circumcision: Risks and benefits (2023).
19. Male circumcision global trends and determinants of prevalence, safety and acceptability. WHO document (2009).

20. Kariem El-Boghdadly, Amit Pawa, et Ki Jinn Chin. Anesthésique local systémique toxicité : perspectives actuelles Publié en ligne - local Reg Anesth. 11 : 35-44 (2018)

21. <https://base-donnees-publique.medicaments.gouv.fr/affichageDoc.php?specifie=61751079&typedoc=N>. ANSM. Mise à jour le : 23/06/2021.

22. Dadure C., Marec P., Veyckemans F., Belœil H. Douleurs chroniques et anesthésie locorégionale chez l'enfant. Archives de Pédiatrie Vol20, Numéro 10, 1149-1157 (2013).

UNDER PEER REVIEW