

Synergistic Effect of Organic Inputs on the Morphophysiological Traits of Sweet Basil (*Ocimum basilicum* L.)

ABSTRACT

The present investigation on "Synergistic Effect of Organic Inputs on the Morphophysiological Traits of Sweet basil (*Ocimum basilicum* L.)" was conducted from 2023 in Chengam, Tiruvannamalai district. The objective was to evaluate the impact of various organic treatments on the growth and yield characteristics of sweet basil (*Ocimum basilicum* L.). The experiment was carried out in Randomized Block Design with ten treatments and replication thrice. Among the ten different treatments applied, the combination of FYM @ 25 t ha⁻¹ + VC @ 2.5 t ha⁻¹ + Seaweed extract @ 2% (T₉) consistently outperformed other treatments. This treatment significantly enhanced growth characteristics viz., plant height (35.01, 60.33, 86.71 and 110.34 cm), number of leaves plant⁻¹ (291.34, 540.56, 850.41 and 1010.66), leaf area (482.33, 1673.20, 3581.23 and 4468.67 cm²) and leaf area index (0.65, 1.33, 2.32 and 3.23), number of primary branches plant⁻¹ (21.45, 25.37, 32.17 and 39.46) and secondary branches plant⁻¹ (64.72, 76.45, 96.56 and 109.34), plant spread (north - south) (25.32, 33.39, 40.26 and 50.38 cm), plant spread (East- west) (40.56, 55.89, 65.47 and 71.45 cm), at 30, 60, 90 and 120 DAT. The findings indicate that the integration of organic inputs of treatment, T₉ (FYM @ 25 t ha⁻¹ + VC @ 2.5 t ha⁻¹ + Seaweed extract @ 2%) is highly effective in optimizing the growth of sweet basil.

Key Words : Sweet basil, Organic farming, FYM, Vermicompost, Seaweed extract.

INTRODUCTION

Sweet Basil (*Ocimum basilicum* L.), a widely cultivated aromatic herb, is known for its flavorful foliage (Davis, 1995). It belongs to the Lamiaceae family, comprising 50-150 species of herbs and shrubs native to tropical regions of Asia, Africa and the Americas (Malav *et al.*, 2015). Sweet basil is tetraploid, with a chromosome number of 2n=48 (Farooqi & Sreeramu, 2005). Known as the "King of herbs," basil holds commercial, culinary, cosmetic and medicinal value (Meena *et al.*, 2013). Overuse of chemical fertilizers depletes soil, while organic fertilizers restore nutrients and enhance soil health. Organic farming of sweet basil boosts pharmaceutical demand for its chemical-free properties, improving oil quality and medicinal value. Organic systems enhance morpho-physiological traits, yield and soil health by increasing organic carbon, phosphorus and microbial activity, ensuring long-term productivity. The overuse of chemical fertilizers has degraded soil health and reduced the quality of medicinal and aromatic plants (MAPs), which are increasingly needed for chemical-free products in the pharmaceutical industry. Organic farming offers a solution by restoring soil fertility, improving crop growth and enhancing essential oil quality. This sustainable approach boosts yields, enhances soil health and meets the demand for high-quality, organically grown MAPs like sweet basil.

Comment [DT1]: Make botanical name italics

Comment [DT2]: Clarify the leaf area whether it is complete area covered by total leaves or it is average leaf area

Comment [DT3]: Add conclusion

Comment [DT4]: Make introduction upto date and more organized

MATERIAL AND METHODS

The present investigation on “Synergistic Effect of Organic Inputs on the Morphophysiological Traits of Sweet basil (*Ocimum basilicum* L.)” was conducted from 2023 in Chengam, Tiruvannamalai district. The experiment was laid out in Randomized Block Design with ten treatments and three replications. Each replication has 30 plants. Five randomly selected plants in each replication were tagged, labelled and used for observation of different growth parameters. The mean of five plants was taken for analysis. The data recorded were subjected to statistical analysis by adopting the standard procedure of Panse and Sukhatme (1985). The critical differences were arrived at 5 per cent probability significance.

Comment [DT5]: Mention the period of experiment

RESULT AND DISCUSSION

The data in Table 1 showed significant differences among treatments, with T₉ (FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Seaweed extract @ 2%) recording the highest plant height (35.01, 60.33, 86.71, 110.34 cm) and number of leaves plant⁻¹ (291.34, 540.56, 850.41, 1010.66) followed by T₁₀ (FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Jeevamrutham @ 3%) with plant height of 33.74, 58.05, 83.77, 106.22 cm and number of leaves plant⁻¹ of 269.69, 510.22, 821.64, 969.12 while the lowest values were observed in the control (T₁) with plant heights of 23.62, 39.81, 60.34 and 73.62 cm and number of leaves plant⁻¹ of 96.49, 267.50, 591.48 and 636.80 at 30, 60, 90 and 120 DAT.

Comment [DT6]: Mention the statistical software and methods used for the analysis of data

Farmyard manure (FYM) promotes microbial growth, accumulates humus and provides phytohormones that aid plant growth, even with reduced chemical fertilizers (Gupta *et al.*, 1983). The basal application of vermicompost (VC) enhances plant height by promoting cell division and enlargement. The combination of organic manures like FYM and VC ensures balanced nutrient supply, increasing the number of nodes and internodal length, thus boosting plant height. Similar results were observed by Malav *et al.* (2015), Baraa *et al.* (2017), Jalil Dehghan Saman *et al.* (2017) and Rajit Ram *et al.* (2019a) in *Ocimum spp.*

Table 1. Effect of organic inputs on plant height (cm) and Number of leaves plant⁻¹ in sweet basil(*Ocimum basilicum* L.)

Treatments	Plant height (cm)				Number of leaves plant ⁻¹			
	30DAT	60 DAT	90 DAT	120 DAT	30DAT	60DAT	90DAT	120DAT
T ₁ - Control	23.62	39.81	60.34	73.62	96.49	267.50	591.48	636.80
T ₂ - FYM @ 25t ha ⁻¹	24.88	42.09	63.26	77.38	118.14	297.83	620.25	678.34
T ₃ - VC @ 5t ha ⁻¹	26.15	44.37	66.20	81.50	139.09	328.18	649.02	719.88
T ₄ - FYM @ 25t ha ⁻¹ + VC @ 2.5t ha ⁻¹	29.59	51.21	74.99	93.86	204.74	419.20	735.32	844.50
T ₅ - FYM @ 25 t ha ⁻¹ + Seaweed extract @2%	28.68	48.93	72.06	89.74	183.09	388.86	706.56	802.96
T ₆ - FYM @ 25 t ha ⁻¹ + Jeevamrutham @ 3%	27.42	46.65	69.13	85.62	161.44	358.51	677.79	761.42
T ₇ - VC @ 5 t ha ⁻¹ + Seaweed extract @2%	32.48	55.77	80.85	102.10	248.04	479.88	792.87	927.58
T ₈ - VC @ 5 t ha ⁻¹ + Jeevamrutham @ 3%	31.21	53.49	77.92	97.98	226.39	449.54	764.10	886.04
T ₉ - FYM @ 25 t ha ⁻¹ + VC @ 2.5 t ha ⁻¹ + Seaweed extract @2%	35.01	60.33	86.71	110.34	291.34	540.56	850.41	1010.66
T ₁₀ - FYM @ 25 t ha ⁻¹ + VC @ 2.5 t ha ⁻¹ + Jeevamrutham @ 3%	33.74	58.05	83.77	106.22	269.69	510.22	821.64	969.12
S.ED	0.57	0.98	1.43	1.80	4.13	8.19	14.1	16.2
CD (P = 0.05)	1.15	1.97	2.89	3.62	8.38	16.47	28.36	32.67

Table 2. Effect of organic inputs on leaf area (cm²) and leaf area index in sweet basil(*Ocimum basilicum* L.)

Treatments	Leaf area (cm ²)				Leaf area index			
	30 DAT	60 DAT	90 DAT	120 DAT	30 DAT	60 DAT	90 DAT	120 DAT
T ₁ - Control	331.85	967.69	2026.57	2447.81	0.38	0.79	1.33	2.15
T ₂ - FYM @ 25t ha ⁻¹	348.57	1046.08	2199.31	2672.35	0.41	0.84	1.44	2.26
T ₃ - VC @ 5t ha ⁻¹	365.29	1124.46	2372.05	2896.89	0.43	0.91	1.55	2.38
T ₄ - FYM @ 25t ha ⁻¹ + VC @ 2.5t ha ⁻¹	415.44	1359.64	2890.26	3570.51	0.53	1.09	1.88	2.75
T ₅ - FYM @ 25 t ha ⁻¹ + Seaweed extract@2%	398.73	1281.25	2717.53	3345.97	0.50	1.03	1.77	2.63
T ₆ - FYM @ 25 t ha ⁻¹ + Jeevamrutham @ 3%	382.01	1202.86	2544.79	3121.43	0.47	0.97	1.66	2.50
T ₇ - VC @ 5 t ha ⁻¹ + Seaweed extract @2%	448.89	1516.42	3235.75	4019.59	0.59	1.21	2.10	2.99
T ₈ - VC @ 5 t ha ⁻¹ + Jeevamrutham @ 3%	432.17	1438.03	3063.01	3795.05	0.56	1.15	1.99	2.87
T ₉ - FYM @ 25 t ha ⁻¹ + VC @ 2.5 t ha ⁻¹ + Seaweed extract @2%	482.33	1673.20	3581.23	4468.67	0.65	1.33	2.32	3.23
T ₁₀ - FYM @ 25 t ha ⁻¹ + VC @ 2.5 t ha ⁻¹ + Jeevamrutham @ 3%	465.61	1597.80	3408.49	4244.13	0.62	1.27	2.21	3.11
S.ED	7.97	26.2	55.9	69.2	0.01	0.20	0.36	0.52
CD (P = 0.05)	16.02	502.82	112.44	139.20	0.021	0.042	0.074	0.106

Note: FYM-farm yard manure, VC- vermicompost

Table 3. Effect of organic inputs on plant spread North – South (cm) and Plant spread East - West (cm) in sweet basil (*Ocimum basilicum* L.)

Treatments	plant spread North – South (cm)				Plant spread East - West(cm)			
	30 DAT	60 DAT	90 DAT	120 DAT	30 DAT	60 DAT	90 DAT	120 DAT
T ₁ – Control	16.68	23.13	24.33	35.16	26.79	32.49	39.37	46.07
T ₂ - FYM @ 25t ha ⁻¹	17.64	24.26	26.10	36.85	28.32	35.09	42.26	48.88
T ₃ - VC @ 5t ha ⁻¹	18.60	25.40	27.87	38.55	29.85	37.69	45.17	51.71
T ₄ -FYM @ 25t ha ⁻¹ + VC @ 2.5t ha ⁻¹	21.47	28.83	33.18	43.62	34.43	45.48	53.87	60.17
T ₅ - FYM @ 25 t ha ⁻¹ + Seaweed extract@2%	20.50	27.69	31.41	41.93	32.91	42.89	50.97	57.35
T ₆ - FYM @ 25 t ha ⁻¹ + Jeevamrutham @ 3%	19.55	26.54	29.64	40.24	31.38	40.29	48.07	54.52
T ₇ - VC @ 5t ha ⁻¹ + Seaweed extract @2%	23.40	31.11	36.72	47.00	37.50	50.68	59.66	65.81
T ₈ - VC @ 5t ha ⁻¹ + Jeevamrutham @ 3%	22.44	29.97	34.95	45.30	35.97	48.09	56.77	62.99
T ₉ - FYM @ 25 t ha ⁻¹ + VC @ 2.5 t ha ⁻¹ + Seaweed extract @2%	25.32	33.39	40.26	50.38	40.56	55.89	65.47	71.45
T ₁₀ - FYM @ 25 t ha ⁻¹ + VC @ 2.5 t ha ⁻¹ + Jeevamrutham @ 3%	24.36	32.25	38.49	48.69	39.03	53.29	62.57	68.63
S.ED	0.40	0.55	0.63	0.83	0.66	0.87	1.03	1.15
CD (P = 0.05)	0.82	1.11	1.28	1.68	1.33	1.76	2.08	2.32

Table 4. Effect of organic inputs on number of primary and secondary branches plant⁻¹ in sweet basil (*Ocimum basilicum* L.)

Treatments	Number of primary branches plant ⁻¹				Number of secondary branches plant ⁻¹			
	30 DAT	60 DAT	90 DAT	120 DAT	30 DAT	60 DAT	90 DAT	120 DAT
T ₁ – Control	12.27	14.48	18.76	23.89	27.46	40.27	45.89	58.58
T ₂ - FYM @ 25t ha ⁻¹	13.28	15.69	20.25	25.62	31.60	44.28	51.52	64.22
T ₃ - VC @ 5t ha ⁻¹	14.31	16.90	21.74	27.35	35.74	48.31	57.14	69.86
T ₄ -FYM @ 25t ha ⁻¹ + VC @ 2.5t ha ⁻¹	17.36	20.53	26.21	32.54	48.16	60.36	74.04	86.78
T ₅ - FYM @ 25 t ha ⁻¹ + Seaweed extract @2%	16.35	19.32	24.72	30.81	44.02	56.35	68.41	81.14
T ₆ - FYM @ 25 t ha ⁻¹ + Jeevamrutham @ 3%	15.33	18.11	23.23	29.08	39.88	52.33	62.78	75.50
T ₇ - VC @ 5t ha ⁻¹ + Seaweed extract @2%	19.41	22.95	29.19	36.00	56.44	68.41	85.30	98.06
T ₈ - VC @ 5t ha ⁻¹ + Jeevamrutham @ 3%	18.39	21.74	27.70	34.27	52.30	64.39	79.67	92.42
T ₉ - FYM @ 25 t ha ⁻¹ + VC @ 2.5 t ha ⁻¹ + Seaweed extract @2%	21.45	25.37	32.17	39.46	64.72	76.45	96.56	109.34
T ₁₀ - FYM @ 25 t ha ⁻¹ + VC @ 2.5 t ha ⁻¹ + Jeevamrutham @ 3%	20.43	24.16	30.68	37.73	60.58	72.43	90.93	103.70
S.ED	0.33	0.39	0.50	0.62	0.95	1.17	1.42	1.68
CD (P = 0.05)	0.67	0.80	1.01	1.26	1.92	2.36	2.92	3.38

Note: FYM–farm yard manure, VC- vermicompost

Seaweed extracts improve plant growth due to growth-promoting hormones (IAA, IBA, cytokinins), trace elements (Fe, Cu, Zn, Co, Mo, Mn, Ni), vitamins and amino acids, as noted by Katarzyna Chojnacka *et al.* (2012), Wajahatullah Khan *et al.* (2009) and Craige (2011). Organic manures enhance leaf production due to the optimal C:N ratio in FYM, which releases nitrogen (ammonium and nitrate) during decomposition. Increased nitrogen levels stimulate leaf production, larger leaf areas and greater plant spread, as nitrogen is vital for amino acids and co-enzymes (Venkatesan, 2009). Similar findings were reported by Ashashri Shinde *et al.* (2013) in ashwagandha, Mansour *et al.* (2017) in sweet basil, Patke *et al.* (2018), Ram *et al.* (2019b) in Indian basil (*Ocimum sanctum* L. cvs. Cim-Ayu and Cim-Angana), Aloe vera and Gunda *et al.* (2022) in sweet basil. Nitrogen in vermicompost also boosts leaf production, supported by findings from Suresh and Senthilnathan (2018). This is consistent with studies by Padmapriya *et al.* (2010) in gymnema, Tiwari and Roy (2014) in gloriosa, Khanzadeh and Naderi (2015) in periwinkle, Hidangmayum and Sharma *et al.* (2017) and Suresh *et al.* (2018) in Japanese mint. Seaweed extracts, containing growth substances, further enhance vegetative growth and yield by influencing cellular metabolism (El-Miniawy *et al.*, 2014).

Comment [DT7]: If possible, use latest references

The data in Table 2 revealed significant differences among treatments in leaf area (cm²) and leaf area index. T₉ (FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Seaweed extract @ 2%) recorded the highest leaf area (482.33, 1673.20, 3581.23 and 4468.67 cm²) and leaf area index (0.65, 1.33, 2.32 and 3.23). This was followed by T₁₀ (FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Jeevamrutham @ 3%) with leaf area of 456.61, 1597.80, 3408.49 and 4244.13 cm² and leaf area index of 0.62, 1.27, 2.21 and 3.11. The lowest values were recorded in T₁ (Control), with leaf area of 331.85, 967.69, 2026.57 and 2447.81 cm² and leaf area index of 0.38, 0.79, 1.33 and 2.15 at 30, 60, 90 and 120 DAT.

The increase in leaf area and leaf area index with the application of organic manure could be related to better nutrition allocated to leaf development, due to nutrient release by microorganisms in the soil, which in turn increased plant growth as a result of the production of more assimilates and increased cell division and cell size (Selosse *et al.*, 2004). This finding is similar to the results of Jayasri (2010), Harishkumar *et al.* (2019) and Ram *et al.* (2019a). Meanwhile, the foliar spraying of seaweed extract (*Ascophyllum nodosum*) influenced the vegetative growth of the plant (Gheorghe Cristian Popescu and Monica Popescu, 2014). Increased vegetative growth might have been due to the presence of nutrients and hormonal levels in brown seaweed. These results are in close agreement with Salama and Raina (2015) and Veeranan Uthirapandi *et al.* (2018) in sacred basil.

The data in Table 3 showed significant differences among treatments, with T₉ (FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Seaweed extract @ 2%) recording the highest number of primary branches plant-1 (21.45, 25.37, 32.17, 39.46) and secondary branches plant-1 (64.72, 76.45, 96.56, 109.34), followed by T₁₀ (FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Jeevamrutham @ 3%) with primary branch counts of 20.43, 24.16, 30.68, 37.73 and secondary branch counts of 60.58, 72.43, 90.93, 103.70. The lowest number of primary and secondary branches was observed in the control (T₁) with 12.27, 14.48, 18.76, 23.89 for primary branches and 27.46, 40.27, 45.89, 58.58 for secondary branches at 30, 60, 90 and 120 DAT.

The increased number of branches as a result of higher nitrogen content from FYM was attributed to the involvement of nitrogen in the physiological processes of the plant, which stimulated growth and,

thus, the increased number of branches per plant at higher nitrogen levels were observed. Almost identical results have been reported by Kandil *et al.* (2009) in Holy Basil, Asgharipour (2011), Rahman *et al.* (2014) in Holy Basil and Naggar *et al.* (2015) and EL-Sayed *et al.* (2015) in basil. The basal application of vermicompost stimulated bacterial activity in the soil, nitrogen accumulation and nutrient availability in the plant, resulting in enhanced plant growth (Mohammad Reza Befrozfar *et al.*, 2013). Similar results were obtained by Soheila Shahriari *et al.* (2015) in basil and Chandana *et al.* (2018) in kalmegh and Mohit Lal *et al.* (2018) in sacred basil. According to Dhriti Battacharyya *et al.* (2015), seaweeds affected plant growth due to the effects of oligosaccharides, hormone-like elicitors, betaines and minerals that promoted cell division, protein synthesis and improved stress tolerance.

The data in Table 4 showed significant differences among treatments, with T₉ (FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Seaweed extract @ 2%) recording the highest plant spread North-South (25.32, 33.39, 40.26, 50.38 cm) and East-West (40.56, 55.89, 65.47, 71.45 cm), followed by T₁₀ (FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Jeevamrutham @ 3%) with plant spread North-South of 24.36, 32.25, 38.49, 48.69 cm and East-West of 39.03, 53.29, 62.57, 68.63 cm. The lowest plant spread was observed in the control (T₁), with North-South measurements of 16.68, 23.13, 24.33, 35.16 cm and East-West measurements of 26.79, 32.49, 39.37, 46.07 cm at 30, 60, 90 and 120 DAT.

The superior performance for branches per plant might have been due to the higher availability of nutrients from planting to harvest. The results are in line with those of Munnu Singh (2011) in geranium and Singh and Wasnik (2013) in rosemary Mansour *et al.* (2017) and Netam *et al.* (2020) in basil.

CONCLUSION

Based on the findings of the present study, it can be concluded that combined application of FYM @ 25 t ha⁻¹ + Vermicompost @ 2.5 t ha⁻¹ + Sea weed extract @ 2% has beneficial effect on the growth of Sweet basil.

REFERENCE

- Akash Hidangmayum and Richa Sharma. 2017. Effect of different concentrations of commercial seaweed liquid extract of (*Ascophyllum nodosum*) as a plant bio stimulant on growth, yield and biochemical constituents of onion (*Allium cepa* L.). **J. Pharmacognosy and Phytochemistry.** **6(4)**: 658-663.
- Ali Khanzadeh and Davood Naderi. 2015. Different growing substrates affect Periwinkle (*Catharanthus roseus* L.) growth and flowering. **J. Biodiversity and Environ. Sci.**, **6(6)**: 179-186.
- Asgharipour M I and Rafiei M. 2011. Effect of different organic amendments and drought on the growth and yield of basil in the greenhouse. **Advances in Environmental Biology** **5(6)**: 1233-1239.

Comment [DT8]: Add recommendations based on conclusion of the study

Comment [DT9]: Add acknowledgement and conflict of interest

- Shinde, A., Gahunge, P., Singh, P., Rath, S. K., & Khemani, N. (2013). Effect of inorganic fertilizers and organic manures on growth, quality and yield of ashwagandha (*Withaniasomnifera*Dunal) cv. Jawahar Ashwagandha-20. **Annals of Pharmacy and Pharmaceutical Sciences**, **4(1)**, 2.
- Baraa AL-mansour, M.A. Suryanarayana, D. Kalaivanan, K. Umesha and M. Vasundhara. 2017. Effect of graded levels of N through FYM, inorganic fertilizers and biofertilizers on growth, herbage yield, oil yield and economics of sweet basil (*Ocimum basilicum* L.). **Medicinal Plants -International Journal of Phytomedicines** **9(4)**: 250-258.
- Chandana, M., Veena Joshi, D. Vijaya and D. Lakshminarayana. 2018. Effect of organic treatments and spacing on growth parameters of kalmegh (*Andrographis paniculata*) var. **Cim-Megha. J. Pharmacognosy and Phytochemistry**, **7(6)**: 1695-1699.
- Craige, J.S. 2011. Seaweed extract stimuli in plant science and agriculture. **J. Appl. Phycol.**, **23**: 371-393.
- Davis JM, 1995. Basil Misc. Publ. Northern Carolina Cooperative Extension Service, North Carolina State University, Raleigh, North Carolina, USA.
- Dhriti Battacharyya, Mahbobeh Zamani Babgohari, Pramod Rathor and Balakrishnan Prithiviraj. 2015. Seaweed extracts as biostimulants in horticulture. **Scientia Horticulturae.**, 196: 39-48.
- El-Miniawy, S.M., M.E. Ragab, S.M. Youssef and A.A. Metwally. 2014. Influence of Foliar Spraying of Seaweed Extract on Growth, Yield and Quality of Strawberry Plants. **J. App. Sci. Res.**, **10(2)**: 88-94.
- El-Sayed A A, El-Hanafy S H and El-Ziat R A. 2015. Effect of chicken manure and humic acid on herb and essential oil production of *Ocimum sp.* **American Eurasian Journal of Agriculture and Environmental Science** **15(3)**:367-379.
- Farooqi, A. A. and sreeramu, B. S., 2005, French basil In: Cultivation of medicinal and aromatic crops. Eds, Ali Farooqi and B. S. Sreeramu, Universities Press. pp. 348-353.
- Gunda, V., M. Padma, B. M. Rajkumar, J. Cheena, D. Vijaya and D. S. Chary. 2022. Studies on effect of organic manures on growth and yield parameters of sweet basil (*Ocimum basilicum* L.) under Telangana. **J. Pharm. Innov.**, **11(12)**: 1548 1552.

Gupta, J. P., R. K. Agarwal, G. N. Gupta and P. Kaul. 1983. Effect of continuous application of FYM and urea on soil properties in west Rajasthan. **Indian J. Agri. Sci.**, **53(1)**:53-56.

Harishkumar, J.M., Karishma C., Meenaloshini N., Nagavalli K., Pavithra P., Sowbejan A., Aruna S.J. and Theradimani M. 2019. Effect of bio-fertilizers and vesicular arbuscularmycorrhizae on Holy Basil (*Ocimum basilicum*). **Inter. J. Curr. Microbiol. App. Sci.**, **8(6)**, 1316-1326.

In grapevine, Gheorghe and Monic (2014) reported that the 170 ml ha⁻¹ concentration of seaweed extract showed the highest growth parameters.

Jalil Dehghan Samani, Abdollah Ghasemi Pirbalouti and Fatemeh Malekpoor. 2017. Effect of Organic and Chemical Fertilizers on Growth Parameters and Essential Oil of Iranian Basil (*Ocimum basilicum* L.). **J. Crop Nut Sci.**, **3(1)**: 14-24.

Jayasri, P. and Anuja S. 2010. Effect of organic nutrients on growth and essential oil content of sweet basil (*Ocimum basilicum* L.). **The Asian J. Hort.**, **5(1)**, 26-29.

Kandil MAM, Khatab M E, Ahmed SS and Schnug E. 2009. Herbal and essential oil yield of Genovese basil (*Ocimum basilicum* L.) grown with mineral and organic fertilizer sources in Egypt. **Journal Fur Kulturpflanzen** **61(12)**:443-449.

Katarzyna Chojnacka, Agnieszka Saeid, Zuzanna Witkowska and Lukasz Tuhy. 2012. Biologically Active Compounds in Seaweed Extracts - the Prospects for the Application. **The Open Conference Proceedings Journal.**, **3(1)**: 20-28.

Malav, P., Pandey A., Bhatt K.C., Gopala Krishnan S. and Bisht I.S. 2015. Morphological variability in holy basil (*Ocimum tenuiflorum* L.) from India. **Gene. Resource Crop Evol.**, DOI10.1007/s 10722-015-0227-5.

Mansour, B. A., D. Kalaivanan, M. A. Suryanarayana and A. K. Nair. 2017. Effect of integrated nutrient management on dry herbage yield, nutrient uptake and profitability of French Basil (*Ocimum basilicum* L.). **J. Hort. Sci.**, **12(2)**: 171-179.

Mansour, B. A., D. Kalaivanan, M. A. Suryanarayana and A. K. Nair. 2017. Effect of integrated nutrient management on dry herbage yield, nutrient uptake and profitability of French Basil (*Ocimum basilicum* L.). **J. Hort. Sci.**, **12(2)**: 171-179.

- Meena, K.C., Gontia A.S., Upadhayay A. and Rao S. 2013. Response of *Ocimum* germplasms to foliar application of plant growth promoters. **J. Multidisc. Adv. Res.**, **2(3)**, 25–30.
- Mohammad Reza Befrozfar, Davood Habibi, Ahmad Asgharzadeh, Mehdi SadeghiShoae and Mohammad Reza Tookaloo. 2013. Vermicompost, plant growth promoting bacteria and humic acid can affect the growth and essence of basil (*Ocimum basilicum* L.). **Ann. Biol. Res.**, **4 (2)**: 8-12.
- Mohit Lal, Purnima Gautam, Manjeet Kumar, Jitendra Kumar and Satish Kumar Gautam. 2018. Standardize the ideal dose of different organic sources in growth and yield of Tulsi (*Ocimum sanctum*) cv. **Krishna. Int. J. Chem. Studies.**, **6(6)**: 2588-2590.
- Munnu Singh. 2011. Effect of integrated nutrient management through vermicompost and inorganic fertilizers on growth, yield, nutrient uptake and oil quality of geranium (*Pelargonium graveolens* L" Her. ex Ait.) grown. **J. Spices and Arom. Crops.**, **20 (2)**: 55–59.
- Naggar –El A H M, Hassan M R A, Hassan H E and Mohamed M F A. 2015. Effect of organic biofertilizers on growth, oil yield and chemical composition of the essential oil of (*Ocimumbasilicum*L.) plants. **Alexandria Journal of Agricultural Research** **60(1)**:1-16.
- Netam, Y., P. Toppo and Y. Dewangan. 2020. Effect of different organic manure on CIM Saumya (*Ocimum basilicum* L.) under Eucalyptus (*Eucalyptus tereticornis*) based agro forestry system. **J. Rural Agric. Res.**, **20(2)**: 80-83.
- Padmapriya, S., K. Kumanan and K. Rajamani. 2010. Studies on effect of organic amendments and bio-stimulants on morphology, yield and quality of (*Gymnemasylvestre*) R.Br. Afr. **J. Agricultural Res.**, **5(13)**: 1655-1661.
- Panse, V. G. and P. V. Sukhatme. 1985. Stastical methods for agricultural workers. **Indian Council of Agricultural Research, New Delhi.**
- Patke, N. K., V. V. Tapre, B. M. Muradi, S. G. Wankhade and R. B. Sarode. 2018. Effect of Spacing and Organic Manures on Growth and Yield of (*Aloe vera*.) Int. **J. Curr. Microbiol. App. Sci.**, **(6)**: 2492-2495.

- Rahman K M, Sattar M A and Rahman G MM. 2014. Effect of fertilizer and manures on growth and yield of Tulsi and Pudina medicinal plant. **Journal of Environmental Science and Natural Resources** **7(2)**: 13-16.
- Ram, R., Prasad V.M., Bahadur V., Dowsan J., Swaroop N. and Kumar A. 2019a. Influence of organic manure and bio fertilizer on growth and yield of Indian Basil (*Ocimum sanctum* L.) cv Cim-Ayu and Cim-Angana. **Inter. J. Curr. Microbiol. App. Sci.**, **8(10)**, 2385-2392.
- Ram, R., V. M. Prasad, V. Bahadur, J. Dawson, N. Swaroop and A. Kumar. 2019b. Influence of organic manures and biofertilizers on yield and yield attributes of Indian Basil (*Ocimum sanctum* L.). **J. Pharmacogn. Phytochem.**, **8(5)**: 1809-1813.
- Salama M. Azza and Rania S. Yousef. 2015. Response of Basil Plant (*Ocimum sanctum* L.) to Foliar Spray with Amino Acids or Seaweed Extract. **J. Horticultural Science & Ornamental Plants.**, **7 (3)**: 94-106.
- Selosse, M. A., Baudoin, E., & Vandenkoornhuysen, P. (2004). Symbiotic microorganisms, a key for ecological success and protection of plants. **Comptes rendus biologies**, **327(7)**, 639-648.
- Singh, M., & Wasnik, K. (2013). Effect of vermicompost and chemical fertilizer on growth, herb, oil yield, nutrient uptake, soil fertility, and oil quality of rosemary. **Communications in soil science and plant analysis**, **44(18)**, 2691-2700.
- Soheila Shahriari, Ghobad Shabani and Masomeh Khan Ahmadi. 2015. The Effect of Vermicompost Application on Growth Characteristics and Essential Oil of Basil (*Ocimum basilicum* L.). **Jordan J. Agricultural Sci.**, **11(1)**: 127-134.
- Suresh, V., J Preethi Fetricia, V. Saranya, S. Sarithra and K. Tamilselvan. 2018. To study the effect of fym, coirpith, vermicompost, humic acid and panchagavya on growth and yield of mint (*Mentha arvensis*). **Horti. Int. J.**, **2(6)**: 417-419.
- Suresh, V and R. Senthinathan. 2018. Integrated nutrient management on growth and yield of Thuduvalai (*Solanum trilobatum* L.). **J. Med. Plants Studies.**, **6(3)**: 01-03.
- Tiwari S. K and A.K. Roy. 2014. Influence of different vermicompost doses on the growth performance of (*Gloriosa superba* Linn) in culturable wasteland soil. **Int. J. Basic & App. Sci. Res.**, **1(1)**: 122-131.

Veeranan Uthirapandi, Selvam suriya, PonnerulanBoomibalagan, Saminathan Eswaran, Subramanian Sivasangari Ramya, Narayanan Vijayanand and Durairaj Karthiresan. 2018. Biofertilizer potential of seaweed liquid extracts of marine algae on growth and biochemical parameters of (*Ocimum sanctum*.) **J. Pharmacognosy and Phytochemistry**, **7(3)**: 3528-3532.

Venkatesan, S. 2009. Crop management in medicinal solanum. Ph.D. Thesis, **Annamalai University, Annamalai Nagar, Tamil Nadu**.

Wajahatullah Khan, Usha P. Rayirath, Sowmyalakshmi Subramanian, Mundaya N. Jithesh, Prasanth Rayorath, D. Mark Hodges, Alan T. Critchley, James S. Craigie, Jeff Norrie and Balakrishan Prithiviraj.2009. Seaweed Extracts as Biostimulants of Plant Growth and Development. **J Plant Growth Regul.**, 28:386–399.

UNDER PEER REVIEW