

Simultaneous Hydatid Cysts of the Liver and Pancreas: A Rare Case Report.

ABSTRACT

Background:

Hydatidosis is a common, generally benign condition prevalent in endemic areas, presenting a significant public health challenge, particularly in regions with traditional farming practices. In these endemic regions, hydatid disease can affect in various anatomical sites. Pancreatic hydatid cysts are rare, accounting for less than 1% of cases even in endemic areas. This unusual location may present as chronic epigastric pain, the most common reason for consultation. While ultrasound, computed tomography (CT), and magnetic resonance imaging (MRI) can easily detect pancreatic cystic lesions, determining their hydatid nature preoperatively is extremely challenging to recognize. The treatment involves surgical excision via pericystectomy. Prophylaxis remains the most effective strategy to control this parasitic disease in endemic areas.

Case Report:

We report a case of a 38-year-old woman from an endemic region, presenting with chronic right hypochondriac pain. Imaging revealed multiple hepatic cysts and a well-defined cyst in the pancreatic tail. Positive hydatid serology confirmed the diagnosis. Surgical resection of the cysts was performed with an uneventful recovery, and no recurrence was observed at the three-month follow-up (should be six months at least). This case underscores the diagnostic challenges due to the rarity of pancreatic hydatid cysts, with surgical resection and drainage being the preferred treatment to minimize complications.

Keywords: Hydatid cyst; Pancreatic hydatidosis; echinococcosis; infection; serology; albendazole.

1. INTRODUCTION

Hydatid disease is a cyclozoonosis, a parasitic infection that requires more than one vertebrate host to complete its life cycle. It is endemic in several Mediterranean countries, posing a significant public health problem. The most common form of hydatid cyst, unilocular, is caused by *Echinococcus granulosus*. [1]

In Morocco, hydatid disease remains a major public health concern due to its frequency, morbidity, and mortality, as well as the economic losses it causes [2]

Hydatid cysts can localize in almost any tissue or organ, with the liver (50%-77%), lungs (15%-47%), spleen (0.5%-8%), and kidneys (2%-4%) being the most commonly affected organs. [3]

Pancreatic hydatid cyst (PHC) disease is rare, even in regions where hydatidosis is endemic. While the reported incidences of PHC have varied in different studies, the rates are consistently below 1% [3] Through this case of hydatid cysts of the pancreas, the authors raise diagnostic and therapeutic difficulties and discuss **introduction should be longer**

2. CASE REPORT

A 38-year-old Moroccan woman, with no significant medical history and originating from an endemic hydatid region, was hospitalized due to right hypochondriac pain persisting for two years. Physical examination revealed a patient in good general condition, afebrile, with a palpable, mobile, and tender mass in the right hypochondrium, **be sure about the clinical finding Because cyst in the tail supposed to be felt in the left hypochondrium not in the right** approximately the size of a large mandarin. An abdominal CT scan revealed multiple thin-walled, multilocular cystic formations in the liver.

Additionally, a thin-walled cystic formation was noted in the pancreatic tail (see Figure 1). **should mention the measurement of the cyst in ct scan**



Fig. 1. Computed tomography scan showing cystic mass in tail region of pancreas

An abdominal MRI confirmed a well-defined cystic lesion in the tail of the pancreas, with a homogeneous content and no septation or wall thickening. This lesion did not communicate with the Wirsung duct and was in close contact with the splenic vascular pedicle at its posterior border, as well as with the posterior wall of the stomach anteriorly(**figure2**).**why this figer here**



Fig. 2. Abdominal MRI showing two hydatid cysts.

(indicated by arrows a. hydatid cyst in the pancreatic tail b. hydatid cyst in the liver).

A chest X-ray was normal, and laboratory tests showed positive hydatid serology. The patient underwent exploratory laparotomy via a right subcostal incision extended to the left. (**incision should be midline why start right**) Intraoperative findings revealed three hydatid cysts in the liver (segments VII and VIII) and a mass in the tail of the pancreas in close proximity to the posterior surface of the stomach. After protecting the viscera with 10% hydrogen peroxide-soaked sponges, the dome of the three cysts was resected, and the hydatid material was aspirated and lavage with hydrogen peroxide, with no biliary fistula observed. The cystic mass in the tail of the pancreas was also resected. (figure3)



Fig. 3. Photograph showing the excised hydatid cyst from the tail of the pancreas

Pathological examination of the pancreatic cystic mass confirmed a hydatid cyst. The postoperative course was uneventful, with a six-day hospital stay. Three months post-surgery, no recurrence of hydatidosis was detected. **Material and methods should be in separate title**

Need operative picture

3.DISCUSION

Hydatidosis results from the development of the larval stage of *Echinococcus granulosus* in humans.[4] Pancreatic involvement accounts for less than 1% of hydatid cysts and only 0.2% of abdominal hydatidosis.[5] The most common location is the head of the pancreas (57% of cases), followed by the body (24%), and the tail (19%). Pancreatic infestation typically occurs through the arterial route, after passing through hepatic and pulmonary filters.[6]

The size of the cyst varies, ranging from a few millimeters to over 20 cm.[7]

Symptoms are often insidious and vary depending on the cyst's location and its prolonged development.[8] In this case, no revealing signs were observed. This unusual location can indeed be indicated by chronic epigastric pain, which is the most frequent reason for consultation, as well as by obstructive jaundice (in the case of cephalic location) or an epigastric mass.[6], [7], [9] Certain complications associated with pancreatic hydatid disease, similar to those found in hepatic locations, may be indicative: suppuration and abscess formation of the cyst, leading to a deterioration of the general condition [11],[12]

Intra- or retro-peritoneal rupture can result in hemorrhagic complications and severe peritoneal or allergic reactions[11] Fistulation of the cyst into the biliary tract may cause jaundice[12] Compression of the splenic vein by the cyst can lead to segmental portal hypertension in 14% of cases [12] Pancreatic complications have also been reported, including the occurrence of obstructive chronic pancreatitis[13] Ultrasound, computed tomography (CT), and magnetic resonance imaging (MRI) easily

allow for the diagnosis of a pancreatic cystic lesion; however, the preoperative diagnosis of the hydatid nature of the cyst is extremely difficult to recognize, as seen in our patient.[1]

However, the challenge lies in linking this lesion to hydatid disease. Nevertheless, certain signs can help suggest the diagnosis, including peri cystic calcifications, the presence of intra-cystic vesicles, detachment of the hydatid membrane, or the association with other more obvious locations of hydatid cysts, particularly in the liver.[7]

It is also important to use epidemiological arguments and hydatid serology to make a preoperative diagnosis.

In case of persistent diagnostic uncertainty, the use of endoscopic ultrasound is highly beneficial as it allows for a better examination of the cystic content[10]

The treatment of a hydatid cyst of the pancreas is surgical. The choice of surgical procedure will depend on the location of the cyst and whether or not there is a cystic duct fistula.[11]

Many surgical procedures are available to remove the cyst. Partial or total cystectomy, marsupialization and external drainage have also been reported in the management of pancreatic hydatid cysts[6]In our observation, we performed a resection of the cystic mass in the tail of the pancreas with the cyst being closed.

For body and tail cysts, a left pancreatectomy is often preferred as it allows for the cyst's removal and suturing of the pancreas on healthy tissue [15,16]. Conservative treatment, such as dome resection, is considered only for large cysts adhering to surrounding organs, making dissection difficult and risky. In the presence of a pancreatic fistula, a cyst digestive anastomosis is performed. For head cysts, treatment involves resection of the prominent dome, combined with a cystojejunostomy in the presence of a fistula[15,6]. Radical procedures, such as duodenopancreatectomy, are generally deemed excessive for benign parasitic conditions[17]. Postoperatively, drainage of the residual cavity and perioperative administration of somatostatin analogs like Sandostatin® are recommended to reduce the risk of pancreatic fistula[18].

4.CONCLUSION

Primary hydatid cysts of the pancreas are exceptionally rare. Diagnosis is based on abdominal ultrasound, CT, and positive hydatid serology. MRI and endoscopic ultrasound are useful in cases of persistent diagnostic uncertainty. Surgical treatment typically involves resecting the cyst's prominent dome and extensive drainage of the residual cavity, with perioperative administration of somatostatin to reduce the risk of postoperative pancreatic fistula.

CONSENT

Written informed consent was obtained from the patient for publication of this case report and any accompanying images.

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

REFERENCES

- [1] "Mathai, V., Jesudason, S. R. B., Muthusami, J. C., Kuruvilla, R., Idikula, J., & Sada, P. (1994). Chronic pancreatitis caused by intraductal hydatid cysts of the pancreas. *British Journal of Surgery*, 81(7), 1029–1029." **should be mathai etal**
- [2] O. Derfoufi et al., "Epidemiological profile of hydatid disease in Morocco from 1980 to 2008," *Ann Biol Clin (Paris)*, vol. 70, no. 4, pp. 457–461, Jul. 2012, doi: 10.1684/abc.2012.0727.

- [3] S. Akbulut, "Hydatid cyst of the pancreas: Report of an undiagnosed case of pancreatic hydatid cyst and brief literature review," *World J Gastrointestinal Surg*, vol. 6, no. 10, p. 190, 2014, doi: 10.4240/wjgs.v6.i10.190.
- [4] R. A. Wani, A. A. Malik, N. A. Chowdri, K. A. Wani, and S. H. Naqash, "Primary extrahepatic abdominal hydatidosis," *International Journal of Surgery*, vol. 3, no. 2, pp. 125–127, 2005, doi: 10.1016/j.ijssu.2005.06.004.
- [5] H. Mohamed, S. Azza, A. Chrif, S. Karim, and C. Adnen, "Kystehydatique du pancréas révéle par une pancréatite aiguë: À propos d'un cas," *Pan African Medical Journal*, vol. 22, pp. 69–81, 2015, doi: 10.11604/pamj.2015.22.166.6242.
- [6] A. Fadil, A. A. Bolbarod, and F. El Fares, "Fait clinique Kystehydatique du pancréas. À propos d'une observation," 2000. [7] H. Bedioui et al., "Kystehydatique primitif du pancréas : diagnostic et modalités chirurgicales. À propos de trois cas," *Gastroenterol Clin Biol*, vol. 32, no. 1, pp. 102–106, Jan. 2008, doi: 10.1016/j.gcb.2007.12.014.
- [8] M. Abid et al., "Kystehydatique du pancréas: Une localization exceptionnelle," *Archives de Pédiatrie*, vol. 17, no. 7, pp. 1056–1058, Jul. 2010, doi: 10.1016/j.arcped.2010.03.005.
- [9] "Lemmer ER, Krige JE, Price SK, Girdwood AH. Hydatid cyst in the head of the pancreas with obstructive jaundice. *J Clin Gastroenterol*. 1995 Mar;20(2)-136-8. doi: 10.1097:00004836-199503000-00013. PMID- 7769194."
- [10] "Palazzo, L., Hammel, P., Cellier, C., & Ruszniewski, P. (2000). Les tumeurs kystiques du pancréas. *Acta Endoscopica*, 30(S2), 361–366".
- [11] H. Bedioui et al., "Kystehydatique primitif du pancréas : diagnostic et modalités chirurgicales. À propos de trois cas," *Gastroenterol Clin Biol*, vol. 32, no. 1, pp. 102–106, Jan. 2008, doi: 10.1016/j.gcb.2007.12.014.
- [12] Belakhdar A, Mokhtari M, Khadiri R, Mansouri A, Bouzidi S. Kystehydatique du pancréas à propos de two observations. *Acta Chir Belg* 1987;87:225—7.
- [13] Cosme A, Orive V, Ojeda E, Aramburu V, Irazusta M, Arenas JL. Hydatid cyst of the head of the . pancreas with spontaneous fist [14] Katkhouda N, Legoff D, Tricarico A, Castillo L, Bertrand M, Mouiel J. Kystehydatique du . pancréas responsable for a recurrent chronic pancréatit. *Presse Med* 1988;17:2021—3.
- [15] MERCADIER M. -- M~thodes et indications du traitement des formations kystiques et pseudo-. . kystiques du pancr6as. LX ° Congr6s de Chirurgie, Paris 1958. [16] Moosavi SR, Kermany HK. Epigastric mass due to a hydatid cyst of the pancreas. At home..... report and review of the literature. *J Pancreas* 2007;8:232–4 [17] Ismail K, Haluk GI, Necati O. Surgical treatment of hydatid cysts of the pancreas. *Int Surg*. 1991;76:185–8.
- [18] 21 Ennabli E, Zaouch A, Kheireddine E, Horchani H, Slim R, Ben Jilani S. Kyste.hydatique du . pancréas avec érosion du Wirsung. *Med Chir Dig* 1984;13:357—60.