

A Study On Matrix Sequence with Generalized Guglielmo Numbers Components

Abstract. We know that the diverse applications of matrix sequences in fields such as physics, engineering, architecture, nature, and art. Numerous authors have delved into the study of these matrix sequences in existing literature. Furthermore, in this study, we define and investigate the generalized Guglielmo matrix sequence. Hence, in detail, we explore four specific cases of that sequence that called triangular matrix sequences, Lucas-triangular matrix sequences, oblong matrix sequences, and pentagonal matrix sequences. We present Binet's formulas, generating functions, and the summation formulas for these sequences. Moreover, we give some identities and matrices related with these sequences. Furthermore, we show that there always exist some interrelation between generalized triangular, Lucas-triangular, oblong and pentagonal matrix sequences.

2020 Mathematics Subject Classification. 11B37, 11B39, 11B83.

Keywords. Triangular, Lucas-triangular, oblong and pentagonal numbers, triangular matrix sequence, Lucas-triangular matrix sequence, oblong matrix sequence and pentagonal matrix sequence.

1. Introduction

In this section, we present some properties of the generalized Guglielmo sequence, such as recurrence relations, Binet's formula, generating function and characteristic equations, that we will need rest of the our study. For more detail see, [10].

A generalized Guglielmo sequence $\{W_n\}_{n \geq 0} = \{W_n(W_0, W_1, W_2)\}_{n \geq 0}$ is given by the third-order recurrence relations.

$$W_n = 3W_{n-1} - 3W_{n-2} + W_{n-3} \tag{1.1}$$

with the initial values $W_0 = c_0, W_1 = c_1, W_2 = c_2$ not all being zero.

The sequence $\{W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$W_{-n} = 3W_{-(n-1)} - 3W_{-(n-2)} + W_{-(n-3)}$$

for $n = 1, 2, 3, \dots$. Therefore, recurrence (1.1) holds for all integer n .

Now we present four special cases of the sequence $\{W_n\}$. Triangular sequence $\{T_n\}_{n \geq 0}$, triangular-Lucas sequence $\{H_n\}_{n \geq 0}$, oblong sequence $\{O_n\}_{n \geq 0}$ and pentagonal sequence $\{p_n\}_{n \geq 0}$ are defined, respectively, by the third-order recurrence relations

$$T_n = 3T_{n-1} - 3T_{n-2} + T_{n-3}, \quad T_0 = 0, T_1 = 1, T_2 = 3, \tag{1.2}$$

$$H_n = 3H_{n-1} - 3H_{n-2} + H_{n-3}, \quad H_0 = 3, H_1 = 3, H_2 = 3, \tag{1.3}$$

$$O_n = 3O_{n-1} - 3O_{n-2} + O_{n-3}, \quad O_0 = 0, O_1 = 2, O_2 = 6, \tag{1.4}$$

$$p_n = 3p_{n-1} - 3p_{n-2} + p_{n-3}, \quad p_0 = 0, p_1 = 1, p_2 = 5. \tag{1.5}$$

The sequences $\{T_n\}_{n \geq 0}$, $\{H_n\}_{n \geq 0}$, $\{O_n\}_{n \geq 0}$ and $\{p_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$T_{-n} = 3T_{-(n-1)} - 3T_{-(n-2)} + T_{-(n-3)},$$

$$H_{-n} = 3H_{-(n-1)} - 3H_{-(n-2)} + H_{-(n-3)},$$

$$O_{-n} = 3O_{-(n-1)} - 3O_{-(n-2)} + O_{-(n-3)},$$

$$p_{-n} = 3p_{-(n-1)} - 3p_{-(n-2)} + p_{-(n-3)}$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrences (1.2)-(1.5) hold for all integer n . Now, we give some properties related to generalized Guglielmo numbers that we need for the rest of the study.

- The Binet formula of generalized Guglielmo numbers. Binet formula of generalized Guglielmo numbers can be given as

$$W_n = A_1 + A_2n + A_3n^2 \tag{1.6}$$

where

$$A_1 = W_0,$$

$$A_2 = \frac{1}{2}(-W_2 + 4W_1 - 3W_0),$$

$$A_3 = \frac{1}{2}(W_2 - 2W_1 + W_0),$$

i.e.,

$$W_n = W_0 + \frac{1}{2}(-W_2 + 4W_1 - 3W_0)n + \frac{1}{2}(W_2 - 2W_1 + W_0)n^2. \tag{1.7}$$

- For all integers n , triangular, triangular-Lucas, oblong and pentagonal numbers (using initial conditions in (1.7)) can be expressed using Binet’s formulas as

$$\begin{aligned} T_n &= \frac{n(n+1)}{2}, \\ H_n &= 3, \\ O_n &= n(n+1), \\ p_n &= \frac{1}{2}n(3n-1) \end{aligned}$$

respectively.

- Suppose that $f_{W_n}(x) = \sum_{n=0}^{\infty} W_n x^n$ is the ordinary generating function of the generalized Guglielmo sequence $\{W_n\}_{n \geq 0}$. Then, $\sum_{n=0}^{\infty} W_n x^n$ is given by

$$\sum_{n=0}^{\infty} W_n x^n = \frac{W_0 + (W_1 - 3W_0)x + (W_2 - 3W_1 + 3W_0)x^2}{1 - 3x + 3x^2 - x^3}. \tag{1.8}$$

- Here, the characteristic equation of the generalized Guglielmo sequence

$$x^3 - 3x^2 + 3x - 1 = 0.$$

- (Simpson’s formula for generalized Guglielmo numbers)For all integers n , we have

$$\begin{vmatrix} W_{n+2} & W_{n+1} & W_n \\ W_{n+1} & W_n & W_{n-1} \\ W_n & W_{n-1} & W_{n-2} \end{vmatrix} = -(W_2 - 2W_1 + W_0)^3.$$

Next, the first few generalized Guglielmo numbers with positive subscript and negative subscript is given in the following Table 1.

Table 1. A few generalized Guglielmo numbers

n	W_n	W_{-n}
0	W_0	W_0
1	W_1	$3W_0 - 3W_1 + W_2$
2	W_2	$6W_0 - 8W_1 + 3W_2$
3	$W_0 - 3W_1 + 3W_2$	$10W_0 - 15W_1 + 6W_2$
4	$3W_0 - 8W_1 + 6W_2$	$15W_0 - 24W_1 + 10W_2$
5	$6W_0 - 15W_1 + 10W_2$	$21W_0 - 35W_1 + 15W_2$
6	$10W_0 - 24W_1 + 15W_2$	$28W_0 - 48W_1 + 21W_2$
7	$15W_0 - 35W_1 + 21W_2$	$36W_0 - 63W_1 + 28W_2$
8	$21W_0 - 48W_1 + 28W_2$	$45W_0 - 80W_1 + 36W_2$
9	$28W_0 - 63W_1 + 36W_2$	$55W_0 - 99W_1 + 45W_2$
10	$36W_0 - 80W_1 + 45W_2$	$66W_0 - 120W_1 + 55W_2$
11	$45W_0 - 99W_1 + 55W_2$	$78W_0 - 143W_1 + 66W_2$
12	$55W_0 - 120W_1 + 66W_2$	$91W_0 - 168W_1 + 78W_2$
13	$66W_0 - 143W_1 + 78W_2$	$105W_0 - 195W_1 + 91W_2$

Next, we present the first few values of the Triangular and Triangular-Lucas, oblong and pentagonal numbers with positive and negative subscripts:

Table 2. The first few values of the special third-order numbers with positive and negative subscripts.

n	0	1	2	3	4	5	6	7	8	9	10	11	12	13
T_n	0	1	3	6	10	15	21	28	36	45	55	66	78	91
T_{-n}		0	1	3	6	10	15	21	28	36	45	55	66	78
H_n	3	3	3	3	3	3	3	3	3	3	3	3	3	3
H_{-n}		3	3	3	3	3	3	3	3	3	3	3	3	3
O_n	0	2	6	12	20	30	42	56	72	90	110	132	156	182
O_{-n}		0	2	6	12	20	30	42	56	72	90	110	132	156
p_n	0	1	5	12	22	35	51	70	92	117	145	176	210	247
p_{-n}		2	7	15	26	40	57	77	100	126	155	187	222	260

For more detail, see [10].

2. The Matrix Sequences of Generalized Guglielmo Numbers

Recently, numerous studies have explored sequences of numbers in the literature, focusing on sequences of Horadam (generalized Fibonacci) numbers and generalized Tribonacci numbers. These include well-known sequences such as Fibonacci, Lucas, Pell, and Jacobsthal numbers, as well as third-order Pell, third-order Pell-Lucas, Padovan, Perrin, Padovan-Perrin, Narayana, third-order Jacobsthal and third-order Jacobsthal-Lucas numbers. The sequences of numbers have found widespread applications in various research areas, including

physics, engineering, architecture, nature and art. Conversely, matrix sequences have garnered attention, particularly for different types of numbers. Next, we present some study published in the literature related to matrix sequences.

- For study related to Generalized Fibonacci matrix sequences see, [2,3,4,11,12,13,14,15,18].
- For study related to Generalized Tribonacci matrix sequences see,[1,6,7,8,9,16,17].
- For study related to Generalized Tetranacci matrix sequences see, [5].

In this section we define generalized Guglielmo matrix sequence and investigate its properties.

DEFINITION 2.1. For any integer $n \geq 0$, the generalized Guglielmo matrix $(\mathcal{M}W_n)$ is defined by

$$\mathcal{M}W_n = 3\mathcal{M}W_{n-1} - 3\mathcal{M}W_{n-2} + \mathcal{M}W_{n-3} \tag{2.1}$$

with initial conditions

$$\begin{aligned} \mathcal{M}W_0 &= \begin{pmatrix} W_1 & W_2 - 3W_1 & W_0 \\ W_0 & W_1 - 3W_0 & 3W_0 - 3W_1 + W_2 \\ 3W_0 - 3W_1 + W_2 & 9W_1 - 8W_0 - 3W_2 & 6W_0 - 8W_1 + 3W_2 \end{pmatrix}, \\ \mathcal{M}W_1 &= \begin{pmatrix} W_2 & W_0 - 3W_1 & W_1 \\ W_1 & W_2 - 3W_1 & W_0 \\ W_0 & W_1 - 3W_0 & 3W_0 - 3W_1 + W_2 \end{pmatrix}, \\ \mathcal{M}W_2 &= \begin{pmatrix} W_0 - 3W_1 + 3W_2 & W_1 - 3W_2 & W_2 \\ W_2 & W_0 - 3W_1 & W_1 \\ W_1 & W_2 - 3W_1 & W_0 \end{pmatrix}. \end{aligned}$$

The sequence $\{\mathcal{M}W_n\}_{n \geq 0}$ can be extended to negative subscripts by defining

$$\mathcal{M}W_{-n} = 3\mathcal{M}W_{-(n-1)} - 3\mathcal{M}W_{-(n-2)} + \mathcal{M}W_{-(n-3)}$$

for $n = 1, 2, 3, \dots$ respectively. Therefore, recurrence (2.1) holds for all integers n .

Three special cases of generalized Guglielmo matrix sequence, taking $W_n = T_n, W_n = H_n, W_n = O_n, W_n = p_n$, respectively, can be defined as follows.

DEFINITION 2.2. For any integer $n \geq 0$, the triangular matrix $(\mathcal{M}T_n)$, Lucas-triangular matrix $(\mathcal{M}H_n)$, oblong matrix $(\mathcal{M}O_n)$ and pentagonal matrix $(\mathcal{M}p_n)$ are defined by

$$\begin{aligned} \mathcal{M}T_n &= 3\mathcal{M}T_{n-1} - 3\mathcal{M}T_{n-2} + \mathcal{M}T_{n-3}, \\ \mathcal{M}H_n &= 3\mathcal{M}H_{n-1} - 3\mathcal{M}H_{n-2} + \mathcal{M}H_{n-3}, \\ \mathcal{M}O_n &= 3\mathcal{M}O_{n-1} - 3\mathcal{M}O_{n-2} + \mathcal{M}O_{n-3}, \\ \mathcal{M}p_n &= 3\mathcal{M}p_{n-1} - 3\mathcal{M}p_{n-2} + \mathcal{M}p_{n-3} \end{aligned}$$

respectively, with initial conditions

$$\begin{aligned}
 \mathcal{MT}_0 &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \mathcal{MT}_1 = \begin{pmatrix} 3 & -3 & 1 \\ 1 & 0 & 0 \\ 0 & 1 & 0 \end{pmatrix}, \mathcal{MT}_2 = \begin{pmatrix} 6 & -8 & 3 \\ 3 & -3 & 1 \\ 1 & 0 & 0 \end{pmatrix}, \\
 \mathcal{MH}_0 &= \begin{pmatrix} 3 & -6 & 3 \\ 3 & -6 & 3 \\ 3 & -6 & 3 \end{pmatrix}, \mathcal{MH}_1 = \begin{pmatrix} 3 & -6 & 3 \\ 3 & -6 & 3 \\ 3 & -6 & 3 \end{pmatrix}, \mathcal{MH}_2 = \begin{pmatrix} 3 & -6 & 3 \\ 3 & -6 & 3 \\ 3 & -6 & 3 \end{pmatrix}, \\
 \mathcal{MO}_0 &= \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}, \mathcal{MO}_1 = \begin{pmatrix} 6 & -6 & 2 \\ 2 & 0 & 0 \\ 0 & 2 & 0 \end{pmatrix}, \mathcal{MO}_2 = \begin{pmatrix} 12 & -16 & 6 \\ 6 & -6 & 2 \\ 2 & 0 & 0 \end{pmatrix}, \\
 \mathcal{Mp}_0 &= \begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 2 \\ 2 & -6 & 7 \end{pmatrix}, \mathcal{Mp}_1 = \begin{pmatrix} 5 & -3 & 1 \\ 1 & 2 & 0 \\ 0 & 1 & 2 \end{pmatrix}, \mathcal{Mp}_2 = \begin{pmatrix} 12 & -14 & 5 \\ 5 & -3 & 1 \\ 1 & 2 & 0 \end{pmatrix}.
 \end{aligned}$$

The sequences (\mathcal{MT}_n) , (\mathcal{MH}_n) , (\mathcal{MO}_n) and (\mathcal{Mp}_n) can be extended to negative subscripts by defining

$$\begin{aligned}
 \mathcal{MT}_{-n} &= 3\mathcal{MT}_{-(n-1)} - 3\mathcal{MT}_{-(n-2)} + \mathcal{MT}_{-(n-3)}, \\
 \mathcal{MH}_{-n} &= 3\mathcal{MH}_{-(n-1)} - 3\mathcal{MH}_{-(n-2)} + \mathcal{MH}_{-(n-3)}, \\
 \mathcal{MO}_{-n} &= 3\mathcal{MO}_{-(n-1)} - 3\mathcal{MO}_{-(n-2)} + \mathcal{MO}_{-(n-3)}, \\
 \mathcal{Mp}_{-n} &= 3\mathcal{Mp}_{-(n-1)} - 3\mathcal{Mp}_{-(n-2)} + \mathcal{Mp}_{-(n-3)}
 \end{aligned}$$

for $n = 1, 2, 3, \dots$ respectively.

The following theorem gives the n th general terms of the generalized Guglielmo matrix sequence.

THEOREM 2.3. *For any integer n , we have the following formulas of the generalized Guglielmo matrix sequence:*

$$\mathcal{MW}_n = \begin{pmatrix} W_{n+1} & -3W_n + W_{n-1} & W_n \\ W_n & -3W_{n-1} + W_{n-2} & W_{n-1} \\ W_{n-1} & -3W_{n-2} + W_{n-3} & W_{n-2} \end{pmatrix}. \tag{2.2}$$

Proof. Suppose that $n \geq 0$. We prove (2.2) by mathematical induction on n . If $n = 0$, since $W_{-1} = 3W_0 - 3W_1 + W_2$, $W_{-2} = 6W_0 - 8W_1 + 3W_2$, $W_{-3} = 10W_0 - 15W_1 + 6W_2$, we have

$$\mathcal{MW}_0 = \begin{pmatrix} W_1 & -3W_0 + W_{-1} & W_0 \\ W_0 & -3W_{-1} + W_{-2} & W_{-1} \\ W_{-1} & -3W_{-2} + W_{-3} & W_{-2} \end{pmatrix} = \begin{pmatrix} W_1 & W_2 - 3W_1 & W_0 \\ W_0 & W_1 - 3W_0 & 3W_0 - 3W_1 + W_2 \\ 3W_0 - 3W_1 + W_2 & 9W_1 - 8W_0 - 3W_2 & 6W_0 - 8W_1 + 3W_2 \end{pmatrix}$$

which is true. Assume that the equality holds for $n \leq k$. For $n = k + 1$, we have

$$\begin{aligned} \mathcal{M}W_{k+1} &= 3\mathcal{M}W_k - 3\mathcal{M}W_{k-1} + \mathcal{M}W_{k-2} \\ &= 3 \begin{pmatrix} W_{k+1} & -3W_k + W_{k-1} & W_k \\ W_k & -3W_{k-1} + W_{k-2} & W_{k-1} \\ W_{k-1} & -3W_{k-2} + W_{k-3} & W_{k-2} \end{pmatrix} - 3 \begin{pmatrix} W_{(k-1)+1} & -3W_{(k-1)} + W_{(k-1)-1} & W_{(k-1)} \\ W_{(k-1)} & -3W_{(k-1)-1} + W_{(k-1)-2} & W_{(k-1)-1} \\ W_{(k-1)-1} & -3W_{(k-1)-2} + W_{(k-1)-3} & W_{(k-1)-2} \end{pmatrix} \\ &\quad + \begin{pmatrix} W_{(k-2)+1} & -3W_{(k-2)} + W_{(k-2)-1} & W_{(k-2)} \\ W_{(k-2)} & -3W_{(k-2)-1} + W_{(k-2)-2} & W_{(k-2)-1} \\ W_{(k-2)-1} & -3W_{(k-2)-2} + W_{(k-2)-3} & W_{(k-2)-2} \end{pmatrix} \\ &= \begin{pmatrix} W_{(k+1)+1} & -3W_{k+1} + W_{(k+1)-1} & W_{k+1} \\ W_{k+1} & -3W_{(k+1)-1} + W_{(k+1)-2} & W_{(k+1)-1} \\ W_{(k+1)-1} & -3W_{(k+1)-2} + W_{(k+1)-3} & W_{(k+1)-2} \end{pmatrix}. \end{aligned}$$

Thus, by strong induction on $k + 1$, this proves (2.2).

For the case $n \leq 0$, similarly,(2.2) can be proved by strong mathematical induction on n . \square

The following corollary gives the n th general terms of the the triangular matrix $(\mathcal{M}T_n)$, Lucas-triangular matrix $(\mathcal{M}H_n)$, oblong matrix $(\mathcal{M}O_n)$ and pentegonal matrix $(\mathcal{M}p_n)$.

COROLLARY 2.4. *For any integer n , we have the following formulas of the matrix sequences:*

$$\begin{aligned} \mathcal{M}T_n &= \begin{pmatrix} T_{n+1} & -3T_n + T_{n-1} & T_n \\ T_n & -3T_{n-1} + T_{n-2} & T_{n-1} \\ T_{n-1} & -3T_{n-2} + T_{n-3} & T_{n-2} \end{pmatrix}, \\ \mathcal{M}H_n &= \begin{pmatrix} H_{n+1} & -3H_n + H_{n-1} & H_n \\ H_n & -3H_{n-1} + H_{n-2} & H_{n-1} \\ H_{n-1} & -3H_{n-2} + H_{n-3} & H_{n-2} \end{pmatrix}, \\ \mathcal{M}O_n &= \begin{pmatrix} O_{n+1} & -3O_n + O_{n-1} & O_n \\ O_n & -3O_{n-1} + O_{n-2} & O_{n-1} \\ O_{n-1} & -3O_{n-2} + O_{n-3} & O_{n-2} \end{pmatrix}, \\ \mathcal{M}p_n &= \begin{pmatrix} p_{n+1} & -3p_n + p_{n-1} & p_n \\ p_n & -3p_{n-1} + p_{n-2} & p_{n-1} \\ p_{n-1} & -3p_{n-2} + p_{n-3} & p_{n-2} \end{pmatrix}. \end{aligned}$$

LEMMA 2.5. *We suppose that $\alpha = \beta = \gamma$, the Binet's formula of generalized (r, s, t) sequence is*

$$W_n = (A_1 + A_2n + A_3n^2) \times \alpha^n \tag{2.3}$$

where α, β, γ are the roots of the characteristic equation of the generalized (r, s, t) sequence and

$$\begin{aligned} A_1 &= W_0, \\ A_2 &= \frac{1}{2}(-W_2 + 4W_1 - 3W_0), \\ A_3 &= \frac{1}{2}(W_2 - 2W_1 + W_0). \end{aligned}$$

We now give the Binet's formula for the generalized Guglielmo matrix sequence.

THEOREM 2.6. *For every integer n , the Binet's formula of the generalized Guglielmo matrix sequence are given by*

$$MW_n = MA_1 + MA_2n + MA_3n^2 \tag{2.4}$$

where

$$\begin{aligned} MA_1 &= MW_0, \\ MA_2 &= \frac{1}{2}(-MW_2 + 4MW_1 - 3MW_0), \\ MA_3 &= \frac{1}{2}(MW_2 - 2MW_1 + MW_0). \end{aligned}$$

Proof. The proof can be done easily by using Lemma 2.5. \square

The following corollary gives the Binet's formulas of the triangular, Lucas-triangular, oblong and pentagonals matrix sequences.

COROLLARY 2.7. *For every integer n , the Binet formulas of the triangular, Lucas-triangular, oblong and pentagonals matrix sequences, respectively, are given by*

(a):

$$MT_n = MB_1 + MB_2n + MB_3n^2$$

where

$$\begin{aligned} MB_1 &= MT_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \\ MB_2 &= \frac{1}{2}(-MT_2 + 4MT_1 - 3MT_0) = \begin{pmatrix} \frac{3}{2} & -2 & \frac{1}{2} \\ \frac{1}{2} & 0 & -\frac{1}{2} \\ -\frac{1}{2} & 2 & -\frac{3}{2} \end{pmatrix}, \\ MB_3 &= \frac{1}{2}(MT_2 - 2MT_1 + MT_0) = \begin{pmatrix} \frac{1}{2} & -1 & \frac{1}{2} \\ \frac{1}{2} & -1 & \frac{1}{2} \\ \frac{1}{2} & -1 & \frac{1}{2} \end{pmatrix}. \end{aligned}$$

i.e.

$$\mathcal{M}T_n = \begin{pmatrix} \frac{1}{2}n^2 + \frac{3}{2}n + 1 & -n^2 - 2n & \frac{1}{2}n^2 + \frac{1}{2}n \\ \frac{1}{2}n^2 + \frac{1}{2}n & 1 - n^2 & \frac{1}{2}n^2 - \frac{1}{2}n \\ \frac{1}{2}n^2 - \frac{1}{2}n & 2n - n^2 & \frac{1}{2}n^2 - \frac{3}{2}n + 1 \end{pmatrix}.$$

(b):

$$\mathcal{M}H_n = \mathcal{M}C_1 + \mathcal{M}C_2n + \mathcal{M}C_3n^2$$

where

$$\mathcal{M}C_1 = \mathcal{M}H_0 = \begin{pmatrix} 3 & -6 & 3 \\ 3 & -6 & 3 \\ 3 & -6 & 3 \end{pmatrix},$$

$$\mathcal{M}C_2 = \frac{1}{2}(-\mathcal{M}H_2 + 4\mathcal{M}H_1 - 3\mathcal{M}H_0) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix},$$

$$\mathcal{M}C_3 = \frac{1}{2}(\mathcal{M}H_2 - 2\mathcal{M}H_1 + \mathcal{M}H_0) = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

i.e.

$$\mathcal{M}H_n = \begin{pmatrix} 3 & -6 & 3 \\ 3 & -6 & 3 \\ 3 & -6 & 3 \end{pmatrix}.$$

(c):

$$\mathcal{M}O_n = \mathcal{M}D_1 + \mathcal{M}D_2n + \mathcal{M}D_3n^2$$

where

$$\mathcal{M}D_1 = \mathcal{M}O_0 = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix},$$

$$\mathcal{M}D_2 = \frac{1}{2}(-\mathcal{M}O_2 + 4\mathcal{M}O_1 - 3\mathcal{M}O_0) = \begin{pmatrix} 3 & -4 & 1 \\ 1 & 0 & -1 \\ -1 & 4 & -3 \end{pmatrix},$$

$$\mathcal{M}D_3 = \frac{1}{2}(\mathcal{M}O_2 - 2\mathcal{M}O_1 + \mathcal{M}O_0) = \begin{pmatrix} 1 & -2 & 1 \\ 1 & -2 & 1 \\ 1 & -2 & 1 \end{pmatrix}.$$

i.e.

$$\mathcal{M}O_n = \begin{pmatrix} n^2 + 3n + 2 & -2n^2 - 4n & n^2 + n \\ n^2 + n & 2 - 2n^2 & n^2 - n \\ n^2 - n & 4n - 2n^2 & n^2 - 3n + 2 \end{pmatrix}.$$

(d):

$$\mathcal{M}p_n = \mathcal{M}E_1 + \mathcal{M}E_2n + \mathcal{M}E_3n^2$$

where

$$\begin{aligned} \mathcal{M}E_1 &= \mathcal{M}p_0 = \begin{pmatrix} 1 & 2 & 0 \\ 0 & 1 & 2 \\ 2 & -6 & 7 \end{pmatrix}, \\ \mathcal{M}E_2 &= \frac{1}{2}(-\mathcal{M}p_2 + 4\mathcal{M}p_1 - 3\mathcal{M}p_0) = \begin{pmatrix} \frac{5}{2} & -2 & -\frac{1}{2} \\ -\frac{1}{2} & 4 & -\frac{7}{2} \\ -\frac{7}{2} & 10 & -\frac{13}{2} \end{pmatrix}, \\ \mathcal{M}E_3 &= \frac{1}{2}(\mathcal{M}p_2 - 2\mathcal{M}p_1 + \mathcal{M}p_0) = \begin{pmatrix} \frac{3}{2} & -3 & \frac{3}{2} \\ \frac{3}{2} & -3 & \frac{3}{2} \\ \frac{3}{2} & -3 & \frac{3}{2} \end{pmatrix}. \end{aligned}$$

i.e.

$$\mathcal{M}p_n = \begin{pmatrix} \frac{3}{2}n^2 + \frac{5}{2}n + 1 & -3n^2 - 2n + 2 & \frac{3}{2}n^2 - \frac{1}{2}n \\ \frac{3}{2}n^2 - \frac{1}{2}n & -3n^2 + 4n + 1 & \frac{3}{2}n^2 - \frac{7}{2}n + 2 \\ \frac{3}{2}n^2 - \frac{7}{2}n + 2 & -3n^2 + 10n - 6 & \frac{3}{2}n^2 - \frac{13}{2}n + 7 \end{pmatrix}.$$

Now, we present some summation formulas for the generalized Guglielmo matrix sequence.

THEOREM 2.8. *For all integers m, j we have*

(a):

$$\sum_{k=0}^{n-1} \mathcal{M}W_{km+j} = \sum_{k=0}^{n-1} (n\mathcal{M}A_1 + \mathcal{M}A_2 \frac{n(2j - m + mn)}{2}) \tag{2.5}$$

$$+ \mathcal{M}A_3 \frac{n(6j^2 + 6jmn - 6jm + 2m^2n^2 - 3m^2n + m^2)}{6}. \tag{2.6}$$

(b):

$$\begin{aligned} \sum_{k=0}^{n-1} k\mathcal{M}W_{km+j} &= \mathcal{M}A_1 \frac{n(n-1)}{2} + \mathcal{M}A_2 \frac{n(n-1)(3j - m + 2mn)}{6} \\ &+ \mathcal{M}A_3 \frac{n(n-1)(6j^2 + 8jmn - 4jm + 3m^2n^2 - 3m^2n)}{12}. \end{aligned}$$

Proof.

(a) Using (2.4), we get

$$\sum_{k=0}^{n-1} \mathcal{M}W_{km+j} = \sum_{k=0}^{n-1} (\mathcal{M}A_1 + \mathcal{M}A_2(j+km) + \mathcal{M}A_3(j+km)^2) \quad (2.7)$$

$$\begin{aligned} &= \sum_{k=0}^{n-1} \mathcal{M}A_1 + \sum_{k=0}^{n-1} \mathcal{M}A_2(j+km) + \sum_{k=0}^{n-1} \mathcal{M}A_3(j+km)^2 \\ &= \sum_{k=0}^{n-1} (n\mathcal{M}A_1 + \mathcal{M}A_2 \frac{n(2j-m+mn)}{2} \\ &\quad + \mathcal{M}A_3 \frac{n(6j^2 + 6jmn - 6jm + 2m^2n^2 - 3m^2n + m^2)}{6}). \end{aligned} \quad (2.8)$$

(b) Using (2.4), we get

$$\begin{aligned} \sum_{k=0}^{n-1} k\mathcal{M}W_{km+j} &= \sum_{k=0}^{n-1} (k\mathcal{M}A_1 + \mathcal{M}A_2k(j+km) + \mathcal{M}A_3k(j+km)^2) \\ &= \sum_{k=0}^{n-1} k\mathcal{M}A_1 + \sum_{k=0}^{n-1} \mathcal{M}A_2k(j+km) + \sum_{k=0}^{n-1} \mathcal{M}A_3k(j+km)^2 \\ &= \mathcal{M}A_1 \frac{n(n-1)}{2} + \mathcal{M}A_2 \frac{n(n-1)(3j-m+2mn)}{6} \\ &\quad + \mathcal{M}A_3 \frac{n(n-1)(6j^2 + 8jmn - 4jm + 3m^2n^2 - 3m^2n)}{12}. \quad \square \end{aligned}$$

From the theorem given above, we obtain the following corollary, which deals with the summation formulas for the triangular, Lucas-triangular, oblong, and pentagonal matrix sequences.

COROLLARY 2.9. *For all integers m, j we have*

(a):

$$\begin{aligned} \sum_{k=0}^{n-1} \mathcal{M}T_{km+j} &= \sum_{k=0}^{n-1} (n\mathcal{M}B_1 + \mathcal{M}B_2 \frac{n(2j-m+mn)}{2} \\ &\quad + \mathcal{M}B_3 \frac{n(6j^2 + 6jmn - 6jm + 2m^2n^2 - 3m^2n + m^2)}{6}), \\ \sum_{k=0}^{n-1} k\mathcal{M}T_{km+j} &= \mathcal{M}B_1 \frac{n(n-1)}{2} + \mathcal{M}B_2 \frac{n(n-1)(3j-m+2mn)}{6} \\ &\quad + \mathcal{M}B_3 \frac{n(n-1)(6j^2 + 8jmn - 4jm + 3m^2n^2 - 3m^2n)}{12} \end{aligned}$$

where

$$\begin{aligned}\mathcal{M}B_1 &= \mathcal{M}T_0, \\ \mathcal{M}B_2 &= \frac{1}{2}(-\mathcal{M}T_2 + 4\mathcal{M}T_1 - 3\mathcal{M}T_0), \\ \mathcal{M}B_3 &= \frac{1}{2}(\mathcal{M}T_2 - 2\mathcal{M}T_1 + \mathcal{M}T_0).\end{aligned}$$

(b):

$$\begin{aligned}\sum_{k=0}^{n-1} \mathcal{M}O_{km+j} &= \sum_{k=0}^{n-1} (n\mathcal{M}C_1 + \mathcal{M}C_2 \frac{n(2j-m+mn)}{2} \\ &\quad + \mathcal{M}C_3 \frac{n(6j^2 + 6jmn - 6jm + 2m^2n^2 - 3m^2n + m^2)}{6}), \\ \sum_{k=0}^{n-1} k\mathcal{M}O_{km+j} &= \mathcal{M}C_1 \frac{n(n-1)}{2} + \mathcal{M}C_2 \frac{n(n-1)(3j-m+2mn)}{6} \\ &\quad + \mathcal{M}C_3 \frac{n(n-1)(6j^2 + 8jmn - 4jm + 3m^2n^2 - 3m^2n)}{12}\end{aligned}$$

where

$$\begin{aligned}\mathcal{M}C_1 &= \mathcal{M}H_0, \\ \mathcal{M}C_2 &= \frac{1}{2}(-\mathcal{M}H_2 + 4\mathcal{M}H_1 - 3\mathcal{M}H_0), \\ \mathcal{M}C_3 &= \frac{1}{2}(\mathcal{M}H_2 - 2\mathcal{M}H_1 + \mathcal{M}H_0).\end{aligned}$$

(c):

$$\begin{aligned}\sum_{k=0}^{n-1} \mathcal{M}O_{km+j} &= \sum_{k=0}^{n-1} (n\mathcal{M}D_1 + \mathcal{M}D_2 \frac{n(2j-m+mn)}{2} \\ &\quad + \mathcal{M}D_3 \frac{n(6j^2 + 6jmn - 6jm + 2m^2n^2 - 3m^2n + m^2)}{6}), \\ \sum_{k=0}^{n-1} k\mathcal{M}O_{km+j} &= \mathcal{M}D_1 \frac{n(n-1)}{2} + \mathcal{M}D_2 \frac{n(n-1)(3j-m+2mn)}{6} \\ &\quad + \mathcal{M}D_3 \frac{n(n-1)(6j^2 + 8jmn - 4jm + 3m^2n^2 - 3m^2n)}{12}\end{aligned}$$

where

$$\begin{aligned}\mathcal{M}D_1 &= \mathcal{M}O_0, \\ \mathcal{M}D_2 &= \frac{1}{2}(-\mathcal{M}O_2 + 4\mathcal{M}O_1 - 3\mathcal{M}O_0), \\ \mathcal{M}D_3 &= \frac{1}{2}(\mathcal{M}O_2 - 2\mathcal{M}O_1 + \mathcal{M}O_0).\end{aligned}$$

(d):

$$\begin{aligned} \sum_{k=0}^{n-1} \mathcal{M}p_{km+j} &= \sum_{k=0}^{n-1} (n\mathcal{M}E_1 + \mathcal{M}E_2 \frac{n(2j-m+mn)}{2} \\ &\quad + \mathcal{M}E_3 \frac{n(6j^2 + 6jmn - 6jm + 2m^2n^2 - 3m^2n + m^2)}{6}), \\ \sum_{k=0}^{n-1} k\mathcal{M}p_{km+j} &= \mathcal{M}E_1 \frac{n(n-1)}{2} + \mathcal{M}E_2 \frac{n(n-1)(3j-m+2mn)}{6} \\ &\quad + \mathcal{M}E_3 \frac{n(n-1)(6j^2 + 8jmn - 4jm + 3m^2n^2 - 3m^2n)}{12} \end{aligned}$$

where

$$\begin{aligned} \mathcal{M}E_1 &= \mathcal{M}p_0, \\ \mathcal{M}E_2 &= \frac{1}{2}(-\mathcal{M}p_2 + 4\mathcal{M}p_1 - 3\mathcal{M}p_0), \\ \mathcal{M}E_3 &= \frac{1}{2}(\mathcal{M}p_2 - 2\mathcal{M}p_1 + \mathcal{M}p_0). \end{aligned}$$

Next, we present the generating function of the generalized Guglielmo matrix sequence.

THEOREM 2.10. *Let $f_{\mathcal{M}W_n}(x) = \sum_{n=0}^{\infty} \mathcal{M}W_n x^n$ and $A = (a_{ij})_{3 \times 3}$ denote the generating function of generalized Guglielmo matrix sequences. Then,*

$$\begin{aligned} f_{\mathcal{M}W_n}(x) &= \sum_{n=0}^{\infty} \mathcal{M}W_n x^n = \frac{\mathcal{M}W_0 + (\mathcal{M}W_1 - 3\mathcal{M}W_0)x + (\mathcal{M}W_2 - 3\mathcal{M}W_1 + 3\mathcal{M}W_0)x^2}{1 - 3x + 3x^2 - x^3} \quad (2.9) \\ &= \frac{1}{1 - 3x + 3x^2 - x^3} A. \end{aligned}$$

where

$$\begin{aligned} a_{11} &= W_1 + (-3W_1 + W_2)x + W_0x^2, \\ a_{21} &= W_0 + (-3W_0 + W_1)x + 3x^2W_0 + (-3W_1 + W_2)x^2, \\ a_{31} &= (3W_0 - 3W_1 + W_2) + (-8W_0 + 9W_1 - 3W_2)x + (6W_0 - 8W_1 + 3W_2)x^2, \end{aligned}$$

and

$$\begin{aligned} a_{12} &= W_2 - 3W_1 + (W_0 + 6W_1 - 3W_2)x + (-3W_0 + W_1)x^2 \\ a_{22} &= W_1 - 3W_0 + (9W_0 - 6W_1 + W_2)x + (-8W_0 + 9W_1 - 3W_2)x^2 \\ a_{32} &= 9W_1 - 8W_0 - 3W_2 + (21W_0 - 26W_1 + 9W_2)x + (-15W_0 + 21W_1 - 8W_2)x^2 \end{aligned}$$

and

$$\begin{aligned} a_{13} &= W_0 + (-3W_0 + W_1)x + (3W_0 - 3W_1 + W_2)x^2, \\ a_{23} &= (3W_0 - 3W_1 + W_2) + (-8W_0 + 9W_1 - 3W_2)x + (6W_0 - 8W_1 + 3W_2)x^2, \\ a_{33} &= (6W_0 - 8W_1 + 3W_2) + (-15W_0 + 21W_1 - 8W_2)x + (10W_0 - 15W_1 + 6W_2)x^2. \end{aligned}$$

Proof. Using the definition of generalized Guglielmo matrix sequences, and subtracting $xf_{\mathcal{M}W_n}(x)$, $x^2f_{\mathcal{M}W_n}(x)$ and $x^3f_{\mathcal{M}W_n}(x)$ from $f_{\mathcal{M}W_n}(x)$ we get

$$\begin{aligned} (1 - 3x + 3x^2 - x^3)f_{\mathcal{M}W_n}(x) &= \sum_{n=0}^{\infty} \mathcal{M}W_n x^n - 3x \sum_{n=0}^{\infty} \mathcal{M}W_n x^n + 3x^2 \sum_{n=0}^{\infty} \mathcal{M}W_n x^n - x^3 \sum_{n=0}^{\infty} \mathcal{M}W_n x^n \\ &= \sum_{n=0}^{\infty} \mathcal{M}W_n x^n - 3 \sum_{n=0}^{\infty} \mathcal{M}W_n x^{n+1} + 3 \sum_{n=0}^{\infty} \mathcal{M}W_n x^{n+2} - \sum_{n=0}^{\infty} \mathcal{M}W_n x^{n+3} \\ &= \sum_{n=0}^{\infty} \mathcal{M}W_n x^n - 3 \sum_{n=1}^{\infty} \mathcal{M}W_{n-1} x^n + 3 \sum_{n=2}^{\infty} \mathcal{M}W_{n-2} x^n - \sum_{n=3}^{\infty} \mathcal{M}W_{n-3} x^n \\ &= (\mathcal{M}W_0 + \mathcal{M}W_1 x + \mathcal{M}W_2 x^2) - 3(\mathcal{M}W_0 x + \mathcal{M}W_1 x^2) + 3\mathcal{M}W_0 x^2 \\ &\quad + \sum_{n=3}^{\infty} (\mathcal{M}W_n - 3\mathcal{M}W_{n-1} + 3\mathcal{M}W_{n-2} - \mathcal{M}W_{n-3}) x^n \\ &= \mathcal{M}W_0 + \mathcal{M}W_1 x + \mathcal{M}W_2 x^2 - 3\mathcal{M}W_0 x - 3\mathcal{M}W_1 x^2 + 3\mathcal{M}W_0 x^2 \\ &= \mathcal{M}W_0 + (\mathcal{M}W_1 - 3\mathcal{M}W_0)x + (\mathcal{M}W_2 - 3\mathcal{M}W_1 + 3\mathcal{M}W_0)x^2. \end{aligned}$$

Rearranging above equation, we obtain

$$\sum_{n=0}^{\infty} \mathcal{M}W_n x^n = \frac{\mathcal{M}W_0 + (\mathcal{M}W_1 - 3\mathcal{M}W_0)x + (\mathcal{M}W_2 - 3\mathcal{M}W_1 + 3\mathcal{M}W_0)x^2}{1 - 3x + 3x^2 - x^3}$$

which equals the identity that stated in the in the Theorem. This completes the proof. \square

The following corollary gives the generating functions of the triangular, Lucas-triangular, oblong and pentagonal matrix sequences.

COROLLARY 2.11. *The generating functions for the triangular, Lucas-triangular, oblong and pentagonal matrix sequences, respectively, are given as*

(a): *If we take $\mathcal{M}W_n = \mathcal{M}T_n$ in the Theorem 2.10, we get,*

$$\sum_{n=0}^{\infty} \mathcal{M}W_n x^n = \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 1 & x(x-3) & x \\ x & 1-3x & x^2 \\ x^2 & x-3x^2 & 3x^2-3x+1 \end{pmatrix}.$$

(b): *If we take $\mathcal{M}W_n = \mathcal{M}H_n$ in the Theorem 2.10, we get,*

$$\sum_{n=0}^{\infty} \mathcal{M}W_n x^n = \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 3(x-1)^2 & -6(x-1)^2 & 3(x-1)^2 \\ 3(x-1)^2 & -6(x-1)^2 & 3(x-1)^2 \\ 3(x-1)^2 & -6(x-1)^2 & 3(x-1)^2 \end{pmatrix}.$$

(c): If we take $MW_n = MO_n$ in the Theorem 2.10, we get,

$$\sum_{n=0}^{\infty} MW_n x^n = \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 2 & 2x(x - 3) & 2x \\ 2x & 2 - 6x & 2x^2 \\ 2x^2 & -2x(3x - 1) & 6x^2 - 6x + 2 \end{pmatrix}.$$

(d): If we take $MW_n = Mp_n$ in the Theorem 2.10, we get,

$$\sum_{n=0}^{\infty} MW_n x^n = \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 2x + 1 & x^2 - 9x + 2 & x(2x + 1) \\ x(2x + 1) & -6x^2 - x + 1 & 7x^2 - 6x + 2 \\ 7x^2 - 6x + 2 & -19x^2 + 19x - 6 & 15x^2 - 19x + 7 \end{pmatrix}.$$

Next, we give the generating function of the generalized Guglielmo matrix sequences with negative indices.

THEOREM 2.12. Let $g_{MW_n}(x) = \sum_{n=0}^{\infty} MW_{-n}x^n$ be generating function of the generalized Guglielmo matrix sequence. For negative indices, the generating function for the generalized Guglielmo matrix sequence is given as follows

$$\begin{aligned} \sum_{n=0}^{\infty} MW_n x^n &= \frac{MW_0 + (MW_{-1} - 3MW_0)x + (MW_{-2} - 3MW_{-1} + 3MW_0)x^2}{1 - 3x + 3x^2 - x^3} \quad (2.10) \\ &= \frac{1}{1 - 3x + 3x^2 - x^3} B \end{aligned}$$

where $B = (b_{ij})_{3 \times 3}$ and

$$\begin{aligned} b_{11} &= W_1 + (W_0 - 3W_1)x + x^2W_2, \\ b_{21} &= W_0 + (-3W_1 + W_2)x + x^2W_1, \\ b_{31} &= (3W_0 - 3W_1 + W_2) + (-3W_0 + W_1)x + x^2W_0 \end{aligned}$$

and

$$\begin{aligned} b_{12} &= W_2 - 3W_1 + (-3W_0 + 10W_1 - 3W_2)x + (W_0 - 3W_1)x^2, \\ b_{22} &= W_1 - 3W_0 + (W_0 + 6W_1 - 3W_2)x + (-3W_1 + W_2)x^2, \\ b_{32} &= 9W_1 - 8W_0 + -3W_2 + (9W_0 - 6W_1 + W_2)x + (-3W_0 + W_1)x^2 \end{aligned}$$

and

$$\begin{aligned} b_{13} &= W_0 + (-3W_1 + W_2)x + x^2W_1, \\ b_{23} &= (3W_0 - 3W_1 + W_2) + (-3W_0 + W_1)x + x^2W_0, \\ b_{33} &= (6W_0 - 8W_1 + 3W_2) + (-8W_0 + 9W_1 - 3W_2)x + (3W_0 - 3W_1 + W_2)x^2. \end{aligned}$$

Proof. Using the definition of generalized Guglielmo matrix sequences, and subtracting $xg_{\mathcal{M}W_n}(x)$, $x^2g_{\mathcal{M}W_n}(x)$ and $x^3g_{\mathcal{M}W_n}(x)$ from $g_{\mathcal{M}W_n}(x)$ we get

$$\begin{aligned}
 (1 - 3x + 3x^2 - x^3)f_{\mathcal{M}W_{-n}}(x) &= \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^n - 3x \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^n + 3x^2 \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^n - x^3 \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^n \\
 &= \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^n - 3 \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^{n+1} + 3 \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^{n+2} - \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^{n+3} \\
 &= \sum_{n=0}^{\infty} \mathcal{M}W_{-n}x^n - 3 \sum_{n=1}^{\infty} \mathcal{M}W_{-(n-1)}x^n + 3 \sum_{n=2}^{\infty} \mathcal{M}W_{-(n-2)}x^n - \sum_{n=3}^{\infty} \mathcal{M}W_{-(n-3)}x^n \\
 &= (\mathcal{M}W_0 + \mathcal{M}W_{-1}x + \mathcal{M}W_{-2}x^2) - 3(\mathcal{M}W_0x + \mathcal{M}W_{-1}x^2) + 3\mathcal{M}W_0x^2 \\
 &\quad + \sum_{n=3}^{\infty} (3\mathcal{M}W_{-(n-1)} - 3\mathcal{M}W_{-(n-2)} + \mathcal{M}W_{-(n-3)})x^n \\
 &= \mathcal{M}W_0 + \mathcal{M}W_{-1}x + \mathcal{M}W_{-2}x^2 - 3\mathcal{M}W_0x - 3\mathcal{M}W_{-1}x^2 + 3\mathcal{M}W_0x^2 \\
 &= \mathcal{M}W_0 + (\mathcal{M}W_{-1} - 3\mathcal{M}W_0)x + (\mathcal{M}W_{-2} - 3\mathcal{M}W_{-1} + 3\mathcal{M}W_0)x^2.
 \end{aligned}$$

Rearranging above equation, we get

$$\sum_{n=0}^{\infty} \mathcal{M}W_nx^n = \frac{\mathcal{M}W_0 + (\mathcal{M}W_{-1} - 3\mathcal{M}W_0)x + (\mathcal{M}W_{-2} - 3\mathcal{M}W_{-1} + 3\mathcal{M}W_0)x^2}{1 - 3x + 3x^2 - x^3}$$

which equals the identity that stated in the Theorem. \square

The following corollary gives the generating functions of the triangular, Lucas-triangular, oblong and pentagonal matrix sequences with negative subscript .

COROLLARY 2.13. *The generating functions for the triangular, Lucas-triangular, oblong and pentagonal matrix sequences with negative subscript, respectively, are given as*

(a): *If we take $\mathcal{M}W_{-n} = \mathcal{M}T_{-n}$ in the Theorem 2.10, we get,*

$$\sum_{n=0}^{\infty} \mathcal{M}T_{-n}x^n = \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 3x^2 - 3x + 1 & x - 3x^2 & x^2 \\ x^2 & 1 - 3x & x \\ x & x(x - 3) & 1 \end{pmatrix}.$$

(b): *If we take $\mathcal{M}W_{-n} = \mathcal{M}H_{-n}$ in the Theorem 2.10, we have,*

$$\sum_{n=0}^{\infty} \mathcal{M}H_{-n}x^n = \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \\ 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \\ 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \end{pmatrix}.$$

(c): *If we take $\mathcal{M}W_{-n} = \mathcal{M}O_{-n}$ in the Theorem 2.10, we obtain,*

$$\sum_{n=0}^{\infty} \mathcal{M}O_{-n}x^n = \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 6x^2 - 6x + 2 & -2x(3x - 1) & 2x^2 \\ 2x^2 & 2 - 6x & 2x \\ 2x & 2x(x - 3) & 2 \end{pmatrix}.$$

(d): If we take $\mathcal{M}W_{-n} = \mathcal{M}p_{-n}$ in the Theorem 2.10, we get,

$$\sum_{n=0}^{\infty} \mathcal{M}p_{-n}x^n = \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 5x^2 - 3x + 1 & -3x^2 - 5x + 2 & x(x + 2) \\ x(x + 2) & 2x^2 - 9x + 1 & x + 2 \\ x + 2 & x^2 - x - 6 & 2x^2 - 6x + 7 \end{pmatrix}.$$

Now, we give two special equality in the following lemma that we need rest of the study.

LEMMA 2.14. ([??]) We assume that $f(x) = \sum_{n=0}^{\infty} a_n x^n$ is the generating function of the sequence $\{a_n\}_{n \geq 0}$. Then the generating functions of the sequences $\{a_{2n}\}_{n \geq 0}$ and $\{a_{2n+1}\}_{n \geq 0}$ are stated as

$$f_{a_{2n}}(x) = \sum_{n=0}^{\infty} a_{2n}x^n = \frac{f(\sqrt{x}) + f(-\sqrt{x})}{2}$$

and

$$f_{a_{2n+1}}(x) = \sum_{n=0}^{\infty} a_{2n+1}x^n = \frac{f(\sqrt{x}) - f(-\sqrt{x})}{2\sqrt{x}}$$

respectively.

THEOREM 2.15. The generating functions of the sequence $\mathcal{M}W_{2n}$ and $\mathcal{M}W_{2n+1}$ are provided by

$$f_{\mathcal{M}W_{2n}}(x) = \frac{\mathcal{M}W_0 + (\mathcal{M}W_2 - 3\mathcal{M}W_0)x + (6\mathcal{M}W_0 - 8\mathcal{M}W_1 + 3\mathcal{M}W_2)x^2}{1 - 3x + 3x^2 - x^3}, \quad (2.11)$$

$$f_{\mathcal{M}W_{2n+1}}(x) = \frac{\mathcal{M}W_1 + (\mathcal{M}W_0 - 6\mathcal{M}W_1 + 3\mathcal{M}W_2)x + (3\mathcal{M}W_0 - 3\mathcal{M}W_1 + \mathcal{M}W_2)x^2}{1 - 3x + 3x^2 - x^3}. \quad (2.12)$$

Proof. We only proof (2.11). From Theorem (2.10) we can obtain following identities:

$$\begin{aligned} f_{\mathcal{M}W_n}(\sqrt{x}) &= \frac{\mathcal{M}W_0 - \sqrt{x}(\mathcal{M}W_1 - 3\mathcal{M}W_0) + x(3\mathcal{M}W_0 - 3\mathcal{M}W_1 + \mathcal{M}W_2)}{3x + 3\sqrt{x} + x^{\frac{3}{2}} + 1} \\ f_{\mathcal{M}W_n}(-\sqrt{x}) &= -\frac{\mathcal{M}W_0 + \sqrt{x}(\mathcal{M}W_1 - 3\mathcal{M}W_0) + x(3\mathcal{M}W_0 - 3\mathcal{M}W_1 + \mathcal{M}W_2)}{3\sqrt{x} - 3x + x^{\frac{3}{2}} - 1}. \end{aligned}$$

Thereby, using lemma (2.14) identity (2.11) can be proved . The other identity can be found similarly. \square

In the next corollary, we present the generating function of generalized Guglielmo matrix sequences with odd and even subscript by using the Theorem 2.15

COROLLARY 2.16. The triangular, Lucas-triangular, oblong, and pentagonal matrix sequences with even and odd subscript, respectively, are given as

(a): If we take $\mathcal{M}W_n = \mathcal{M}T_n$ in the Theorem 2.15, we get,

$$\begin{aligned} f_{\mathcal{M}T_{2n}}(x) &= \frac{\mathcal{M}T_0 + (\mathcal{M}T_2 - 3\mathcal{M}T_0)x + (6\mathcal{M}T_0 - 8\mathcal{M}T_1 + 3\mathcal{M}T_2)x^2}{1 - 3x + 3x^2 - x^3} \\ &= \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 3x + 1 & -8x & x(x + 3) \\ x(x + 3) & -3x^2 - 6x + 1 & x(3x + 1) \\ x(3x + 1) & -8x^2 & 6x^2 - 3x + 1 \end{pmatrix}, \end{aligned}$$

$$\begin{aligned}
 f_{\mathcal{M}T_{2n+1}}(x) &= \frac{\mathcal{M}T_1 + (\mathcal{M}T_0 - 6\mathcal{M}T_1 + 3\mathcal{M}T_2)x + (3\mathcal{M}T_0 - 3\mathcal{M}T_1 + \mathcal{M}T_2)x^2}{1 - 3x + 3x^2 - x^3} \\
 &= \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} x + 3 & x^2 - 6x - 3 & 3x + 1 \\ 3x + 1 & -8x & x(x + 3) \\ x(x + 3) & -3x^2 - 6x + 1 & x(3x + 1) \end{pmatrix}.
 \end{aligned}$$

(b): If we take $\mathcal{M}W_n = \mathcal{M}H_n$ in the Theorem 2.15, we get,

$$\begin{aligned}
 f_{\mathcal{M}H_{2n}}(x) &= \frac{\mathcal{M}H_0 + (\mathcal{M}H_2 - 3\mathcal{M}H_0)x + (6\mathcal{M}H_0 - 8\mathcal{M}H_1 + 3\mathcal{M}H_2)x^2}{1 - 3x + 3x^2 - x^3} \\
 &= \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \\ 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \\ 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \end{pmatrix},
 \end{aligned}$$

$$\begin{aligned}
 f_{\mathcal{M}H_{2n+1}}(x) &= \frac{\mathcal{M}H_1 + (\mathcal{M}H_0 - 6\mathcal{M}H_1 + 3\mathcal{M}H_2)x + (3\mathcal{M}H_0 - 3\mathcal{M}H_1 + \mathcal{M}H_2)x^2}{1 - 3x + 3x^2 - x^3} \\
 &= \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \\ 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \\ 3(x - 1)^2 & -6(x - 1)^2 & 3(x - 1)^2 \end{pmatrix}.
 \end{aligned}$$

(c): If we take $\mathcal{M}W_n = \mathcal{M}O_n$ in the Theorem 2.15, we get,

$$\begin{aligned}
 f_{\mathcal{M}O_{2n}}(x) &= \frac{\mathcal{M}O_0 + (\mathcal{M}O_2 - 3\mathcal{M}O_0)x + (6\mathcal{M}O_0 - 8\mathcal{M}O_1 + 3\mathcal{M}O_2)x^2}{1 - 3x + 3x^2 - x^3} \\
 &= \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 6x + 2 & -16x & 2x(x + 3) \\ 2x(x + 3) & -6x^2 - 12x + 2 & 2x(3x + 1) \\ 2x(3x + 1) & -16x^2 & 12x^2 - 6x + 2 \end{pmatrix}.
 \end{aligned}$$

$$\begin{aligned}
 f_{\mathcal{M}O_{2n+1}}(x) &= \frac{\mathcal{M}O_1 + (\mathcal{M}O_0 - 6\mathcal{M}O_1 + 3\mathcal{M}O_2)x + (3\mathcal{M}O_0 - 3\mathcal{M}O_1 + \mathcal{M}O_2)x^2}{1 - 3x + 3x^2 - x^3} \\
 &= \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 2x + 6 & 2x^2 - 12x - 6 & 6x + 2 \\ 6x + 2 & -16x & 2x(x + 3) \\ 2x(x + 3) & -6x^2 - 12x + 2 & 2x(3x + 1) \end{pmatrix}.
 \end{aligned}$$

(d): If we take $\mathcal{M}O_n = \mathcal{M}p_n$ in the Theorem 2.15, we get,

$$\begin{aligned}
 f_{\mathcal{M}p_{2n}}(x) &= \frac{\mathcal{M}p_0 + (\mathcal{M}p_2 - 3\mathcal{M}p_0)x + (6\mathcal{M}p_0 - 8\mathcal{M}p_1 + 3\mathcal{M}p_2)x^2}{1 - 3x + 3x^2 - x^3} \\
 &= \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 2x^2 + 9x + 1 & -6x^2 - 20x + 2 & x(7x + 5) \\ x(7x + 5) & -19x^2 - 6x + 1 & 15x^2 - 5x + 2 \\ 15x^2 - 5x + 2 & -38x^2 + 20x - 6 & 26x^2 - 21x + 7 \end{pmatrix}.
 \end{aligned}$$

$$\begin{aligned}
 f_{\mathcal{M}p_{2n+1}}(x) &= \frac{\mathcal{M}p_1 + (\mathcal{M}p_0 - 6\mathcal{M}p_1 + 3\mathcal{M}p_2)x + (3\mathcal{M}p_0 - 3\mathcal{M}p_1 + \mathcal{M}p_2)x^2}{1 - 3x + 3x^2 - x^3} \\
 &= \frac{1}{1 - 3x + 3x^2 - x^3} \begin{pmatrix} 7x + 5 & x^2 - 22x - 3 & 2x^2 + 9x + 1 \\ 2x^2 + 9x + 1 & -6x^2 - 20x + 2 & x(7x + 5) \\ x(7x + 5) & -19x^2 - 6x + 1 & 15x^2 - 5x + 2 \end{pmatrix}.
 \end{aligned}$$

3. Some Identities

In this section we give some identities related to generalized Guglielmo matrix sequences. Moreover, we deal with some special identities such as the Catalan’s identity for the triangular, Lucas-triangular, oblong and pentagonal matrix sequences. First, we investigate relation between $\mathcal{M}H_n$ and $\mathcal{M}W_n$ in the following theorem.

THEOREM 3.1. *For any integer n the following equalities are true:*

- (a): $(W_0 - 2W_1 + W_2)\mathcal{M}H_n = 3\mathcal{M}W_{n+4} - 6\mathcal{M}W_{n+3} + 3\mathcal{M}W_{n+2}$.
- (b): $(W_0 - 2W_1 + W_2)\mathcal{M}H_n = 3\mathcal{M}W_{n+3} - 6\mathcal{M}W_{n+2} + 3\mathcal{M}W_{n+1}$.
- (c): $(W_0 - 2W_1 + W_2)\mathcal{M}H_n = 3\mathcal{M}W_{n+2} - 6\mathcal{M}W_{n+1} + 3\mathcal{M}W_n$.
- (d): $(W_0 - 2W_1 + W_2)\mathcal{M}H_n = 3\mathcal{M}W_{n+1} - 6\mathcal{M}W_n + 3\mathcal{M}W_{n-1}$.
- (e): $(W_0 - 2W_1 + W_2)\mathcal{M}H_n = 3\mathcal{M}W_n - 6\mathcal{M}W_{n-1} + 3\mathcal{M}W_{n-2}$.

Proof. For the prove identity (a) we use the mathematical induction on n . First we derive the proof for $n \geq 0$. If we take $n = 0$, the identity (a) is true. We assume that the identity (a) holds for $n \leq k$ and then ,for $n = k + 1$ and using (2.2) and (1.6), we get

$$\begin{aligned}
 3\mathcal{M}W_{(k+1)+4} - 6\mathcal{M}W_{(k+1)+3} + 3\mathcal{M}W_{(k+1)+2} &= \begin{pmatrix} 3W_0 - 6W_1 + 3W_2 & 3W_0 - 6W_1 + 3W_2 & 3W_0 - 6W_1 + 3W_2 \\ 3W_0 - 6W_1 + 3W_2 & 3W_0 - 6W_1 + 3W_2 & 3W_0 - 6W_1 + 3W_2 \\ 3W_0 - 6W_1 + 3W_2 & 3W_0 - 6W_1 + 3W_2 & 3W_0 - 6W_1 + 3W_2 \end{pmatrix} \\
 &= (W_0 - 2W_1 + W_2) \begin{pmatrix} 3 & 3 & 3 \\ 3 & 3 & 3 \\ 3 & 3 & 3 \end{pmatrix} \\
 &= (W_0 - 2W_1 + W_2)\mathcal{M}H_{k+1}.
 \end{aligned}$$

For the case $n < 0$ the proof can be done in the same way. Consequently the other equalities can be prove similarly. Thus the proof completed. \square

Next, the relation between generalized Guglielmo matrix sequences and oblong matrix sequences.

THEOREM 3.2. *Let n is an arbitrary integer then the following equalities are true:*

- (a): $2\mathcal{M}W_n = (10W_0 - 15W_1 + 6W_2)\mathcal{M}O_{n+4} + (37W_1 - 24W_0 - 15W_2)\mathcal{M}O_{n+3} + (15W_0 - 24W_1 + 10W_2)\mathcal{M}O_{n+2}$.

- (b): $2\mathcal{M}W_n = (6W_0 - 8W_1 + 3W_2)\mathcal{M}O_{n+3} + (21W_1 - 15W_0 - 8W_2)\mathcal{M}O_{n+2} + (10W_0 - 15W_1 + 6W_2)\mathcal{M}O_{n+1}$.
- (c): $2\mathcal{M}W_n = (3W_0 - 3W_1 + W_2)\mathcal{M}O_{n+2} + (9W_1 - 8W_0 - 3W_2)\mathcal{M}O_{n+1} + (6W_0 - 8W_1 + 3W_2)\mathcal{M}O_n$.
- (d): $2\mathcal{M}W_n = W_0O_{n+1} + (W_1 - 3W_0)\mathcal{M}O_n + (3W_0 - 3W_1 + W_2)\mathcal{M}O_{n-1}$.
- (e): $2\mathcal{M}W_n = W_1O_n + (W_2 - 3W_1)\mathcal{M}O_{n-1} + W_0\mathcal{M}O_{n-2}$.
- (f): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}O_n = -2(-3W_1^2 - W_2^2 + W_0W_1 + 3W_1W_2)\mathcal{M}W_{n+4} + 2(-8W_1^2 - 3W_2^2 + 3W_0W_1 - W_0W_2 + 9W_1W_2)\mathcal{M}W_{n+3} + 2(W_0^2 + 9W_1^2 + 3W_2^2 - 6W_0W_1 + 3W_0W_2 - 10W_1W_2)\mathcal{M}W_{n+2}$.
- (g): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}O_n = -2(-W_1^2 + W_0W_2)\mathcal{M}W_{n+3} + 2(W_0^2 - 3W_0W_1 + 3W_0W_2 - W_1W_2)\mathcal{M}W_{n+2} - 2(-3W_1^2 - W_2^2 + W_0W_1 + 3W_1W_2)\mathcal{M}W_{n+1}$.
- (h): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}O_n = 2(W_0^2 + 3W_1^2 - 3W_0W_1 - W_1W_2)\mathcal{M}W_{n+2} - 2(-W_2^2 + W_0W_1 - 3W_0W_2 + 3W_1W_2)\mathcal{M}W_{n+1} - 2(-W_1^2 + W_0W_2)\mathcal{M}W_n$.
- (i): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}O_n = 2(3W_0^2 + 9W_1^2 + W_2^2 - 10W_0W_1 + 3W_0W_2 - 6W_1W_2)\mathcal{M}W_{n+1} - 2(3W_0^2 + 8W_1^2 - 9W_0W_1 + W_0W_2 - 3W_1W_2)\mathcal{M}W_n + 2(W_0^2 + 3W_1^2 - 3W_0W_1 - W_1W_2)\mathcal{M}W_{n-1}$.
- (j): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}O_n = 2(6W_0^2 + 19W_1^2 + 3W_2^2 - 21W_0W_1 + 8W_0W_2 - 15W_1W_2)\mathcal{M}W_n - 2(8W_0^2 + 24W_1^2 + 3W_2^2 - 27W_0W_1 + 9W_0W_2 - 17W_1W_2)\mathcal{M}W_{n-1} + 2(3W_0^2 + 9W_1^2 + W_2^2 - 10W_0W_1 + 3W_0W_2 - 6W_1W_2)\mathcal{M}W_{n-2}$.

For the prove identity (a) we use the mathematical induction on n . First we derive the proof for $n \geq 0$. If we take $n = 0$ the identity (a) is true. We assume that the identity (a) holds for $n \leq k$ and then ,for $n = k + 1$ and using (2.2) and (1.6), we get

$$\begin{pmatrix} (10W_0 - 15W_1 + 6W_2)\mathcal{M}O_{(k+1)+4} + (37W_1 - 24W_0 - 15W_2)\mathcal{M}O_{(k+1)+3} + (15W_0 - 24W_1 + 10W_2)\mathcal{M}O_{(k+1)+2} \\ 2W_{n+1} & -6W_n + 2W_{n-1} & 2W_n \\ 2W_n & -6W_{n-1} + 2W_{n-2} & 2W_{n-1} \\ 2W_{n-1} & -6W_{n-2} + 2W_{n-3} & 2W_{n-2} \end{pmatrix} = 2\mathcal{M}W_{k+1}.$$

For the case $n < 0$ the proof can be done in the same way. Consequently the other equalities can be prove similarly. Thus the proof completed. \square

Next, the relation between generalized Guglielmo Matrix Sequences and pentagonal Matrix Sequences

THEOREM 3.3. *For any integer n the following equalities are true:*

- (a): $27\mathcal{M}W_n = (64W_0 - 89W_1 + 34W_2)\mathcal{M}p_{n+4} + (229W_1 - 158W_0 - 89W_2)\mathcal{M}p_{n+3} + (103W_0 - 158W_1 + 64W_2)\mathcal{M}p_{n+2}$.
- (b): $27\mathcal{M}W_n = (34W_0 - 38W_1 + 13W_2)\mathcal{M}p_{n+3} + (109W_1 - 89W_0 - 38W_2)\mathcal{M}p_{n+2} + (64W_0 - 89W_1 + 34W_2)\mathcal{M}p_{n+1}$.
- (c): $27\mathcal{M}W_n = (13W_0 - 5W_1 + W_2)\mathcal{M}p_{n+2} + (25W_1 - 38W_0 - 5W_2)\mathcal{M}p_{n+1} + (34W_0 - 38W_1 + 13W_2)\mathcal{M}p_n$.
- (d): $27\mathcal{M}W_n = (W_0 + 10W_1 - 2W_2)\mathcal{M}p_{n+1} + (10W_2 - 23W_1 - 5W_0)\mathcal{M}p_n + (13W_0 - 5W_1 + W_2)\mathcal{M}p_{n-1}$.
- (e): $27\mathcal{M}W_n = (7W_1 - 2W_0 + 4W_2)\mathcal{M}p_n + (10W_0 - 35W_1 + 7W_2)\mathcal{M}p_{n-1} + (W_0 + 10W_1 - 2W_2)\mathcal{M}p_{n-2}$.

- (f): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}p_n = (2W_0^2 + 21W_1^2 + 7W_2^2 - 13W_0W_1 + 6W_0W_2 - 23W_1W_2)\mathcal{M}W_{n+4} + (-6W_0^2 + 37W_0W_1 - 19W_0W_2 - 56W_1^2 + 63W_1W_2 - 19W_2^2)\mathcal{M}W_{n+3} + (7W_0^2 + 47W_1^2 + 15W_2^2 - 36W_0W_1 + 19W_0W_2 - 52W_1W_2)\mathcal{M}W_{n+2}$.
- (g): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}p_n = (7W_1^2 + 2W_2^2 - 2W_0W_1 - W_0W_2 - 6W_1W_2)\mathcal{M}W_{n+3} + (W_0^2 - 16W_1^2 - 6W_2^2 + 3W_0W_1 + W_0W_2 + 17W_1W_2)\mathcal{M}W_{n+2} + (2W_0^2 + 21W_1^2 + 7W_2^2 - 13W_0W_1 + 6W_0W_2 - 23W_1W_2)\mathcal{M}W_{n+1}$.
- (h): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}p_n = (W_0^2 + 5W_1^2 - 3W_0W_1 - 2W_0W_2 - W_1W_2)\mathcal{M}W_{n+2} + (2W_0^2 + W_2^2 - 7W_0W_1 + 9W_0W_2 - 5W_1W_2)\mathcal{M}W_{n+1} + (7W_1^2 + 2W_2^2 - 2W_0W_1 - W_0W_2 - 6W_1W_2)\mathcal{M}W_n$.
- (i): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}p_n = (5W_0^2 + 15W_1^2 + W_2^2 - 16W_0W_1 + 3W_0W_2 - 8W_1W_2)W_{n+1} + (-3W_0^2 + 7W_0W_1 + 5W_0W_2 - 8W_1^2 - 3W_1W_2 + 2W_2^2)\mathcal{M}W_n + (W_0^2 + 5W_1^2 - 3W_0W_1 - 2W_0W_2 - W_1W_2)\mathcal{M}W_{n-1}$.
- (j): $(W_0 - 2W_1 + W_2)^3 \mathcal{M}p_n = (12W_0^2 + 37W_1^2 + 5W_2^2 - 41W_0W_1 + 14W_0W_2 - 27W_1W_2)\mathcal{M}W_n + (-14W_0^2 + 45W_0W_1 - 11W_0W_2 - 40W_1^2 + 23W_1W_2 - 3W_2^2)\mathcal{M}W_{n-1} + (5W_0^2 + 15W_1^2 + W_2^2 - 16W_0W_1 + 3W_0W_2 - 8W_1W_2)\mathcal{M}W_{n-2}$.

For the prove identity (a) we use the mathematical induction on n . First we derive the proof for $n \geq 0$. If we take $n = 0$ the identity (a) is true. We assume that the identity (a) holds for $n \leq k$ and then ,for $n = k + 1$ and using (2.2) and (1.6), we get,

$$(64W_0 - 89W_1 + 34W_2)\mathcal{M}p_{k+5} + (229W_1 - 158W_0 - 89W_2)\mathcal{M}p_{k+4} + (103W_0 - 158W_1 + 64W_2)\mathcal{M}p_{k+3} = \begin{pmatrix} 27W_{k+2} & 27W_k & 27W_{k+1} \\ 27W_{k+1} & 27W_{k-1} & 27W_k \\ 27W_k & 27W_{k-2} & 27W_{k-1} \end{pmatrix} = 27\mathcal{M}W_{k+1}.$$

For the case $n < 0$ the proof can be done in the same way. Consequently the other equalities can be prove similarly. Thus the proof completed. \square

THEOREM 3.4. *For all integers m, n the following identity holds:*

$$\mathcal{M}W_{m+n} = T_{m-1}\mathcal{M}W_{n+2} + (T_{m-3} - 3T_{m-2})\mathcal{M}W_{n+1} + T_{m-2}\mathcal{M}W_n.$$

Proof. First, we assume that $m, n \geq 0$. the theorem (3.4) can be proved by mathematical induction on m . If $m = 0$ we get

$$\mathcal{M}W_n = T_{-1}\mathcal{M}W_{n+2} + (T_{-3} - 3T_{-2})\mathcal{M}W_{n+1} + T_{-2}\mathcal{M}W_n$$

which is true since $T_{-1} = 0, T_{-2} = 1, T_{-3} = 3$. We assume that the identity given holds for $m \leq k$. For $m = k + 1$, we get

$$\begin{aligned}
 \mathcal{M}W_{(k+1)+n} &= 3\mathcal{M}W_{n+k} - 3\mathcal{M}W_{n+k-1} + \mathcal{M}W_{n+k-2} \\
 &= 3(T_{k-1}\mathcal{M}W_{n+2} + (T_{k-3} - 3T_{k-2})\mathcal{M}W_{n+1} + T_{k-2}\mathcal{M}W_n) \\
 &\quad - 3(T_{k-2}\mathcal{M}W_{n+2} + (T_{k-4} - 3T_{k-3})\mathcal{M}W_{n+1} + T_{k-3}\mathcal{M}W_n) \\
 &\quad + (T_{k-3}\mathcal{M}W_{n+2} + (T_{k-5} - 3T_{k-4})\mathcal{M}W_{n+1} + T_{k-4}\mathcal{M}W_n) \\
 &= (3T_{k-1} - 3T_{k-2} + T_{k-3})\mathcal{M}W_{n+2} + ((3T_{k-3} - 3T_{k-4} + T_{k-5}) \\
 &\quad - 3(3T_{k-2} - 3T_{k-3} + T_{k-4}))\mathcal{M}W_{n+1} + (3T_{k-2} - 3T_{k-3} + T_{k-4})\mathcal{M}W_n \\
 &= T_k\mathcal{M}W_{n+2} + (T_{k-2} - 3T_{k-1})\mathcal{M}W_{n+1} + T_{k-1}\mathcal{M}W_n \\
 &= T_{(k+1)-1}\mathcal{M}W_{n+2} + (T_{(k+1)-3} - 3T_{(k+1)-2})\mathcal{M}W_{n+1} + T_{(k+1)-2}\mathcal{M}W_n.
 \end{aligned}$$

Consequently, by mathematical induction on m , this proves Theorem 3.4. The case $m, n < 0$ can be proved similarly. \square

For $n \geq 0$, $m \geq 0$ and taking $\mathcal{M}W_n = \mathcal{M}T_n$ or $\mathcal{M}W_n = \mathcal{M}H_n$ or $\mathcal{M}W_n = \mathcal{M}O_n$ or $\mathcal{M}W_n = \mathcal{M}p_n$, respectively, we get,

$$\begin{aligned}
 \mathcal{M}T_{m+n} &= T_{m-1}\mathcal{M}T_{n+2} + (T_{m-3} - 3T_{m-2})\mathcal{M}T_{n+1} + T_{m-2}\mathcal{M}T_n, \\
 \mathcal{M}H_{m+n} &= T_{m-1}\mathcal{M}H_{n+2} + (T_{m-3} - 3T_{m-2})\mathcal{M}H_{n+1} + T_{m-2}\mathcal{M}H_n, \\
 \mathcal{M}O_{m+n} &= T_{m-1}\mathcal{M}O_{n+2} + (T_{m-3} - 3T_{m-2})\mathcal{M}O_{n+1} + T_{m-2}\mathcal{M}O_n, \\
 \mathcal{M}p_{m+n} &= T_{m-1}\mathcal{M}p_{n+2} + (T_{m-3} - 3T_{m-2})\mathcal{M}p_{n+1} + T_{m-2}\mathcal{M}p_n.
 \end{aligned}$$

Next Lemma, we give some identities related to $\mathcal{M}A_1, \mathcal{M}A_2, \mathcal{M}A_3, \mathcal{M}B_1, \mathcal{M}B_2, \mathcal{M}B_3, \mathcal{M}C_1, \mathcal{M}C_2, \mathcal{M}C_3, \mathcal{M}D_1, \mathcal{M}D_2, \mathcal{M}D_3, \mathcal{M}E_1, \mathcal{M}E_2$ and $\mathcal{M}E_3$.

LEMMA 3.5. *The following identities holds:*

$$\begin{aligned}
 \mathcal{M}B_1\mathcal{M}A_1 &= \mathcal{M}B_1\mathcal{M}C_1 = \mathcal{M}B_1\mathcal{M}D_1 = \mathcal{M}B_1\mathcal{M}E_1 \\
 \mathcal{M}B_3^2 &= \mathcal{M}B_2\mathcal{M}B_3 = \mathcal{M}B_3\mathcal{M}B_2 = 0 \\
 \mathcal{M}B_2\mathcal{M}C_2 &= \mathcal{M}C_2\mathcal{M}B_2 = \mathcal{M}E_2\mathcal{M}C_2 = \mathcal{M}C_2\mathcal{M}E_2 = 0 \\
 \mathcal{M}B_3\mathcal{M}E_3 &= \mathcal{M}E_3\mathcal{M}B_3 = \mathcal{M}B_3\mathcal{M}D_3 = \mathcal{M}D_3\mathcal{M}B_3 = 0
 \end{aligned}$$

Proof. Using matrix multiplication the proof is done. \square

Note that $\mathcal{M}B_1 = \mathcal{M}T_0$ is identity matrix.

Next Lemma, we give some relation linked to Guglielmo matrix sequences.

LEMMA 3.6. *For all integers m and n , we have the following identities.*

- (a): $\mathcal{M}T_0\mathcal{M}W_n = \mathcal{M}W_n\mathcal{M}T_0 = \mathcal{M}W_n$.
- (b): $\mathcal{M}W_0\mathcal{M}T_n = \mathcal{M}T_n\mathcal{M}W_0 = \mathcal{M}W_n$.
- (c): $\mathcal{M}T_n\mathcal{M}T_m = \mathcal{M}T_m\mathcal{M}T_n = \mathcal{M}T_{n+m}$.
- (d): $\mathcal{M}T_m\mathcal{M}W_n = \mathcal{M}W_n\mathcal{M}T_m = \mathcal{M}W_{m+n}$.
- (e): $\mathcal{M}T_m\mathcal{M}H_n = \mathcal{M}H_n\mathcal{M}T_m = \mathcal{M}H_{m+n}$.
- (f): $\mathcal{M}T_m\mathcal{M}O_n = \mathcal{M}O_n\mathcal{M}T_m = \mathcal{M}O_{m+n}$.
- (g): $\mathcal{M}T_m\mathcal{M}p_n = \mathcal{M}p_n\mathcal{M}T_m = \mathcal{M}p_{m+n}$.
- (h): $\mathcal{M}W_0\mathcal{M}W_n = \mathcal{M}W_n\mathcal{M}W_0$.
- (i): $\mathcal{M}W_n\mathcal{M}W_m = \mathcal{M}W_m\mathcal{M}W_n = \mathcal{M}W_0\mathcal{M}W_{n+m}$.
- (j): $\mathcal{M}T_{-n} = (\mathcal{M}T_n)^{-1}$.
- (k): $\mathcal{M}W_{-n} = (\mathcal{M}W_0)^{1-n}(\mathcal{M}W_{-1})^n$.

Proof. Identities can be established easily.

- (a): The proof can be easily seen since $\mathcal{M}T_0$ is the identity matrix.
- (b): The proof can be seen by using Theorem 2.3 and Corollary 2.4.
- (c): Using Lemma 3.5 we have the following equalities,

$$\begin{aligned}
 \mathcal{M}T_n\mathcal{M}T_m &= (\mathcal{M}B_1 + \mathcal{M}B_2n + \mathcal{M}B_3n^2)(\mathcal{M}B_1 + \mathcal{M}B_2m + \mathcal{M}B_3m^2) \\
 &= \mathcal{M}B_1^2 + \mathcal{M}B_1\mathcal{M}B_2m + \mathcal{M}B_1\mathcal{M}B_3m^2 + \mathcal{M}B_1\mathcal{M}B_2n + \mathcal{M}B_2\mathcal{M}B_2mn + \mathcal{M}B_2\mathcal{M}B_3nm^2 + \mathcal{M}B_1\mathcal{M}B_3n^2 + \\
 &= (\mathcal{M}B_1 + \mathcal{M}B_2(n+m) + \mathcal{M}B_3(n+m)^2) \\
 &= \mathcal{M}T_{n+m}.
 \end{aligned}$$

The identity $\mathcal{M}T_m\mathcal{M}T_n = \mathcal{M}T_{n+m}$ can be seen similarly. This completes the proof that we need.

- (d): From (b) and (c), we have

$$\begin{aligned}
 \mathcal{M}T_m\mathcal{M}W_n &= \mathcal{M}T_m\mathcal{M}T_n\mathcal{M}W_0 \\
 &= \mathcal{M}T_{n+m}\mathcal{M}W_0 \\
 &= \mathcal{M}W_{m+n}.
 \end{aligned}$$

- (e): Take $\mathcal{M}W_n = \mathcal{M}H_n$ in (d).
- (f): Take $\mathcal{M}W_n = \mathcal{M}O_n$ in (d).
- (g): Take $\mathcal{M}W_n = \mathcal{M}p_n$ in (d).
- (h): If we perform matrix multiplication and compare the entries of the resulting rows and columns, the required proof is complete.

(i): If we use (d) and (h) and (b), we have

$$\begin{aligned} \mathcal{M}W_0\mathcal{M}W_{n+m} &= \mathcal{M}W_0\mathcal{M}W_n\mathcal{M}T_m \\ &= \mathcal{M}W_n\mathcal{M}W_0\mathcal{M}T_m \\ &= \mathcal{M}W_n\mathcal{M}W_m \end{aligned}$$

The identity $\mathcal{M}W_0\mathcal{M}W_{n+m} = \mathcal{M}W_m\mathcal{M}W_n$ can be seen similarly. This completes the proof of (h).

(j): First, we establish the proof the case $n \geq 0$ by using mathematical induction on n . If $n = 0$ then we have

$$\mathcal{M}T_0 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}^{-1} = (\mathcal{M}T_0)^{-1}$$

and

$$\mathcal{M}T_{-1} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -3 & 3 \end{pmatrix} = \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -3 & 3 \end{pmatrix}^{-1} = (\mathcal{M}T_{-1})^{-1}.$$

Let the equality given in the (j) holds for $n \leq k$. For $n = k + 1$, by using (c), we obtain

$$\begin{aligned} (\mathcal{M}T_{k+1})^{-1} &= (\mathcal{M}T_k\mathcal{M}T_{-1})^{-1} = (\mathcal{M}T_{-1})^{-1}(\mathcal{M}T_k)^{-1} = \mathcal{M}T_{-1}\mathcal{M}T_{-k} \\ &= \begin{pmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \\ 1 & -3 & 3 \end{pmatrix} \begin{pmatrix} T_{-n+1} & -3T_{-n} + T_{-n-1} & T_{-n} \\ T_{-n} & -3T_{-n-1} + T_{-n-2} & T_{-n-1} \\ T_{-n-1} & -3T_{-n-2} + T_{-n-3} & T_{-n-2} \end{pmatrix} \\ &= \begin{pmatrix} T_{-k} & -3T_{-k-1} + T_{-k-2} & T_{-k-1} \\ T_{-k-1} & -3T_{-k-2} + T_{-k-3} & T_{-k-2} \\ 3T_{-k-1} - 3T_{-k} + T_{-k+1} & 10T_{-k-1} - 12T_{-k-2} + 3T_{-k-3} - 3T_{-k} & 3T_{-k-2} - 3T_{-k-1} + T_{-k} \end{pmatrix} \\ &= \begin{pmatrix} T_{-(k+1)+1} & -3T_{-(k+1)} + T_{-(k+1)-1} & T_{-(k+1)} \\ T_{-(k+1)} & -3T_{-(k+1)-1} + T_{-(k+1)-2} & T_{-(k+1)-1} \\ T_{-(k+1)-1} & -3T_{-(k+1)-2} + T_{-(k+1)-3} & T_{-(k+1)-2} \end{pmatrix} \\ &= \mathcal{M}T_{-(k+1)}. \end{aligned}$$

Thus, by induction on n , this proves (j) for $n \geq 0$. For the other case $n < 0$ the proof can be done similarly. This completes proof.

(k): If we take $m = -n + 1$ and $n = -1$ in the identity (i), we obtain that

$$\mathcal{M}W_0\mathcal{M}W_{-n} = \mathcal{M}W_{-n+1}\mathcal{M}W_{-1}. \quad (3.1)$$

If we multiply both side of the equation that we obtain above with $\mathcal{M}W_0$ and using (i) we have the relation

$$\begin{aligned} \mathcal{M}W_0\mathcal{M}W_0\mathcal{M}W_{-n} &= \mathcal{M}W_0\mathcal{M}W_{-n+1}\mathcal{M}W_{-1} \\ &= \mathcal{M}W_{-n+1}\mathcal{M}W_{-1}\mathcal{M}W_{-1}. \end{aligned}$$

Repeating this process then we obtain

$$(\mathcal{M}W_0)^{n-1}\mathcal{M}W_{-n} = (\mathcal{M}W_{-1})^n.$$

Thus, it follows that

$$\mathcal{M}W_{-n} = (\mathcal{M}W_0)^{1-n}(\mathcal{M}W_{-1})^n.$$

This completes the proof. \square

Note that using Lemma 3.6 (k) and (d), we obtain

$$\begin{aligned} \mathcal{M}W_{-n} &= (\mathcal{M}W_0)^{1-n}(\mathcal{M}W_{-1})^n \\ &= (\mathcal{M}W_n\mathcal{M}W_{-n})^{1-n}(\mathcal{M}W_{-1})^n \\ &= (\mathcal{M}W_{-n})^{1-n}(\mathcal{M}W_n)^{1-n}(\mathcal{M}W_{-1})^n \end{aligned}$$

and then by Lemma (j), we get

$$\mathcal{M}W_{-n} = (\mathcal{M}T_n)^{1-n}(\mathcal{M}W_n)^{1-n}(\mathcal{M}W_{-1})^n.$$

Using Lemma 3.6 and comparing matrix entries, we obtain next result.

THEOREM 3.7. *For all integers m, n and r , the following identities hold:*

- (a): $\mathcal{M}T_n^m = \mathcal{M}T_{mn}$,
- (b): $\mathcal{M}T_{n+1}^m = \mathcal{M}T_1^m\mathcal{M}T_{mn}$,
- (c): $\mathcal{M}T_{n-r}\mathcal{M}T_{n+r} = \mathcal{M}T_n^2 = \mathcal{M}T_2^n$.

Proof. We prove for $m, n, r \geq 0$. The other cases can be proved similarly.

- (a): We can write \mathcal{N}_n^m as

$$\mathcal{M}T_n^m = \mathcal{M}T_n\mathcal{M}T_n\dots\mathcal{M}T_n \text{ (} m \text{ times)}.$$

Using Lemma 3.6 (c) iteratively, we obtain the required result:

$$\begin{aligned}
 \mathcal{M}T_n^m &= \underbrace{\mathcal{M}T_n \mathcal{M}T_n \dots \mathcal{M}T_n}_{m \text{ times}} \\
 &= \mathcal{M}T_{2n} \underbrace{\mathcal{M}T_n \mathcal{M}T_n \dots \mathcal{M}T_n}_{m-1 \text{ times}} \\
 &= \mathcal{M}T_{3n} \underbrace{\mathcal{M}T_n \mathcal{M}T_n \dots \mathcal{M}T_n}_{m-2 \text{ times}} \\
 &\vdots \\
 &= \mathcal{M}T_{(m-1)n} \mathcal{M}T_n \\
 &= \mathcal{M}T_{mn}.
 \end{aligned}$$

(b): As a similar approach in (a) we have

$$\mathcal{M}T_{n+1}^m = \mathcal{M}T_{n+1} \mathcal{M}T_{n+1} \dots \mathcal{M}T_{n+1} = \mathcal{M}T_{m(n+1)} = \mathcal{M}T_m \mathcal{M}T_{mn} = \mathcal{M}T_1 \mathcal{M}T_{m-1} \mathcal{M}T_{mn}.$$

Using Lemma 3.6 (c), we can write iteratively,

$$\begin{aligned}
 \mathcal{M}T_m &= \mathcal{M}T_1 \mathcal{M}T_{m-1}, \\
 \mathcal{M}T_{m-1} &= \mathcal{M}T_1 \mathcal{M}T_{m-2}, \\
 &\vdots \\
 \mathcal{M}T_2 &= \mathcal{M}T_1 \mathcal{M}T_1.
 \end{aligned}$$

Now it follows that

$$\begin{aligned}
 \mathcal{M}T_{n+1}^m &= \mathcal{M}T_1 \mathcal{M}T_{m-1} \mathcal{M}T_{mn} \\
 &= \underbrace{\mathcal{M}T_1 \mathcal{M}T_1 \dots \mathcal{M}T_1}_{m \text{ times}} \mathcal{M}T_{mn} = \mathcal{M}T_1^m \mathcal{M}T_{mn}.
 \end{aligned}$$

(c): Lemma 3.6 (c) gives

$$\mathcal{M}T_{n-r} \mathcal{M}T_{n+r} = \mathcal{M}T_{2n} = \mathcal{M}T_n \mathcal{M}T_n = \mathcal{M}T_n^2$$

and also

$$\mathcal{M}T_{n-r} \mathcal{M}T_{n+r} = \mathcal{M}T_{2n} = \underbrace{\mathcal{M}T_2 \mathcal{M}T_2 \dots \mathcal{M}T_2}_{n \text{ times}} = \mathcal{M}T_2^n.$$

Next Theorem, we encounter similar outcomes for the matrix sequence $\mathcal{M}W_n$.

THEOREM 3.8. *For all integers m, n and r , the following identities hold:*

(a): $\mathcal{M}W_{n-r} \mathcal{M}W_{n+r} = \mathcal{M}W_n^2,$

(b): $\mathcal{M}W_n^m = \mathcal{M}W_0^m \mathcal{M}T_{mn}.$

Proof.

(a): Using Binet’s formula of generalized Guglielmo sequence given in Theorem 2.6 and matrix multiplication, we obtain

$$\begin{aligned} \mathcal{M}W_{n-r}\mathcal{M}W_{n+r} - \mathcal{M}W_n^2 &= (\mathcal{M}A_1 + \mathcal{M}A_2(n+r) + \mathcal{M}A_3(n+r)^2)(\mathcal{M}A_1 + \mathcal{M}A_2(n-r) \\ &\quad + \mathcal{M}A_3(n-r)^2) - (\mathcal{M}A_1 + \mathcal{M}A_2n + \mathcal{M}A_3n^2)^2 \\ &= -2n^2r^2\mathcal{M}A_3^2 - 2nr^2\mathcal{M}A_2\mathcal{M}A_3 + r^4\mathcal{M}A_3^2 - r^2\mathcal{M}A_2^2 + 2\mathcal{M}A_1r^2\mathcal{M}A_3 \\ &= 0. \end{aligned}$$

At least, we get the result as required.

(b): By Theorem 3.7, we have

$$\mathcal{M}W_0^m \mathcal{M}T_{mn} = \underbrace{\mathcal{M}W_0\mathcal{M}W_0\dots\mathcal{M}W_0}_{m \text{ times}} \underbrace{\mathcal{M}T_n\mathcal{M}T_n\dots\mathcal{M}T_n}_{m \text{ times}}.$$

By applying Lemma 3.6 (b) iteratively, it follows that

$$\begin{aligned} \mathcal{M}W_0^m \mathcal{M}T_{mn} &= (\mathcal{M}W_0\mathcal{M}T_n)(\mathcal{M}W_0\mathcal{M}T_n)\dots(\mathcal{M}W_0\mathcal{M}T_n) \\ &= \mathcal{M}W_n\mathcal{M}W_n\dots\mathcal{M}W_n = \mathcal{M}W_n^m. \end{aligned}$$

This completes the proof. \square

4. Conclusion

In this study we define generalized Guglielmo matrix sequence and give four special cases that we named triangular matrix sequence, Lucas-triangular matrix sequence, oblong matrix sequence and pentagonal matrix sequence. We also define the Binet’s formula of the generalized Guglielmo matrix sequence and its special cases. Moreover, we give generating function and sum formulas related to generalized Guglielmo matrix sequence. Furthermore, we give some lemma and theorem using matrix properties and some formulas. We know that the matrix sequences are fundamental issue in linear algebra and have applications in various fields including numerical analysis, differential equations, control theory, and computer graphics. Thus, we believe that our study will contribute to ongoing research in this field.

References

- [1] Cerda-Morales, G., On the Third-Order Jacobsthal and Third-Order Jacobsthal–Lucas Sequences and Their Matrix Representations. *Mediterranean Journal of Mathematics*, **16** (2019) 1-12.
- [2] Civciv, H., Turkmen, R., On the (s; t)-Fibonacci and Fibonacci matrix sequences, *Ars Combin.* **87** (2008) 161-173.
- [3] Civciv, H., Turkmen, R., Notes on the (s; t)-Lucas and Lucas matrix sequences, *Ars Combin.* **89** (2008) 271-285.
- [4] Gulec, H.H., Taskara, N., On the (s; t)-Pell and (s; t)-Pell-Lucas sequences and their matrix representations, *Appl. Math. Lett.* **25** (2012), 1554-1559, doi.org/10.1016/j.aml.2012.01.014.

- [5] Soykan, Y., Matrix Sequences of Tetranacci and Tetranacci-Lucas Numbers, *Int. J. Adv. Appl. Math. and Mech.* 7(2), 57-69, 2019.
- [6] Soykan, Y., Matrix Sequences of Tribonacci and Tribonacci-Lucas Numbers, *Communications in Mathematics and Applications*, 11(2), 281-295, 2020. DOI: 10.26713/cma.v11i2.1102
- [7] Soykan, Y., Tribonacci and Tribonacci-Lucas Matrix Sequences with Negative Subscripts, *Communications in Mathematics and Applications*, 11(1), 141159, 2020. DOI: 10.26713/cma.v11i1.1103.
- [8] Soykan, Y., Göcen, M., Çevikel, S., On Matrix Sequences of Narayana and Narayana-Lucas Numbers, *Karaelmas Science and Engineering Journal*, 11(1), 83-90, 2021. DOI: 10.7212/karaelmasfen.824911.
- [9] Soykan, Y., Taşdemir, E., İrge, V., A Study on Matrix Sequence of Adjusted Tribonacci-Lucas Numbers, *J. Int. Math. Virtual Inst.*, in Print.
- [10] Soykan, Y., Generalized Guglielmo Numbers: An Investigation of Properties of Triangular, Oblong and Pentagonal Numbers via Their Third Order Linear Recurrence Relations, *Earthline Journal of Mathematical Sciences*, 9(1), 1-39, 2022.
- [11] Uslu, K., Uygun, Ş., On the (s,t) Jacobsthal and (s,t) Jacobsthal-Lucas Matrix Sequences, *Ars Combin.* **108** (2013), 13-22.
- [12] Uygun, Ş., Uslu, K., (s,t) -Generalized Jacobsthal Matrix Sequences, *Springer Proceedings in Mathematics & Statistics, Computational Analysis*, Amat, Ankara, (May 2015), 325-336.
- [13] Uygun, Ş., Some Sum Formulas of (s,t) -Jacobsthal and (s,t) -Jacobsthal Lucas Matrix Sequences, *Applied Mathematics*, **7** (2016), 61-69, <http://dx.doi.org/10.4236/am.2016.71005>.
- [14] Uygun, Ş., The binomial transforms of the generalized (s,t) -Jacobsthal matrix sequence, *Int. J. Adv. Appl. Math. and Mech.* 6(3) (2019), 14–20.
- [15] Yazlik, Y., Taskara, N., Uslu K., Yilmaz, N., The generalized $(s; t)$ -sequence and its matrix sequence, *Am. Inst. Phys. (AIP) Conf. Proc.* 1389 (2012), 381-384, <https://doi.org/10.1063/1.3636742>.
- [16] Yilmaz, N., Taskara, N., Matrix Sequences in Terms of Padovan and Perrin Numbers, *Journal of Applied Mathematics*, Volume 2013 (2013), Article ID 941673, 7 pages, <http://dx.doi.org/10.1155/2013/941673>.
- [17] Yilmaz, N., Taskara, N., On the Negatively Subscripted Padovan and Perrin Matrix Sequences, *Communications in Mathematics and Applications*, **5**(2) (2014), 59-72.
- [18] Wani, A.A., Badshah, V.H., and Rathore, G.B.S., Generalized Fibonacci and k -Pell Matrix Sequences, *Punjab University Journal of Mathematics*, **50**(1) (2018), 68-79.