

Original Research Article

Effect of different planting dates and spacing on growth parameters of cabbage (*Brassica oleracea* var. *capitata* L.)

Comment [u1]:

Comment [u2]: Use Sentence case, font style should be Times new Roman

Abstract

Aim: To investigate the impact of various planting dates and spacing on cabbage growth.

Study Design: The experimental design Factorial Randomized Block Design (FRBD) was used.

Place and Duration of study: An experiment was conducted in the rabi seasons of 2021–22 and 2022–23 at the Horticulture Research Farm, Babasaheb Bhimrao Ambedkar University, Lucknow (U.P.)

Methodology: There were twelve treatments with three replications. The variety selected for experiment was Pusa Mukta.

Results: The combined data of two years showed that treatment P₁ (30 November) had significantly higher plant survivability (95.28%), plant height (14.67 cm, 21.31 cm and 28.39 cm) at 30, 45 and 60 DAT, respectively, stem diameter (2.11 cm), no. of non-wrapper leaves (16.28), no. of wrapper leaves (27.19), plant spread E-W (49.40 cm) and plant spread N-S (52.63 cm). Among spacings, The findings showed that the S₄ (60 cm x 60 cm) plant spacing achieved maximum plant survivability at 91.59%. It also led to the greatest plant height at 30, 45 and 60 days after transplanting (DAT) with measurements of 15.73 cm, 22.45 cm and 31.12 cm, respectively. Additionally, this spacing resulted in the largest stem diameter (2.31 cm), the highest no. of non-wrapper leaves (17.40), no. of wrapper leaves (28.30), greatest plant spread in both East-West (52.79 cm) and North-South (56.08 cm) directions.

Conclusion: The combined data of two years showed that treatment P₁ (30 November) and Spacing 60 cm x 60 cm had significantly higher growth parameters.

Keywords: Planting dates, plant spacing, cabbage and growth parameters.

Comment [u3]: Abstract should be single line spaced, blocked and should not be divided into sub headings. It should begin with statement of the problem and study design and place and duration of study together with details of the treatments and their combination should come as methodology. It should also explain Data collection procedure and the parameters involved, Data analysis and the tool used for the analysis together with means separation. There should be suggestion/recommendation after conclusion

1. INTRODUCTION

Cabbage (*Brassica oleracea* var. *capitata* L.) is among the most popular cole crops which is consumed globally, related to Brassicaceae (Cruciferae) family & has chromosome number $2n=2x=18$. The "head," or edible part of the cabbage plant, is composed of many smooth, thick leaves

Comment [u4]: Remove number

Comment [u5]:

that overlap one another. Cabbage varieties vary widely in terms of colour (from green to purple), leaf characteristics (from smooth to savoy), head form (from flat to pointed), and maturity (from early to late maturing). The most prevalent kinds are the green, round-headed ones (Splittstoesser, 1979 and Phillips and Rix, 1993). All of these crops descended from the wild cabbage, which was originally from the Mediterranean region and gave rise to a wide range of cultivars, including *B. oleracea* vars. *capitata* and *botrytis* (Kochhar, 2010; Singh and Sharma, 2001). India and other countries grow a lot of cabbage. It came to India during fifteenth century from Portugal (Singh *et al.*, 2004). Crop yield is determined by a multitude of factors, two of the most critical ones being plant population and the ideal planting season. The current cropping pattern and the surrounding conditions determine the ideal planting timing. Cool temperatures are necessary for cabbage to grow to its full potential and form a head. The timing of planting is crucial, and meticulous seed sowing is required to ensure that the crop maximises the benefits of the prolonged chilled season. It is discovered that planting early produced larger and more uniformly distributed plants, primarily due to the longer daylight hours and more consistent temperatures. Plant morphology is altered in a variety of ways by competition brought through different spacing. In order to maximize the use of natural resources like nutrients, sunlight, soil moisture etc., as well as achieve a satisfactory yield and appropriate land use, it is important to maintain optimal plant spacing (Ullah *et al.*, 2013). Crop spacing can vary based on climate, soil fertility and the suitability of different cultivars for a given area. Lowest plant density generally results in more robust plants with larger leaves, likely because there is less rivalry for nutrition, light and water in comparison to highest plant density (Bairwa *et al.*, 2017). Although high population densities are often used for various crops to enhance vegetable production, closer spacing can provide benefits such as better soil protection, improved weed control, more efficient use of fertilizers and increased yields (Neto *et al.*, 2016).

Comment [u6]: Are there no recent experiments results?

2. MATERIALS AND METHODS

The research trial conducted at Horticulture Farm of Babasaheb Bhimrao Ambedkar University in Lucknow, Uttar Pradesh, India. The site is located at 80° 92' East longitude, 26° 76' North latitude, and 123 meters above MSL. The trial took place during the rabi seasons of 2021-22 and 2022-23. The soil at the site was sandy loam with a slightly alkaline pH of 8.5, and comprised of 83.57 kg/ha of available Nitrogen, 14.40 kg/ha of available Phosphorus and 140.21 kg/ha of available Potash. The seedbed was ~~one meter long and two meters~~ 1m × 2m wide, and the seeds used were of the Pusa Mukta cabbage variety was used. For transplanted the seeds were evenly spaced at a distance of 5 cm, and they were seeded three times according to treatments on the seedbed in each year to preserve the same age. After sowing the seeds at 2 cm depth, they were lightly watered and covered with a fine layer of dirt. One month's old healthy young plants transferred on 30 November (P₁), 7 December (P₂) and 14 December (P₃) at four spacings viz. 45 cm x 30 cm (S₁), 45 cm x 45 cm (S₂), 60 cm x 45 cm (S₃) and 60 cm x 60 cm (S₄). The trial was conducted in Factorial RBD. There were twelve treatments with three replications. The Farm Yard Manure was mixed in the soil at the rate 20 kg/ha fifteen days before transplantation. The recommended fertilizer dose of 180 kg N, 120 kg P and 100 kg K per hectare was applied using Urea, DAP & MOP respectively as basal dose. During transplantation, a half quantity of N applied to the soil; the second dose was divided in equivalent

Comment [u7]: Introduction should be rounded up by statement of problem and objectives of the study

amounts i.e., 1st top dressing applied at 30 DAT and 2nd at 45 DAT. All of the suggested practices were implemented throughout the duration of the cropping season. In each plot, five plants were chosen at random, and data was collected regarding the plants' survival, height (30, 45, and 60 DAT), stem diameter, number of wrapper and non-wrapper leaves, plant spread E-W, and plant spread N-S. As indicated by Panse and Sukhatme (1985), the conventional technique was followed for statistical analysis of the growth parameters measurements from various sets of the trials. After analysis, the data were shown at a significance level of 5 percent.

Comment [u8]: This should be separated as fertilizer application

Comment [u9]: Management practices observed should be stated clearly

Comment [u10]: INSTEAD: Data were collected from the five randomly selected and tagged plants on each plot

Comment [u11]: Data collected at what interval?

Comment [u12]: State the statistical software and the version

3. RESULTS AND DISCUSSION

3.1 Effect of planting dates on growth parameters of cabbage

3.1.1 Plant survivability

Planting dates had a major effect on the vegetative characters. The data on a pooled average basis (Table 1 and Fig. 1) showed that the planting date of 30 November (P₁), had the significantly higher plant survival (95.28%), while the planting date of 14 December, (P₃), had the lowest plant survival (84.81%). Early planting usually results in a longer growing season for the plant, which might enhance establishment and resilience. Plants benefit from positive weather conditions because they have more time to develop robust root systems before adverse ones appear. These results support Yasmin's (2008) findings in potatoes.

3.1.2 Plant height

It is revealed from Table 1 and Fig. 1 that the maximum plant height at 30 DAT (14.67 cm), 45 DAT (21.31 cm), and 60 DAT (28.39 cm) were reported under the planting date on 30 November (P₁). The minimum plant height at 30 DAT (13.34 cm), 45 DAT (18.17 cm), and 60 DAT (25.22 cm) were reported under planting date on 14 December (P₃). Temperature variations were the primary cause of changes in plant height, however favourable weather throughout growth period, may also have played a role. These findings validate observations of Saikia *et al.*, (2010) and Kanase *et al.*, (2018) in broccoli, Yadav *et al.*, (2021) in cabbage.

3.1.3 Stem diameter

The data on pooled mean basis revealed that stem diameter significantly influenced by planting time (Table 1 and Fig. 1). The maximum stem diameter (2.11 cm) was recorded from the plants which were transplanted on 30 November (P₁) while the minimum stem diameter (1.81 cm) recorded from the planting dates on 14 December (P₃). When comparing the vegetative development phase to an earlier planting date, the minimal diameter of the stem may be the result of decreased in both night and day average temperatures. These findings concur with Abed *et al.*, (2015) in cabbage, Singhal *et al.*, (2009) and Kanase *et al.*, (2018) in broccoli.

Comment [u13]: et al should come with full stop and comma only if the name and year of the cited reference are both inside the brackets. If only the year is inside, it should be et al. only

3.1.4 No. of Non-Wrapper Leaves and Wrapper Leaves

It is revealed on the basis of pooled data of two years that number of non-wrappers leaves significantly influenced by distinct planting dates. The maximal no. of non-wrapper leaves (16.28) was counted from the planting dates on 30 November (P₁) and the least number of non-wrapper leaves (13.94) were counted in plant that are planted on 14 December (P₃).

Comment [u14]: this should be written in full

Number of wrapper leaves were significantly affected by various the planting dates used. The data according to average basis are shown in Table 1 and graphically illustrated in Fig. 1. The transplanting date i.e., of 30th November (P₁) performed better having maximum wrapper leaves (27.19) as when compared to other dates while a smaller minimum number of wrapper leaves (25.36) were found recorded under the planting date of 14th December (P₃). This is because the plants have might be as a result of optimum climatic conditions that favours cabbage growth and a longer period to grow before unfavourable weather conditions impact their development. The outcomes of the experiments is in line with the findings of are confined along with the studies of Ullah *et al.*, (2013) in cabbage.

3.1.5 Plant Spread (East-West and North-South)

The timing of planting significantly impacts the canopy area of cabbage plants in both of north to south and east to west orientations. The results in data according to average basis (Table 1 and Fig. 1) concluded that highest plant spread E-W (49.40 cm) was recorded on planting date of 30 November (P₁) while the lowest plant spread E-W (44.43 cm) were found observed under the planting date of 14 December (P₃). Plant spread N-S was significantly affected by various planting dates. The data according to average basis are shown in Table 1 and graphically illustrated in Figure. 1. Maximal plant spread N-S (52.63 cm) was recorded under planting date of 30 November (P₁) while the lowest plant spread E-W (47.63 cm) were found under the planting date of 14 December (P₃). Typically, early planting early results in more consistent growth. This can result in more uniform east-west and north-south spread, as the plants have an extended growing period to develop and grow. These findings concur with Abed *et al.*, (2015) in cabbage, Singhal *et al.*, (2009) Thirupal *et al.*, (2014) & Kanase *et al.*, (2018) in broccoli.

Table 1. Effect of planting dates on growth parameters of cabbage (Pooled data)

Planting Date	Plant survivability (%)	Plant height (cm)			Stem diameter (cm)	No. of non-wrapper leaves	No. of wrapper leaves	Plant Spread E-W	Plant Spread N-S
		30 DAT	45 DAT	60 DAT					
P ₁	95.28	14.67	21.31	28.39	2.11	16.28	27.19	49.40	52.63
P ₂	88.39	14.01	19.45	26.72	1.95	15.17	26.34	47.54	50.20
P ₃	84.81	13.34	18.17	25.22	1.81	13.94	25.36	44.43	47.63
SEm (±)	1.81	0.18	0.23	0.24	0.02	0.13	0.19	0.32	0.44
C.D. (P=0.05)	3.66	0.37	0.47	0.49	0.04	0.27	0.38	0.65	0.89

Comment [u15]: there should be consistency of indenting all through

Comment [u16]: Is it data or results?

Comment [u17]: Apply the same for all

Comment [u18]:

Comment [u19]: ?

Comment [u20]: What was the findings of Ullah *et al.* (2013)?


Comment [u21]: Headings should be written in full beginning with "effect of....."

Comment [u22]: Research results should come as reported speech

Comment [u23]: Not listed in the reference

Comment [u24]: Include the findings of these authorities in your discussions on only their names and years

Comment [u25]: Tables should not be having more than three lines. Where necessary, use dotted lines



3.2 Effect of spacings on growth parameters of cabbage

3.2.1 Plant survivability

Growth characters of cabbage were considerably impacted by various plant densities. The data on pooled mean basis (Table 2 and Fig. 2) revealed that the significantly higher plant survivability (91.59%) was observed under plant spacings at 60 cm x 60 cm (S₄) while minimum plant survivability (86.57%) recorded found under spacing at 45 cm x 30 cm (S₁). The reason behind the plants' exceptional growth and survival may be traced back to the abundance of vital nutrients, water, light, and carbon dioxide in the 60 cm x 60 cm spacing. These findings concur with Muhammad *et al.* (2004) on onion investigation.

3.2.2 Plant height

The data on pooled mean basis (Table 2 and Fig. 2) revealed concluded that the greatest plant height at 30 days after transplanting (15.73 cm), at 45 days after transplanting (22.45 cm), and at 60 days after transplanting (31.12 cm) was recorded under plant spacing at 60 cm x 60 cm (S₄). The least height at 30 DAT (11.31 cm), at 45 DAT (16.07 cm) and at 60 DAT (21.66 cm) were reported observed under the plant spacings at 45 cm x 30 cm (S₁). This indicated it demonstrated that as plant spacing increases raises, plant height also increased. This It could be because of plants are getting enough light and nourishment. The current experiment's trend was found to be consistent with Moniruzzaman (2011), Ullah *et al.*, (2013), Singh *et al.*, (2007) and Yadav *et al.*, (2021) in cabbage.

3.2.3 Stem diameter

The result data according to average basis disclosed that stem diameter was significantly influenced by planting time (Table 1 and Fig. 1). The maximum stem diameter (2.31 cm) observed in these plants which were transplanted at 60 cm x 60 cm (S₄) spacing, while the minimum stem diameter (1.57 cm) was recorded measured in these plants which were transplanted at 45 cm x 30 cm (S₁) spacing. This could be because they received an adequate amount of light and nutrients. Similar findings are closely aligned with results of Abed *et al.*, (2015), and Yadav *et al.*, (2021) in cabbage.

3.2.4 Number of non-wrapper leaves and wrapper leaves

It is revealed from the pooled data (Table. 2 and Figure. 2) of both years that no. of non-wrapper leaf greatly impacted by various plant configurations. As plant spacing was enhanced, so did the no. of non-wrapper leaves. The maximal no. of non-wrapper leaves (17.40) were counted from the plot which

Comment [u26]: Not listed in the reference

Comment [u27]: Whose report says what?

Comment [u28]: Present their findings

Comment [u29]: ???????

Comment [u30]: Write the appropriate word

Comment [u31]: ??????

have less plant densities(60 cm x 60 cm) and the least no. of non-wrapper leaves (12.81) were obtained from plants which were transplanted at 45 cm x 30 cm (S₁) plant spacing.

Different plant spacing had a substantial impact on the number of wrapper leaves. Table 2 presents the pooled mean basis data, while Fig. 2 provides a visual representation of the same. In comparison with other spacings, the plant spacing of 60 cm x 60 cm (S₄) performed better, yielding the largest number of wrapper leaves (28.30). Conversely, the plants transplanted at 45 cm x 30 cm (S₁) plant spacing had the lowest number of wrapper leaves (23.15). More light, moisture and nutrients were available to plants planted wider apart than to those grown closer spacing together, which probably helped explain why the latter group's plants performed better during growth. The current experiment's tendency was found to be consistent with Moniruzzaman's (2011) findings in cabbage.

Comment [u32]: Recast

Comment [u33]: Recast

Comment [u34]: ????????

3.2.5 Plant spread (East-West and North-South)

The data on pooled mean basis (Table 2 and Fig. 2) revealed that highest plant spread E-W (52.79 cm) were recorded from the plant which were planted at 60 cm x 60 cm (S₄) whereas minimum plant spread E-W (40.02 cm) were found from the plots in which plants are transplanted at 45 cm x 30 cm (S₁) plant spacing.

Plant spread N-S was significantly affected by various plant densities. The data on pooled mean basis are shown in Table 2 and graphically illustrated in Fig. 2. The maximal plant spread N-S (56.08 cm) was recorded under plant spacing at 60 cm x 60 cm (S₄) whereas least plant spread E-W (42.77 cm) were found from the plants which were transplanted at 45 cm x 30 cm (S₁) plant spacing. Higher spacings typically enable cabbage plants to grow larger and spread more in both directions. With more space, plants can develop a more even and balanced spread. The decreased competition for resources fosters healthier growth and a greater overall spread. The patterns of current findings confirmed with the results of Sandhu *et al.*, (1999) in cabbage, Thirupal *et al.*, (2014) in broccoli.

Comment [u35]: ????????

Table 2. Effect of spacing on growth parameters of cabbage (Pooled data)

Spacing	Plant survivability (%)	Plant height (cm)			Stem diameter (cm)	No. of non-wrapper leaves	No. of wrapper leaves	Plant spread E-W	Plant spread N-S
		30 DAT	45 DAT	60 DAT					
S ₁	86.57	11.31	16.07	21.66	1.57	12.81	23.15	40.02	42.77
S ₂	89.23	13.71	18.53	24.80	1.84	14.01	25.97	44.03	47.23
S ₃	90.58	15.28	21.52	29.53	2.19	16.31	27.77	51.65	54.54
S ₄	91.59	15.73	22.45	31.12	2.31	17.40	28.30	52.79	56.08
S.Em(±)	1.57	0.15	0.20	0.21	0.01	0.11	0.16	0.28	0.38
C.D.	3.17	0.32	0.40	0.43	0.03	0.24	0.33	0.57	0.77

(P=0.05)




Fig. 2 Effect of spacing on growth parameters

4. CONCLUSION

The experimental findings of these experiments concluded that the planting time and spacing showed significant variation impact on among the different parameters observed. Generally, the early planting date (30 November) performed better among the different dates of planting while wider plant spacing 60 cm x 60 cm gave outstanding results on in terms of growth parameters of cabbage.

5. REFERENCES

- Abed, M., El-Said, E., &Shebl, E. Effect of planting date and spacing on yield and quality of cabbage (*Brassica oleracea* var. *capitata* L.). *Journal of Plant Production*, 2015;6(12), 2093–2102
- Bairwa, R.K., Mahawar, A.K., Singh, S.P. and Gocher, P. Influence of sulfur dose and spacing on quality attributes and economics of knolkhol (*Brassica oleracea* var. *gongylodes* L.) variety early White Vienna. *Chemical Science Review and Letters*.2017;6(22): 933-938.
- Kanase, V. J., Bhosale, A. M., Shinde, V. N. Studies on effect of planting dates on growth, yield and quality of broccoli (*Brassica oleracea* L. var. *italica*) cv. Green Magic. *International Journal of Current Microbiology and Applied Sciences*.2018;6:8-86
- Kochhar, S.L. *Economic Botany in the Tropics*, Mac Millan India Ltd., Daryaganj, New Delhi. 2010. pp. 604
- Moniruzzaman, M. Effect of plant spacings on the performance of hybrid cabbage (*Brassica oleracea* var. *capitata*) varieties. *Bangladesh Journal of Agricultural Research*.2011; 36 (3): 495-506.
- Neto, E.E., Benett, C.G.S. Santos, E.C.M.D. Reboucas, T.N.H. and Costa, E. Plant spacing and boron (B)topdressing fertilization for purple cabbage crop (*Brassica oleracea* var. *capitata*) variety purple giant. *Australian Journal of Crop Science*.2016;10(11): 1529-1533
- Panse, V.G. and Sukhatme, P.V. *Statistical Methods for Agricultural Workers*. Fourth Edition. ICAR Publication, New Delhi, 1985;187-196
- Phillips R, Rix M. *The Random House Book of Vegetables*, Random House, NY;1993.

Comment [u36]: Not cited in the text

- Saikia, D.B. Phokan and Sanchita Brahma. Effect of time of planting and planting densities on growth, yield and economic production of broccoli (*Brassica oleracea* L. var. *italica*) cv. Pusa Broccoli KTS-1. *Journal of Hill Agriculture*. 2010;1(2):135-139
- Sandhu, K. S., Chahal, B. S., Singh, D. and Sandha, M. S. Effect of plant population density and nitrogen on growth characters and yield of cabbage (*Brassica oleracea* L. var. *capitata*). *Vegetable Science*.1999; 26(2):146-148
- Singh, J.P., Munib, K., Rakshanda, M. Effect of nitrogen levels and spacing on the growth and yield of cabbage. *Environment and Ecology*. 2007; 25(1), 153-155.
- Singh, R. and Sharma, S.R. Cole crops: In-Text Book of Vegetables, Tuber crops and Spices (Eds. Thamburaj and Singh, N.) ICAR, New Delhi, 2001;76-146.
- Singh, S.K., Singh, T., Singh, B.N. and Verma, R.B. Response of fertility levels and plant density on growth, yield, and quality of hybrid cabbage. *Vegetable Science*.2004; 31(1): 69-72.
- Singhal, B. K., Preeti, B. K., Srivastava, Singh, M.P., Singh, P. K. Effect of date of planting and spacing on the performance of broccoli. *Indian Journal of Horticulture*. 2009;66(1):137-140
- Splitstoeser W.E. (1979). *Vegetable growing handbook*. The AVI Publishing Company, Inc., USA.,
- Ullah, A., Islam, M. N., Hossain, M. I., Sarkar, M. D., & Moniruzzaman, M. Effect of planting time and spacing on growth and yield of cabbage. *International Journal of Bio-resource and Stress Management*. 2013;4(2), 182–186.
- Yasmin, M. F. Effect of planting time and plant density of top shoot cuttings in potato production (Doctoral dissertation, MS thesis, Salna, Gazipur: Department of Horticulture, Bangabandhu Sheikh Mujibur Rahman Agricultural University);2008
- Yadav, R., Ahamad, S., Sharma, N., & Raj, R. Effect of different planting time and spacing on growth, yield and quality of cabbage (*Brassica oleracea* var. *capitata*) cv. Golden Acre under Bundelkhand region in U.P. *International Journal of Chemical Studies*, 2021;9(2), 358–363.

Comment [u37]: Not cited in the text