

THE USE OF ANTI-INFLAMMATORY, ANALGESICS, AND ANTIPYRETICS DURING PREGNANCY: A POPULATION COHORT STUDY IN A CAPITAL CITY OF THE NORTH REGION OF BRAZIL.

Abstract

Objective: To analyze the utilization of anti-inflammatory drugs and associated factors during pregnancy. **Materials and Methods:** This is a population-based cross-sectional study conducted in Rio Branco, AC. The study included all women who gave birth between April 6 and July 10, 2015. Demographic, socioeconomic, reproductive, maternal habits, prenatal care, and newborn status were investigated. Multivariate analysis with logistic regression was performed, considering $p < 0.05$ for association. **Results:** 1190 postpartum women were interviewed. Anti-inflammatories were not used by only 13.2% of pregnant women. The prevalence of dipyron use was 72.7%, paracetamol 50.3%, nimesulide 16.1%, and diclofenac 5.2%. Women with lower educational levels (up to high school $OR=1.55$, 95% CI 1.07-2.25), those who consumed alcohol ($OR=1.96$, 95% CI 1.03-3.73), and those with a higher number of living children (more than 4 children $OR=1.6$, 95% CI 1.06-2.24, 2 to 3 children $OR=1.52$, 95% CI 1.10-2.27) had a higher chance of using anti-inflammatories during pregnancy. Primiparous women ($OR=0.66$, 95% CI 0.47-0.93), those with more than eight prenatal visits ($OR=0.55$, 95% CI 0.35-0.85), and those experiencing moderate to severe food insecurity ($OR=0.42$, 95% CI 0.25-0.69) had a lower chance of using anti-inflammatories during pregnancy. After adjusted analysis, alcohol consumption and the number of living children lost statistical significance and were not included in the final model as independent variables. **Conclusion:** The prevalence of anti-inflammatory use was high. Alternative therapies and restricting prescriptions to necessary cases could be proposed to reduce the use of non-steroidal anti-inflammatory drugs (NSAIDs) during pregnancy.

Key-words: Anti-inflamatórios, gestantes, pré-natal.

Introduction

Medications with anti-inflammatory, analgesic, and antipyretic properties serve multiple physiological functions, often overlapping in their efficacy. While certain drugs may excel in one aspect over another, their shared mechanism of action—primarily the inhibition of cyclooxygenases (COX-1 and COX-2) and subsequent suppression of prostaglandin synthesis allow them to effectively address pain, inflammation, and fever across the spectrum (1).

In the context of pregnancy, where ensuring the safety of both the mother and the developing fetus is paramount, the decision to utilize such medications demands a careful, personalized approach. Factors such as the specific risks associated with the medication, the gestational period, optimal dosage, and duration of therapeutic intervention must be meticulously considered. Recognizing the inherent risks accompanying the use of any pharmacological agent, a thorough evaluation of the potential benefits against these risks becomes imperative (2).

Conditions during pregnancy that may necessitate the use of these therapeutic agents are commonplace, including complaints such as lower back pain, headaches, leg pain, and fever resulting from infections (3). Importantly, it should be noted that alternative therapies exist for some of these conditions, offering a medication-free approach that also prepares the body for the anticipated changes associated with pregnancy. This approach becomes crucial in minimizing exposure to medications, as their potential adverse effects on the reproductive cycle warrant further in-depth exploration (4).

Non-steroidal anti-inflammatory drugs (NSAIDs), including ibuprofen and aspirin, are commonly used analgesics and antipyretics. Their mechanism involves the inhibition of prostaglandin synthesis, which, while effective in managing symptoms, can pose significant risks during pregnancy. For instance, NSAIDs are known to affect the fetal cardiovascular system, particularly causing premature closure of the ductus arteriosus, a vital part of fetal circulation (5). Additionally, the use of these medications during the third trimester is associated with an increased risk of prolonged labor and postpartum hemorrhage due to their effect on platelet function and uterine contractility (6).

Paracetamol, also known as acetaminophen, is another widely used analgesic and antipyretic considered safer for use during pregnancy. However, recent studies have raised concerns about its potential association with neurodevelopmental disorders in children when used extensively during pregnancy. Despite being a first-line treatment for fever and mild to moderate pain, its use should be carefully monitored and limited to the lowest effective dose (7).

The prevalence of self-medication during pregnancy is a significant public health concern. Many pregnant women opt to self-medicate due to the ease of access to over-the-counter medications and the commonality of ailments such as headaches and back pain. This behavior underscores the need for better education and counseling regarding the risks and safe practices associated with medication use during pregnancy. Healthcare providers play a crucial role in guiding pregnant women to make informed decisions about their health and the health of their unborn children (8-10).

In Brazil, the use of medications during pregnancy is influenced by various factors, including socioeconomic status, access to healthcare, and cultural practices. Studies have shown that women with lower educational levels and those with limited access to healthcare services are more likely to self-medicate. This highlights the disparities in healthcare access and the need for targeted interventions to educate and support these vulnerable populations (9).

This article seeks to delve into the prevalence of anti-inflammatory, analgesic, and antipyretic use during pregnancy, along with the associated factors, focusing on data from the municipality of Rio Branco/AC in the year 2015. Understanding the patterns of medication use and the underlying factors can inform public health strategies to improve maternal and fetal health outcomes.

The decision-making process regarding medication use during pregnancy is complex and multifaceted. It involves not only the clinical assessment of the benefits and risks but also the social, cultural, and economic contexts in which pregnant women live (9). Therefore, a comprehensive approach that includes medical guidance, patient education, and supportive policies is essential to ensure the safe use of medications during pregnancy (10,11).

The primary aim of this study is to analyze the utilization of anti-inflammatory drugs and associated factors among pregnant women in Rio Branco, AC. By examining demographic, socioeconomic, reproductive, and maternal health variables, we aim to identify the determinants of medication use and the potential risks involved. This knowledge is crucial for developing targeted interventions to promote safe medication practices and ultimately improve the health of mothers and their babies.

Methods

Study Design

This study employs a cross-sectional, population-based design.

Population and Sample

Rio Branco, the capital of the state of Acre, had an estimated population of 377,057 inhabitants in 2015, with the majority residing in the urban area (89.4%). The city hosts two maternity

hospitals: the Santa Juliana Hospital and Maternity (HSJ) and the Bárbara Heliodora Maternity, the latter serving only the public system. The sample was based on 6,943 deliveries in these hospitals in 2014. With a 3% sampling error, 80% test power, and an estimated odds ratio of 2.0, 926 pregnant women were initially required. The final sample included 1,190 postpartum women.

Instruments

Data were collected through interviews with postpartum women and medical record reviews. A standardized and semi-structured research questionnaire was used. Interviews were conducted by trained interviewers, including undergraduate students and/or professionals in health-related fields, who underwent prior training and were financially supported during the data collection period.

Data Collection Procedures

Data collection occurred between April 6 and July 10, 2015, at the two maternity hospitals in the municipality, specifically among parturients residing in the urban area. Pregnant women were identified through the hospital admission record, invited to participate in the research, and asked to sign an informed consent form. Interviews took place within 24 hours after delivery. The collected information was reviewed, coded, and entered into the database. Inconsistencies were addressed through questionnaire review and/or telephone contact with the parturients.

Statistical Procedures

Data analysis was performed using IBM SPSS 22.0 for Windows. The study began with a description of the study population using prevalence rates for each independent variable. The outcome's prevalence was measured according to the studied variables. Both crude and adjusted analyses were conducted using logistic regression to calculate odds ratios and 95% confidence intervals (95% CI). Variables with a p-value < 0.05 in Wald tests were considered associated with the outcome.

Ethical Considerations

This study adhered to ethical principles in line with Resolution No. 466/12 of the National Health Council. Authorization was obtained from the institutions where data collection took place. The main project was approved by the Research Ethics Committee of the Federal University of Acre, and this subproject was approved by the Research Ethics Committee of the National School of Public Health.

Results

Of the 1,190 pregnant women in the sample, the majority were aged between 25 and 34 years (38.7%), had completed high school (51.3%), had an income above 1.5 minimum wages (56.4%), belonged to economic classes C, D, or E (79.4%), and reported having a partner (84.0%) (Table 1). Regarding prenatal care, most pregnant women received it in the public health system (85.1%), with six or more consultations (60.7%), and 63.4% reported unplanned pregnancies. The majority of pregnant women were multiparous (60.7%) and reported having 2 to 3 living children (31.1%). Smoking was reported by 9.6% of these women, 12.3% reported alcohol consumption, and only 10.7% engaged in physical activity during pregnancy. The prevalence of cesarean section was 48.0% (Table 1).

*Minimum wage in force at the time (R\$ 788,00)

**ABEP – Brazilian Association of Research Companies

Table 1: Socioeconomic, demographic, maternal and prenatal care habits characteristics of the study population in the municipality of Rio Branco - AC, 2015. (N=1190)

Variable	N	%
Age (years)		
13 -18	221	18,6
19 – 24	386	32,4
25 – 34	460	38,7
≥ 35 anos	123	10,3
Education		
Elementary school I	77	6,5
Elementary school II	232	19,5
High school	611	51,3
Higher education	270	22,7
Family income (minimum wages) *		
Up to 1.5 MW	444	43,6
≥ 1.5 MW	574	56,4
ABEP Class**		
A and B	242	20,6
Receipt of Bolsa Família	935	79,4
No	914	80,9
Yes	216	19,1
Marital status		
No partner	190	16,0
With partner	999	84,0
Self-declared skin color		
White	125	10,5
Non-white	1064	89,5
Food and Nutritional Security		
Mild Security and Insecurity	777	65,3
Moderate and severe insecurity	413	34,7
Prenatal Care		
Public	982	85,1
Private	172	14,9
Number of Prenatal Consultations		
None	17	1,4
1 to 5	450	37,8
6 to 8	494	41,5
> 8	229	19,2
Planned pregnancy		
No	751	63,4
Yes	434	36,6
Primigravida		
No	718	60,7
Yes	464	39,3

Number of living children (including newborns)		
One	466	39,3
2 or 3	369	31,1
4 or more	352	29,7
Smoked during pregnancy		
No	1076	90,4
Yes	114	9,6
Drank alcoholic beverages during pregnancy		
No	1035	87,7
Yes	145	12,3
Practicipated physical activities		
No	1046	89,3
Yes	125	10,7
Type of delivery		
Normal	618	52,0
Cesarean section	570	48,0
Use of anti-inflammatory medication by pregnant woman		
Did not take	157	13,2
Took one type	485	40,8
Took two types	427	35,9
Took three types	105	8,8
Took four types	16	1,3
Dipyron		
No	325	27,3
Yes	865	72,7
Paracetamol		
No	591	49,7
Yes	599	50,3
Nimesulide		
No	998	83,9
Yes	192	16,1
Diclofenac		
No	1128	94,8
Yes	62	5,2

Only 13.2% of the respondents did not use anti-inflammatory drugs during pregnancy. The prevalence of dipyron use was 72.7%, paracetamol 50.3%, nimesulide 16.1%, and diclofenac 5.2%. It is noteworthy that 46.0% of the women reported using two or more of these drugs during pregnancy (Table 1).

The frequency of anti-inflammatory drug use varied according to the type of medication. For dipyron, the proportion was higher when the pregnant woman had completed lower secondary and upper secondary education, belonged to economic classes C, D, or E, had fewer prenatal consultations, consumed alcoholic beverages during pregnancy, did not engage in physical activity, was not a primipara, and had a higher number of living children. The use was higher among those who had prenatal care in the public health system (Table 2).

The proportion of paracetamol consumption was higher when pregnancy was unplanned and lower among pregnant women with one living child (Table 2).

UNDER PEER REVIEW

For nimesulide, the highest proportion of use occurred in women aged 25 or older, with lower educational attainment, experiencing food insecurity, having fewer prenatal consultations, reporting smoking during pregnancy, and receiving prenatal care in the public service (Table 2).

Diclofenac showed a higher proportion of use by pregnant women with lower educational attainment, experiencing food insecurity, having fewer prenatal consultations, and reporting smoking during pregnancy (Table 2)

Table 2 :Prenatal consultations, and reporting smoking during pregnancy

Age (years)	Dipyron			Paracetamol			Nimesulide			Diclofenac		
	No (%)	Yes (%)	p-value	No (%)	Yes (%)	p-value	No (%)	Yes (%)	p-value	No (%)	Yes (%)	p-value
13 - 18	53 (24.0%)	168 (76.0%)	0.387	110 (49.8%)	111 (50.2%)	0.275	181 (81.9%)	40 (18.1%)	0.016	210 (95.0%)	11 (5.0%)	0.064
19 – 24	108 (28.0%)	278 (72.0%)		178 (46.1%)	208 (53.9%)		311 (80.6%)	75 (19.4%)		358 (92.7%)	28 (7.3%)	
25 – 34	124 (27.0%)	336 (73.0%)		235 (51.1%)	225 (48.9%)		393 (85.4%)	67 (14.6%)		445 (96.7%)	15 (3.3%)	
≥ 35 years	40 (32.5%)	83 (67.5%)		68 (55.3%)	55 (44.7%)		113 (91.9%)	10 (8.1%)		115 (93.5%)	8 (6.5%)	
Education												
Fundamental I	24 (31.2%)	53 (68.8%)	0.014	38 (49.4%)	39 (50.6%)	0.385	62 (80.5%)	15 (19.5%)	0.002	70 (90.9%)	7 (9.1%)	0.044
Fundamental II	60 (25.9%)	172 (74.1%)		127 (54.7%)	105 (45.3%)		197 (84.9%)	35 (15.1%)		217 (93.5%)	15 (6.5%)	
High School	148 (24.2%)	463 (75.8%)		294 (48.1%)	317 (51.9%)		494 (80.9%)	117 (19.1%)		577 (94.4%)	34 (5.6%)	
College	93 (34.4%)	177 (65.6%)		132 (48.9%)	138 (51.1%)		245 (90.7%)	25 (9.3%)		264 (97.8%)	6 (2.2%)	
Family Income												

Up to 1.5 SM	108 (24.3%)	336 (75.7%)	0.181	215 (48.4%)	229 (51.6%)	0.739	365 (82.2%)	79 (17.8%)	0.259	414 (93.2%)	30 (6.8%)	0.248
≥ 1.5 SM	161 (28.0%)	413 (72.0%)		284 (49.5%)	290 (50.5%)		487 (84.8%)	87 (15.2%)		545 (94.9%)	29 (5.1%)	
ABEP Class												
A and B	90 (37.2%)	152 (62.8%)	0.001	120 (49.6%)	122 (50.4%)	0.961	209 (86.4%)	33 (13.6%)	0.206	229 (94.6%)	13 (5.4%)	0.935
C, D, and E	230 (24.6%)	705 (75.4%)		462 (49.4%)	473 (50.6%)		776 (83.0%)	159 (17.0%)		886 (94.8%)	49 (5.2%)	
Receives Bolsa Família												
No	258 (28.2%)	656 (71.8%)	0.074	456 (49.9%)	458 (50.1%)	0.735	774 (84.7%)	140 (15.3%)	0.070	863 (94.4%)	51 (5.6%)	0.578
Yes	48 (22.2%)	168 (77.8)										

The prevalence of any anti-inflammatory drug use during pregnancy was 86.8%. In bivariate analysis, women with lower educational attainment, up to complete high school (Crude OR = 1.55, 95% CI 1.07-2.25), with reference to those with a higher education level, those who consumed alcoholic beverages (Crude OR = 1.96, 95% CI 1.03-3.73), and those with a higher number of living children (more than 4 children, Crude OR = 1.6, 95% CI 1.06-2.24, and 2 to 3 children, Crude OR = 1.52, 95% CI 1.10-2.27), had a higher chance of using anti-inflammatory drugs. Primiparous women (Crude OR = 0.66, 95% CI 0.47-0.93), those who had more than eight prenatal consultations (Crude OR = 0.55, 95% CI 0.35-0.85), and those in a situation of moderate to severe food insecurity (Crude OR = 0.42, 95% CI 0.25-0.69) had a lower chance of using anti-inflammatory drugs during pregnancy. After adjusted analysis, alcohol consumption and the number of living children lost statistical significance and did not remain in the final model as independent variables associated with the use of anti-inflammatory drugs (Table 3).

Table 3 : Independent variables associated with the use of anti-inflammatory drugs

Variable	Yes (n/%)	OR (Crude)	95% CI	p-value	OR (Adjusted)	95% CI
Age (years)						
Up to 35 years		931 (87.3%)	1	0.179		
≥ 35 years		102 (82.9%)	0.71	(0.42-1.17)		

Education					
College	223 (82.6%)	1	0.021	1	
Up to complete high school	810 (88.0%)	1.55	(1.07-2.25)	1.46	(0.99-2.17)
Family Income (minimum wages)					
Up to 1.5 SM	392 (88.3%)	1	0.374		
≥ 1.5 SM	496 (86.4%)	0.84	(0.58-1.22)		
ABEP Class					
A and B	203 (83.9%)	1	0.107		
C, D, and E	821 (87.8%)	1.38	(0.93-1.05)		
Receives Bolsa Família					
No	792 (86.7%)	1	0.608		
Yes	190 (88.0%)	1.12	(0.71-1.76)		
Marital Status					
No partner	163 (85.8%)	1	0.655		
With partner	869 (87.0%)	1.10	(0.70-1.73)		
Skin Color					
White	114 (91.2%)	1	0.128		
Non-White	918 (86.3%)	0.60	(0.31-1.15)		
Food and Nutrition Security					
Mild insecurity	937 (87.7%)	1	0.011	1	
Moderate and severe insecurity	96 (79.3%)	0.54	(0.33-0.87)	0.421	(0.25-0.69)
Number of Prenatal Consultations					
1 to 5	398 (88.4%)	1	0.014	1	
6 to 8	434 (87.9%)	0.95	(0.63-1.40)	0.95	(0.63-1.42)
> 8	185 (80.8%)	0.55	(0.35-0.85)	0.57	(0.36-0.90)
Smoked during pregnancy					
No	932 (86.6%)	1	0.553		
Yes	101 (88.6%)	1.2	(0.65-2.19)		
Consumed alcohol during pregnancy					
No	891 (

UNDER PEER REVIEW

Discussion

A prevalent overall use of these medications during pregnancy was observed at 86.8%. Dipyron was the most commonly used medication (72.7%), followed by paracetamol (50.3%) and nimesulide (16.1%), while 5.2% of pregnant women reported using diclofenac. Comparing these results with the scientific literature proved challenging, as studies classifying analgesics, anti-inflammatories, and antipyretics in the same group were scarce. However, prevalence results from publications analyzing medication use during pregnancy were obtainable.

The prevalence of using anti-inflammatories, analgesics, and antipyretics during pregnancy (86.8%) can be considered high compared to various studies analyzing the use of this therapeutic class. A population-based study analyzing medication consumption among 5,564 pregnant women in six Brazilian cities reported a prevalence of analgesic/anti-inflammatory use at 22.2% (10,11). In Santa Rosa/RS, a population-based study with 470 pregnant women found a prevalence of 17.6% for non-opioid analgesics (12). Another population-based study with 1,091 pregnant women in Santo Antônio de Jesus/BA reported a prevalence of analgesic use at 21.9% (13).

In Maceió/AL, a study with 130 pregnant women treated at the university hospital reported that 19.0% of pregnant women used analgesics, anti-inflammatories, and antipyretics (14). Another study with a convenience sample of military pregnant women (n=100) in Belo Horizonte/MG reported that 4.6% reported using analgesics and anti-inflammatories (15).

Observing studies from other countries also indicates that the percentage of usage is lower than found in this research. It is essential to consider the diverse study designs, but prevalence rates for the use of these medications in each respective group could be extracted. In a historical cohort study involving 65,547 women in Israel, exposure to anti-inflammatories was 6.9%, with the analysis including the use of ibuprofen, diclofenac, naproxen, etodolac, indomethacin, lornoxicam, or nabumetone (16). In Canada, a case-control study with 4,705 women diagnosed with spontaneous abortion reported a prevalence of anti-inflammatory use at 7.5% among cases and 2.6% among controls (17). In Norway, a cohort study with 69,929 women found that 4.32% of women reported using anti-inflammatories early in pregnancy (4). Another cohort study, also in Norway, with 90,417 pregnant women, reported that 7.2% of them used at least one of the four analyzed anti-inflammatories (18). In Quebec, Canada, in a case-control study with 36,387 pregnant women, the prevalence of anti-inflammatory use was 2.9% in the first trimester of gestation (19), and in San Francisco, USA, a cohort with 1,055 pregnant women, 5% reported using anti-inflammatories during pregnancy and around conception (3).

Dipyron was the most used analgesic in this study, reported by 72.7% of pregnant women. Dipyron has been cited as the drug of choice in some other studies in Brazil (10,20,21). There is controversy regarding dipyron's general use, as the Food and Drug Administration (FDA) prohibited it in 1977 in the United States, citing the potential risk of irreversible agranulocytosis (22). This American initiative has been widely debated, but given the global influence of the FDA, other countries (England, Canada, Denmark, Norway, Sweden, Japan, and Australia) follow this guidance (23,24). In Brazil, in 2001, the National Health Surveillance Agency (ANVISA) conducted an "International Panel for the Assessment of Dipyron Safety," recognizing its undeniable effectiveness as an analgesic and antipyretic. The reported risks were deemed low and insufficient to remove the drug from the market. The recommendation was not to withdraw it (23,25).

Caution is required in the use of this medication among pregnant groups, as the risk of developing agranulocytosis cannot be the sole attribution. Literature reports suggest an association of dipyron with cases of childhood acute leukemia (20,26,27), orofacial clefts in newborns (28), Wilms tumor (29), spontaneous abortion, and congenital malformation (30).

Paracetamol, the second most commonly reported medication among pregnant women in Rio Branco/AC, was mentioned by 50.3%. The literature designates this drug as the preferred choice for prescription during pregnancy, considering the strong controversies surrounding dipyrene. The FDA classifies paracetamol as low risk for pregnant women based on studies in animals and humans. In Brazil, it is widely used (31,32). Even though it presents a lower prevalence than observed for dipyrene, it is still considerably high when compared to other research in the country. A study in Rio de Janeiro/RJ found a prevalence of 35.5% (20), and in a study analyzing medication use in six Brazilian capitals, the prevalence was 18.3% (10). In a population-based study in Santa Rosa/RS, paracetamol had the highest prevalence at 17.3% (12), and in another study in Santo Antônio de Jesus/BA, it appeared as the most chosen analgesic at 21.3% (13). Convenience sample studies report lower prevalence rates, such as those observed in João Pessoa/PB at 4.93% (33), Natal/RN at 13.1% (34), Piracicaba/SP at 6.3% (35), and Campinas/SP in 5.5% of pregnant women (21).

International studies indicate higher prevalence rates than in Brazil, likely due to the prohibition and/or contraindication of dipyrene in their countries. In three studies in Denmark, with cohort and cross-sectional designs, the following prevalence rates were reported: in a cohort study with 63,652 pregnant women, paracetamol use during pregnancy was 56.3% (36); another cohort with 64,322 participants where 56% reported paracetamol use during pregnancy (38); in a cross-sectional study with 10,209 pregnant women, prevalence rates of 35.2% and 6.5% before and at the beginning of pregnancy, respectively (37). In New Zealand, a cohort study with 871 pregnant women reported a prevalence of paracetamol exposure during pregnancy at 49.8% (39), and in Chicago, USA, a cohort study with 345 women observed that 70% declared paracetamol use at least once during pregnancy (40).

Risks associated with the use of paracetamol, as mentioned in the literature and warranting further exploration, include an association with respiratory problems such as asthma, wheezing, and chest tightness in children under one year. These effects are believed to be a result of paracetamol use after the twentieth week of gestation. Additionally, some studies suggest an increased risk of childhood autism, attention deficit disorders, and the occurrence of the congenital malformation gastroschisis (36,38,40–42).

Nimesulide, reported by 16.1% of pregnant women, is not a preferred anti-inflammatory for use during pregnancy, as the risks to the fetus due to its use are not well elucidated in humans and have been deemed inappropriate in animal trials (31,43). Some studies describe the risk of this medication being associated with fetal cardiac abnormalities, but these studies have methodological limitations, and their conclusions should be approached with due care and caution (44,45).

The use of diclofenac was reported by 5.2%. This medication is not among the first-choice anti-inflammatories, although it is classified as low risk for the gestational period (43). This finding aligns with low prevalences reported in studies conducted in other locations (17,18,46). Risks associated with its use include low birth weight when used in the second trimester and a significant association with maternal vaginal bleeding when used in the third trimester (OR 1.8; 95% CI 1.1-3.0) (18). In other studies, diclofenac has been strongly associated with episodes of spontaneous abortion (17,46).

When analyzing the use of anti-inflammatories collectively, this study indicated a higher odds ratio among pregnant women who consumed alcohol. On the other hand, the likelihood of exposure to anti-inflammatories was lower in pregnant women with a college education compared to those with only a basic education, as well as among primiparous women, those with six to eight prenatal consultations, and pregnant women experiencing moderate to severe food insecurity. Higher educational attainment appears to lead to a better understanding of the inherent risks of medication use, while being a first-time mother reflects self-care and maternal

inexperience. Adequate prenatal care is consistently identified in various studies as a protective factor against the use of risky medications. Alcohol consumption, besides indicating a lack of care during pregnancy, also necessitates addressing the adverse effects resulting from alcohol use (11,47–49).

Given the cross-sectional design and the use of interviews as a data collection technique, albeit with the utilization of institutional records from the maternity card and medical records, it is essential to describe limitations. There is a possibility of memory bias, confusion with medication names, and the reluctance of pregnant women to report use, knowing the potential implications for the neonate. These limitations were duly considered during the research implementation, with all measures taken to overcome them.

Despite knowing the gestational complications that require the use of medications in this therapeutic class and recognizing the influence of the service with excellent prenatal coverage, it is concluded that the prevalence of using these medications was high. In this regard, measures can and should be taken to restrict this practice. Despite studies emphasizing the benefits that outweigh the risks, these risks should not be disregarded. It is possible to introduce alternative therapies such as physiotherapy, acupuncture, and physical activities, among others, capable of alleviating pain and inflammation, thus limiting drug therapy only to cases where it is essential.

What does this article add to the field?

This article contributes to the existing literature by providing insights into the prevalence and patterns of using anti-inflammatory, analgesic, and antipyretic medications during pregnancy in Rio Branco/AC, Brazil. The study not only presents the prevalence rates of medication use but also delves into specific drugs such as dipyron, paracetamol, nimesulide, and diclofenac.

The findings shed light on the high prevalence of medication use during pregnancy in the studied population, with a particular emphasis on the widespread use of dipyron and paracetamol. Moreover, the study identifies associated factors, including demographic, socioeconomic, reproductive, and lifestyle factors, providing a comprehensive understanding of the context in which these medications are utilized.

Comparisons with existing literature highlight the unique aspects of medication use in this specific region, as prevalence rates may differ across populations and geographies. The article also discusses the controversies surrounding certain medications, such as dipyron, providing a nuanced perspective on the risks and benefits associated with their use during pregnancy.

Furthermore, the study emphasizes the need for alternative therapeutic approaches, suggesting that interventions like physiotherapy, acupuncture, and physical activities could be considered to reduce reliance on medication, particularly in cases where the risks of drug use need careful evaluation.

In summary, this article adds valuable information to the literature by offering a detailed analysis of medication use during pregnancy in a specific Brazilian region, contributing to a broader understanding of prescription patterns, associated factors, and potential avenues for promoting safer therapeutic practices during gestation.

Although our manuscript has a robust conjunct of data we need to consider limitations. Firstly, its cross-sectional design restricts establishing causal relationships. Secondly, reliance on self-reported data introduces potential recall and social desirability biases. Thirdly, the sample's representativeness might be skewed, as it focuses on specific maternity hospitals. Fourthly, the findings' generalizability is constrained to the Rio Branco/AC population. Lastly, incomplete medication assessment and the absence of longitudinal data limit the study's depth and understanding of medication use during pregnancy.

Conclusions

In conclusion, this study sheds light on the prevalent use of anti-inflammatory, analgesic, and antipyretic medications during pregnancy in Rio Branco/AC. The high prevalence, especially considering potential associated risks, underscores the need for cautious evaluation and alternative approaches. The findings emphasize the importance of informed decision-making, personalized care, and promoting non-pharmacological interventions. Despite the study's contributions, addressing this complex issue requires a comprehensive, multidisciplinary approach, integrating healthcare, education, and public awareness initiatives to optimize maternal and fetal well-being during pregnancy.

References

1. Jasięcka A, Maślanka T, Jaroszewski JJ. Pharmacological characteristics of metamizole. *Pol J Vet Sci* [Internet]. 2014 Jan 1 [cited 2016 Feb 24];17(1). Available from: <http://www.degruyter.com/view/j/pjvs.2014.17.issue-1/pjvs-2014-0030/pjvs-2014-0030.xml>
2. Silva P. *Farmacologia*. Rio de Janeiro: Guanabara koogan; 2010.
3. Li D-K, Liu L, Odouli R. Exposure to non-steroidal anti-inflammatory drugs during pregnancy and risk of miscarriage: population based cohort study. *Bmj*. 2003;327(7411):368.
4. van Gelder MMHJ, Roeleveld N, Nordeng H. Exposure to Non-Steroidal Anti-Inflammatory Drugs during Pregnancy and the Risk of Selected Birth Defects: A Prospective Cohort Study. *Morty RE*, editor. *PLoS ONE*. 2011 Jul 18;6(7):e22174.
5. Brito JLOP de, Torquato IMB, Trigueiro JVS, Medeiros HA de, Souza Neto VL de, Albuquerque AM de. Lombalgia: prevalência e repercussões na qualidade de vida de gestantes. *Rev Enferm UFSM* [Internet]. 2014 Aug 27 [cited 2017 May 16];4(2). Available from: <http://cascavel.ufsm.br/revistas/ojs-2.2.2/index.php/reufsm/article/view/12231>
6. Cleves MA, Savell VH, Raj S, Zhao W, Correa A, Werler MM, et al. Maternal use of acetaminophen and nonsteroidal anti-inflammatory drugs (NSAIDs), and muscular ventricular septal defects. *Birt Defects Res A Clin Mol Teratol*. 2004 Mar;70(3):107–13.
7. Feldkamp ML, Meyer RE, Krikov S, Botto LD. Acetaminophen use in pregnancy and risk of birth defects: findings from the National Birth Defects Prevention Study. *Obstet Gynecol*. 2010;115(1):109–115.
8. MHB ZM, Melo Jr EF, Simões R. Pregnancy and Analgesia [Internet]. Brazilian Federation of Gynecology and Obstetrics Associations; 2011 [cited 2016 Feb 24]. 18 p. Available from: http://www.projetodiretrizes.org.br/ans/diretrizes/gestacao_e_analgesia.pdf
9. Carvalho MECC, Lima LC, de Lira Terceiro CA, Pinto DRL, Silva MN, Cozer GA, et al. Low back pain during pregnancy. *Braz J Anesthesiol*. 2017 May;67(3):266–70.
10. Mengue SS, Schenkel EP, Duncan BB, Schmidt MI. Medication use by pregnant women in six Brazilian cities. *Rev Saúde Pública*. 2001;35(5):415–420.
11. Mengue SS, Schenkel EP, Schmidt MI, Duncan BB. Factors associated with medication use during pregnancy in six Brazilian cities. *Cad Saúde Pública*. 2004;20(6):1602–1608.
12. Silva Brum LF da, Pereira P, Felicetti LL, Silveira RD da. Medication use by pregnant women using the Unified Health System in the city of Santa Rosa (RS, Brazil). *Ciênc Saúde Coletiva* [Internet]. 2011 [cited 2017 Jul 22];16(5). Available from: <http://www.redalyc.org/html/630/63018749012/>
13. Costa DB, Coelho HLL, Santos DB dos. Medication use before and during pregnancy: prevalence and associated factors. *Cad Saúde Pública* [Internet]. 2017 [cited 2017 Jul 22];33(2). Available from: http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0102-311X2017000205005&lng=pt&tlng=pt
14. Filho AD de O, Gama DP da, Dias JMG, Leopardi M a ria das G, Júnior DP de L, Neves SJF. Self-reported adherence to prescribed medications during pregnancy. *Rev Bras Gynecol Obstet*. 2012;34(4):147–52.

15. Freitas TCSB, Lamounier JA, Chaves RG, Silva SC da. Use of medications during pregnancy and lactation in military women in the metropolitan region of Belo Horizonte and its association with breastfeeding duration. *Rev MÉDICA MINAS GERAIS-RMMG* [Internet]. 2012 [cited 2014 Sep 1];22(2). Available from: <http://www.medicina.ufmg.br/rmmg/index.php/rmmg/article/viewArticle/516>
16. Daniel S, Koren G, Lunenfeld E, Bilenko N, Ratzon R, Levy A. Fetal exposure to nonsteroidal anti-inflammatory drugs and spontaneous abortions. *Can Med Assoc J*. 2014;186(5):E177–E182.
17. Nakhai-Pour HR, Broy P, Sheehy O, Berard A. Use of nonaspirin nonsteroidal anti-inflammatory drugs during pregnancy and the risk of spontaneous abortion. *Can Med Assoc J*. 2011 Oct 18;183(15):1713–20.
18. Nezvalová-Henriksen K, Spigset O, Nordeng H. Effects of ibuprofen, diclofenac, naproxen, and piroxicam on the course of pregnancy and pregnancy outcome: a prospective cohort study. *BJOG Int J Obstet Gynaecol*. 2013 Jul;120(8):948–59.
19. Ofori B, Oraichi D, Blais L, Rey E, Bérard A. Risk of congenital anomalies in pregnant users of non-steroidal anti-inflammatory drugs: a nested case-control study. *Birth Defects Res B Dev Reprod Toxicol*. 2006 Aug;77(4):268–79.
20. Couto AC, Ferreira JD, Pombo-de-Oliveira MS, Koifman S. Pregnancy, maternal exposure to analgesic medications, and leukemia in Brazilian children below 2 years of age: *Eur J Cancer Prev*. 2015 May;24(3):245–52.
21. Fonseca MRCC da, Fonseca E da, Bergsten-Mendes G. Prevalence of medication use during pregnancy: a pharmacoepidemiological approach. *Rev Saúde Pública*. 2002;36(2):205–12.
22. US, Food and Drug Administration. Rx Drug Study Bulletin #231: Drug Products Containing Dipyrone [Internet]. USA: Food and Drug Administration; 1977. Available from: http://www.fda.gov/ohrms/dockets/ac/00/backgrd/3634b1a_tab2b.pdf
23. Danieli P, Leal MB. Safety assessment of dipyrone: a review. *Rev Bras Farm*. 2003;84(1):17–20.
24. Hedenmalm K, Spigset O. Agranulocytosis and other blood dyscrasias associated with dipyrone (metamizole). *Eur J Clin Pharmacol*. 2002 Jul;58(4):265–74.
25. Brazil, ANVISA. Dipyrone International Safety Assessment Panel. Brasília: Ministry of Health; 2001. 11 p.
26. Emerenciano M, Koifman S, Pombo-de-Oliveira MS. Acute leukemia in early childhood. *Braz J Med Biol Res* 2007;40(6):749–760.
27. Pombo-de-Oliveira MS, Koifman S, Brazilian Collaborative Study Group of Infant Acute Leukemia. Infant Acute Leukemia and Maternal Exposures during Pregnancy. *Cancer Epidemiol Biomarkers Prev*. 2006 Nov 28;15(12):2336–41.
28. Leite ICG, Paumgarten FJR, Koifman S. Orofacial clefts in newborns and the use of medications and maternal health conditions: a case-control study in the city of Rio de Janeiro, Brazil. *Rev Bras Saúde Materno Infant*. 2005;5(1):35–43.
29. Sharpe C, Franco E. Use of dipyrone during pregnancy and risk of Wilm's tumor. *Epidemiology*. 1996;7:533–535.
30. Bar-Oz B, Clementi M, Giantonio ED, Greenberg R, Beer M, Merlob P, et al. Metamizole (dipyrone, optalgin) in pregnancy, is it safe? A prospective comparative study. *Eur J Obstet Gynecol Reprod Biol*. 2005 Apr;119(2):176–9.
31. Brasil. National therapeutic formulary 2010: RENAME 2010. Secretariat of Science, Technology and Strategic Inputs. Brasília: Ministry of Health; 2011.
32. Brazil. National List of Essential Medicines, RENAME, 2014. Department of Pharmaceutical Assistance. 9th ed. Brasília: Ministry of Health; 2015. 230 p.
33. Ribeiro NKR, Leite LLB, da Silva Pontes ZBV. Pharmacoepidemiological study: the use of medicines by pregnant women. *Rev Eletrônica Farmácia*. 2013;10(1):11.

34. Guerra GCB, Silva AQB da, França LB, Assunção PMC, Cabral RX, Ferreira AA de A. Use of medicines during pregnancy in the city of Natal, Rio Grande do Norte, Brazil. *Rev Bras Ginecol Obstet.* 2008;30(1):12–8.
35. Carmo TA do, Nitrini SMO. Drug prescriptions for pregnant women: a pharmacoepidemiological study. *Cad Saúde Pública.* 2004;20(4):1004–1013.
36. Liu X, Liew Z, Olsen J, Pedersen LH, Bech BH, Agerbo E, et al. Association of prenatal exposure to acetaminophen and coffee with childhood asthma: Prenatal Acetaminophen Exposure and Asthma. *Pharmacoepidemiol Drug Saf.* 2016 Feb;25(2):188–95.
37. Ersbøll AS, Hedegaard M, Damm P, Johansen M, Tabor A, Hegaard HK. Changes in the pattern of paracetamol use in the periconception period in a Danish cohort. *Acta ObstetGynecol Scand.* 2015 Aug;94(8):898–903.
38. Liew Z, Ritz B, Rebordosa C, Lee P-C, Olsen J. Acetaminophen Use During Pregnancy, Behavioral Problems, and Hyperkinetic Disorders. *JAMA Pediatr.* 2014 Apr 1;168(4):313.
39. Thompson JMD, Waldie KE, Wall CR, Murphy R, Mitchell EA, the ABC study group. Associations between Acetaminophen Use during Pregnancy and ADHD Symptoms Measured at Ages 7 and 11 Years. Hashimoto K, editor. *PLoS ONE.* 2014 Sep 24;9(9):e108210.
40. Persky V, Piorkowski J, Hernandez E, Chavez N, Wagner-Cassanova C, Vergara C, et al. Prenatal exposure to acetaminophen and respiratory symptoms in the first year of life. *Ann Allergy Asthma Immunol.* 2008;101(3):271–278.
41. Scialli AR, Ang R, Breitmeyer J, Royal MA. A review of the literature on the effects of acetaminophen on pregnancy outcome. *ReprodToxicol.* 2010 Dec;30(4):495–507.
42. Shaheen SO, Newson RB, Sherriff A, Henderson AJ, Heron JE, Burney PGJ, et al. Paracetamol use in pregnancy and wheezing in early childhood. *Thorax.* 2002 Nov;57(11):958–63.
43. Food and Drugs Administration. Prescription Drug Advertising: Content and Format for Labeling of Human Prescription Drugs. Rules and Regulations. 1979 Jun 26;44:37451.
44. Lopes LM, Francisco RPV, Zugaib M, others. Anti-inflamatórios e alterações cardíacas fetais. *Rev Bras Ginecol E Obstetrícia.* 2010;32(1):1–3.
45. Sousa JM, Araújo Neto MF, Partata AK. Ação anti-inflamatória da nimesulida e seu grau de hepatotoxicidade. *Rev Científica ITPAC [Internet].* 2016 [cited 2017 May 17];9(1). Available from: http://www.itpac.br/arquivos/Revista/77/Artigo_6.pdf
46. Daniel S, Koren G, Lunefeld E, Levy A. NSAIDs and spontaneous abortions - true effect or an indication bias?: Indication bias in drug safety cohort studies. *Br J Clin Pharmacol.* 2015 Oct;80(4):750–4.
47. Andrade AM de, Ramalho AA, Koifman RJ, Dotto LMG, Cunha M de A, Opitz SP. Fatores associados ao uso de medicamentos na gestação em primigestas no Município de Rio Branco, Acre, Brasil. *Cad Saúde Pública.* 2014 May;30(5):1042–56.
48. Kassada DS, Miasso AI, Waidman MAP, Marcon SS. Prevalence and factors associated with drug use in pregnant women assisted in primary care. *Texto Contexto - Enferm.* 2015 Sep;24(3):713–21.
49. Rocha RS, Bezerra SC, Lima JWO, Costa FS. Consumption of medications, alcohol and tobacco during pregnancy and assessment of teratogenic risks. *Rev Gaúcha Enferm.* 2013;34(2):37–45.