

Chronic Kidney Disease: Pattern of Admission and Outcomes in a Nigerian
Tertiary Institution.

ABSTRACT:

Background: Chronic kidney disease is a global issue with a rising incidence and hospitalization rate. It coexists with cardiovascular diseases and presents higher morbidity and mortality.

AIMS; We aim at examining the admission and outcomes of patients with chronic kidney disease admitted to medical wards over a 5 year period; (November 2018 to October 2023).

Study Design: This is a single-center retrospective descriptive study examining patients with chronic kidney disease admitted to medical wards over a 5 year period; from November 2018 to October 2023.

Place and Duration of Study: This study was carried out in the University of Uyo Teaching Hospital, Nigeria. Data of patients admitted into the medical ward from November 2018 to October 2023 (5 years period) were retrieved.

Methodology: Patients' case notes were retrieved from the hospital's records department. Demographic data such as age, sex, occupation, and religion were retrieved from the case notes. Other data that were retrieved included diagnosis, date of admission and discharge, and duration of stay.

Results: A total of 567 patients were admitted during the study period consisting of 237(41.8%) females and 330(58.2%) males. The mean age of the patients was 49.75 ± 15.44 years. Median duration of stay was 10 days. Diabetic Nephropathy was the leading cause of CKD 160(28.2%), followed by hypertensive Nephrosclerosis 152(26.8%).

Approximately 60.1% of the patients were discharged, while 19.6% left against medical advice and 20.3%. Patients with diagnosis of chronic kidney of unknown cause (CKDu) had a higher likelihood of dying (54.05%).

Conclusion: This study observed an increasing trend of CKD patients, with the majority presenting late to the hospital. It also revealed that diabetic nephropathy is gradually becoming the most common cause of CKD in this environment.

Keywords: Chronic kidney disease, Admission, Outcomes. Tertiary Institution.

1. INTRODUCTION:

Chronic kidney disease (CKD) is defined as a chronic progressive reduction in renal function characterized by glomerular filtration rate (GFR) of $\leq 60 \text{ ml/min } 1.73 \text{ m}^2$ and or present of structural or functional abnormalities lasting up to $\geq 3 \text{ months}$ [1] The commonest functional abnormality is proteinuria.

Chronic kidney disease is a global issue with a rising incidence and hospitalization[2]. It coexists with cardiovascular diseases and present higher mobility and mortality. Mortality due to chronic kidney is 20 times more common than in the general population[3]. This is attributed to the increasing prevalence of risk factors for CKD, such as hypertension, obesity and diabetes[4]. CKD affects an estimated 843.6 million people worldwide[4], which is approximately 10% of the global population[5]. A systematic analysis of 33 population-based studies highlights the significant differences in CKD prevalence by geographic region, classified by income level. The age-standardized prevalence of 8.6% and 9.6% in men and women in high-income countries, and 10.6% in men 12.5% in women in low- and middle-income countries[6]. However, a more recent systemic review and meta-analysis of 100 studies observed a global prevalence rate of 13.4% for CKD stages 1-5 and 10.6% for CKD stages 3-5.[7] The prevalence rates for CKD ranged from 6.4 to 8.7% in South Africa[8-9], 10.7–13.9% in sub-Saharan Africa [10-11] and 4.6–10.1% across the whole of Africa [10], [12]. A systematic review on the prevalence of chronic kidney in Nigeria showed a rate of 2.5% to 26% depending of the formula used in calculating the glomerular filtration rate[13]. This wide variation is attributed to lack of uniformity in their definition of CKD. In a community study, Ulasi et al observed a prevalence rate 11.4%[14].

With recent shift from communicable to non-communicable diseases in low and middle-income countries, the burden of chronic kidney disease also been increasing[15]. A study in Southwestern, Uganda regional referral hospital revealed a prevalence rate of 15.3%[16].

Global Burden of Disease (GBD) studies have shown that CKD has emerged as a leading cause of worldwide mortality[17-18]. A study conducted in a specialised University Hospital in Southwest Ethiopia found that chronic kidney accounted for 4.6% of admissions [19]. Meanwhile, 6.6% of patients admitted into medicals wards in a hospital in Lahore were attributed to chronic kidney disease[20]. In Nigeria, a study conducted in a secondary care hospital reported a high prevalence of non-communicable diseases with mortality rate of 19.9%.[21]. This study did not consider CKD, although there were a high prevalence of hypertension and diabetes which are considered the leading cause of CKD worldwide[22]. Understanding the disease pattern and outcomes is necessary for proper planning, prevention, and better outcomes.

This study aims to review the prevalence and outcomes of patients admitted with chronic kidney in our hospital's medical wards in a middle- income economy.

Methodology: This is single center retrospective descriptive study of patients with chronic kidney disease admitted into medical wards from November 2018 to October 2023. There are five medical wards at the University of Uyo Teaching Hospital: three male wards and two female wards, each with 14 beds, giving a total of 70 beds.

Patients' case notes were retrieved from the hospital's records department. Demographic data such as age, sex, occupation, and religion were retrieved from the case notes. Other data that were retrieved included diagnosis, date of admission and discharge, and duration of stay. The reasons for signing against medical advice was recorded for those who did so. The number of patients who died were retrieved. Patients' investigation results were also recorded.

Glomerular filtration rate was calculated using Chronic Kidney Disease Epidemiology Collaboration (CKD-EPI) formula[23]. Chronic disease staging was performed using Kidney Disease: Improving Global Outcomes (KDIGO) guidelines[24].

Data handling: Data was analyzed using SPSS version 23. Continuous variables were presented as mean with standard deviation. While categorical variables were represented as frequencies and percentage. Chi-square was used for calculation of P value. A p-value < 0.05 was considered as significant. Regression analysis was done to compare the association between 2 variables.

3. RESULTS AND DISCUSSION

RESULTS: Table 1 shows the socio-demographic characteristics of patients with chronic kidney disease. A total of 567 patients were admitted during the study period, consisting of 237(41.8%) females and 330(58.2%) males. Minimum 18 years, maximum 90 years with mean age of 49.75±15.44 years. Most of the admitted patients were within age range of 41-60 years 256(45.1%), followed by 21-40years, 155(27.3%), 61 years and above 144(25.4%), and those below 20 years 129(2.1%)

TABLE 1: SOCIO-DEMOGRAPHIC CHARACTERISTICS OF CHRONIC KIDNEY DISEASE PATIENTS

Socio-demography	Frequency(n=567)	Percentages	
Females	237	41.8	P=0.001
Males	330	58.2	
Age Range			
20 years and below	12	2.1	Age
21 - 40 years	155	27.3	Mean = 49.75
41 - 60 years	256	45.1	SD = ±15.44
61 years and above	144	25.4	Maximum = 90
		100.0	Minimum = 18

Figure 1 illustrate the yearly trend of admissions from November, 2018 to October, 2023. The highest number of admission occurred in 2023, with 159 (28.00%) followed by 2022, 118(20.80%), 2020, 104(18.30%), 2019, 78(13.80%) and 2021, 77(13.60%).

Figure 1. Admission trend from November 2018 to October 2023 with a dip only in 2021




Figure 2 depicts duration of hospital stay. The minimum stay was 1day, the maximum stay was 85 days, and median stay of 10 days. The majority of patients stayed in the hospital was between 1and10days, 301(53.10%), followed by 11-20 days, 164(28.92%), 21-30days 62(10.93%) 31-40 days,25(4.41%) and more than 40 days, 15(2.64%)

Figure 2: Duration of Hospital Stay.




Table2 presents the causes of CKD.

Diabetic Nephropathy was the leading cause of CKD 160(28.2%), followed by hypertensive nephrosclerosis 152(26.8%), HIV related nNephropathy 88 (15.5%) ,chronic glomerulonephritis 81(14.3), obstructive nephropathy32(5.6%). Chronic kidney disease of unknown origin (CKDu) 37(6.5%), and lupus nephritis 2(0.4%). While, chronic allograft dysfunction, focal segmental sclerosis, multiple myeloma nephritis, seronegative vacuities , mixed connective tissue disease and single left kidney with calculi contributed 1(0.2%) each.

Table 2.. Causes of chronic kidney disease among the patients

Causes of chronic kidney disease	Frequency	Percentages
CGN*	81	14.3
Chronic allograft dysfunction	1	0.2
CIN**	8	1.4
DM nephropathy	160	28.2
FSGS ***	1	0.2
HIV related Nephropathy	88	15.5
Hypertensive Nephrosclerosis	152	26.8
Lupus nephritis	2	0.4
Mixed connective tissue disease	1	0.2
Multiple myeloma, nephrotic syndrome	1	0.2
Nephrolithiasis	1	0.2
Obstructive nephropathy	32	5.6
Seronegative vasculitis	1	0.2
Single left kidney with calculi	1	0.2

CKDu****	37	6.5
Total	567	100.0

*CGN- chronic glomerulonephritis **CIN – chronic interstitial nephritis, *** Focal segmental glomerulosclerosis. **** CKDu- chronic kidney disease of unknown origin

Figure 3 illustrates the outcomes of patient’s admissions. A majority (60.1%) were treated successfully and discharged, while, 19.6% left against medical advice, and 20.3% died.

Figure 3: Outcome of patients admitted with chronic kidney disease




Figure 4 depicts death outcomes based on etiology of CKD. Patients diagnosed chronic kidney disease of unknown etiology (CKDu) had the highest mortality rate (54.05%), followed by those chronic interstitial nephritis(37.50%), hypertensive nephrosclerosis (20.39%), diabetic nephropathy(19.38%), HIV- related nephropathy(19.32%) and obstructive uropathy (18.75%). Chronic glomerulonephritis was least associated with the tendency to die.

Figure 4: Death outcome and etiology of chronic kidney disease




Table 3 presents the reasons why patients left the hospital against medical advice with financial constraints being the predominant factor (89.2%).

Table 3: Reasons for leaving against medical advice.(LAMA)

Reasons	Frequency (n=111)	Percentages (%)
Financial constraint	99	89.20
Home treatment	1	0.9
No improvement	3	2.7
No functioning dialysis machine	1	0.9
Personal reason	3	2.7
Strike	2	1.8
To another hospital	2	1.8

Table 4 shows that most of the patients were in CKD stage five 378(66.7%), followed by stage four 87(15.3%), stage three 40(8.8%), stage two 27(4.8%) and stage one 25(4.4%) .It also shows the mortality outcomes across different s CKD. Stages. Out of 25 patients in stage 1, 1(0.2%) died, stage 2 , 3(0.6%) out of 27 patients, stage 3a , 2(0.4%) out of 22 patients , stage 3b, 5(1.0%) of 28, stage 4, 17(3.5%) out of 83 and stage 5, 62(12.9%) out of 382. ($P= 0.396$).

Table 4. Stages of chronic and death outcomes kidney disease

CKD Stages (n=567)

	Stage 1	Stage 2	Stage 3a	Stage 3b	Stage 4	Stage 5	P=0.396
	F(%)n=25	F(%)n=27	F(%)n=22	F(%)n=28	F(%)n=83	F(%)n=382	
Death							
No	24(4.2)	24(4.2)	20(3.5)	23(4.0)	66(11.6)	320(56.4)	
Yes	1(0.2)	3(0.6)	2(0.4)	5(1.0)	17(3.5)	62(12.9)	

Table 5 Death outcomes of patients who received dialysis versus those who did not. This table compares the death outcomes of patients who received dialysis versus those who did not. Of the 180 patients (31.7%) who had dialysis, 38 (6.7%) died. Meanwhile, of the 387 patients (68.3%) who did not have dialysis, 77 (13.6%) died ($P=0.73$).

Table 5 Shows relationship of death outcome and dialysis:

	Dialysis n=567		
	No	Yes	
Death	F(%) n =387	F(%) n=180	P=0.73
No	310(54.0)	142(25.0)	
Yes	77(13.6)	38(6.7)	

Discussion: The study shows that more males were admitted than females. This observation, however, is not consistent with other studies that have found CKD to be more prevalent in females than males.[13], [25] This discrepancy may be attributed to lifestyle factors that expose males to greater risks for chronic kidney disease, such as diabetes and hypertension. Additionally, men may prioritize raising funds for their families and spend more time working, often neglecting their health. Another possible explanation is the higher incidence of obstructive nephropathy in men, which can result from conditions like benign prostatic hyperplasia (BPH) and prostate cancer, both of which occur solely in men. The study also revealed that most of the patients were between the ages of 21-60. This contrasts with developed economies, where CKD is more prevalent in older age groups [5],[7]. This difference may be due to the high prevalence of chronic glomerulonephritis (CGN) and HIV associated nephropathy among the young and middle population in low-income and middle-income economies like ours. CGN primarily affects the young and middle aged, and it has been identified as one of the leading causes of chronic kidney disease in Nigeria[26-27]. In this study, HIV-related nephropathy and CGN were the 3rd and 4th leading

causes of CKD. A previous study in Uyo noted that HIV-related kidney disease was the leading cause of CKD[28]. The high prevalence of CKD in the younger and middle-aged groups, as observed in this study, poses a significant burden on the low and middle-income economies, which are ill-equipped to manage the long-term consequences. This age group represents the most productive segment of population, and without urgent intervention, the economic impact could be severe. Therefore, concerted efforts should be made by the government, private organizations, and non-governmental organizations to prioritize prevention.

The study observed an increasing trend of hospital admissions, with the exception of a small dip in 2021(Figure1). This rising trend may be attributed growing prevalence of non-communicable diseases in low- income middle- income economies[29] driven by increasing rates of diabetes and hypertension, both of which were identified as the leading causes of CKD in this study. The decline in admissions in 2021 could be explained by the general decrease in hospital admission observed post covid-19, as noted in other studies[30- 31]. Covid-19 was first diagnosed in Nigeria on February 27, 2020, in 44-year old Italian Citizen February 27, 2020. This led to a drop in hospital admission between 2020 and 2021[31]. The median length of stay among patients was 10 days, which is lower than 14 days reported in study from Port Harcourt[32]. The disparity may be due to inclusion of all patients with renal diseases, including those with acute kidney injury in Port Harcourt study. We observed that diabetes nephropathy was the leading cause of chronic kidney disease, followed by hypertensive nephrosclerosis, HIV related nephropathy and chronic glomerulonephritis. Diabetes mellitus remains the leading cause CKD worldwide [7], [33]. However, this finding contrasts with other Nigeria studies, where CGN and hypertension were identified as the leading cause of CKD [34]. Most of those studies were community-based studies, while ours focused on hospitalized patients with CKD. Chronic glomerulonephritis is often linked to infectious diseases like streptococcal throat and skin infections. Therefore, with reduction in infectious diseases in low and middle income-economies[29], the prevalence of CGN is also likely to reduce.

The prominence of diabetic nephropathy in this study may be attributed to the shift from communicable to non-communicable diseases in low-income and middle-income economies driven by a transition from the traditional lifestyle to more westernized habits [29]. Chronic kidney disease of unknown origin (CKDu) was seen in 6.5% of the patients. Patient is said to have CKDu after all the common etiologies of CKD has been excluded, and it predominantly affects young and middle-aged adults, particularly males engaged work, such as agriculture and manual labor[35]. Patients with CKDu typically have minimal or no proteinuria, are non-diabetic and have either normal blood pressure or only mildly hypertensive[35-36]. CKDu is increasingly recognized as a growing public health concern affecting working age population. The 6.5% prevalence observed in this study is lower than the 20.5% found by Suliaman et al in Northern Nigeria [37]. Other studies in Nigeria revealed that up to 21.5% of all renal admissions and 15% of renal deaths were attributable to CKDu[14],[32].

Over the review period 60.1% of patients were discharged, a higher rate than 53.3% observed than the study renal admission in Port Harcourt.[32]. The mortality rate in this study was 20.3% which aligns with earlier study in Nigerian tertiary institution [32]. Alarming, 114(19.6%) of patients left against medical advice (LAMA), a higher rate than reported in most other studies[32],[38], [39-40].The primary reason for LAMA was financial constraints, accounting for 99(89.2%) of cases. Other reasons included lack of improvement, personal reason, strike by medical personnel, preference for other hospitals and alternative treatment at home. Financial difficulties as the cause of LAMA have been noted in other Nigerian studies [39 -40]. This is due likely because healthcare in Nigeria is mostly out-of-pocket. The Federal Government of Nigeria only cover for six sessions of

dialysis for the enrollees in the National Health Insurance scheme. There is pressing need for the government to include all patients with CKD under the National Health Insurance scheme (NHIS) and take full responsibility for their medical care. States government should also consider to establishing state insurance scheme for their residents.

Most patients in this presented with CKD stage 5, requiring urgent dialysis. This is consistent with earlier studies in Nigeria[27], [41]. This may be because our patients often opt for alternative care from traditional and spiritual care -givers before presenting at a late stage for orthodox care.[42]. Additionally, CKD patients often remains asymptomatic until late stages of the disease. Therefore, there an urgent need for public health campaigns and educational efforts through schools, churches, town criers and mass media at regular interval to promote early detection and intervention.

Patients diagnosed with CKDu were more likely to die than those with other causes of CKD (54.05%). An earlier study had observed HIV-related kidney was the leading cause of death among renal patients with 15% of death attributed to CKDu[32]. This high mortality rate associated CKDu may be due to its insidious onset and lack of symptoms until the disease is advanced. CKDu often progresses rapidly and can be fatal due genetic and environmental factors[43]. Chronic interstitial nephritis, hypertensive nephrosclerosis were also associated with high mortality in this study.

There was no statistically significant difference between the various stages of CKD and death outcomes in this study, which may be attributed to late presentation of patients. Similarly, dialysis also did not significantly affect mortality outcome, possibly due to the inability of many patients to sustain long-term dialysis and delays in initiating treatment due to financial constraints. An earlier study at the same center indicated that majority of the patients were unable maintain regular hemodialysis[44].

Limitation: A major limitation of this study is the manual storage of patient records. Not all patient folders were retrievable, and some folders were damaged or incomplete, resulting in the loss of vital information. .

4. CONCLUSION

This study observed an increasing trend in the number of patients with CKD, with the majority presenting late to the hospital. It also highlighted that diabetic nephropathy is gradually becoming the most common cause of CKD in low- and middle-income countries. Additionally, the study revealed an increasing number of patients leaving the hospital against medical advice due to financial constraints.

There is a need to place greater emphasis on prevention, early detection, and control of CKD risk factors. Furthermore, the government should consider expanding the National Health Insurance Scheme to cover patients with chronic kidney disease, improving access to care and reducing financial burdens

CONSENT

Not applicable (retrospective study)

ETHICAL APPROVAL All authors hereby declare that the research was performed in accordance with the ethical standards laid down in the 1964 Declaration of Helsinki.”

Ethical approval was sought for and obtained from the University of Uyo Health Research Ethics Committee. UUTH NHREC Registration Number- NHREC/24/06/22/UUTH with HREC Protocol assigned number UUTH/AD/S/96/VOL.XXI/854.

REFERENCES:

1. Levey AS, Atkins R, Coresh J, Cohen EP, Collins AJ, Eckardt KU, et al. Chronic kidney disease as a global public health problem: approaches and initiatives - a position statement from Kidney Disease Improving Global Outcomes. *Kidney Int* 2007;72(3):247–59.
2. Cesarino CB, Borges PP, Ribeiro RDCHM, Ribeiro DF, Kusumota Assessment of cardiovascular risk in patients with chronic kidney disease according to Framingham's criteria L. *Acta Paul Enferm* 2013;26(1):101–7.
3. Barbosa DA, Gunji CK, Bittencourt AR de C, Belasco AGS, Diccini S, Vattimo F, et al. Co-morbidity and mortality of patients in dialysis treatment. *Acta Paul Enferm* 2006;19(3):304–9.
4. Jager KJ, Kovesdy C, Langham R, Rosenberg M, Jha V, Zoccali C. A single number for advocacy and communication-worldwide more than 850 million individuals have kidney diseases. *Kidney Int* 2019;96(5):1048–50.
5. Kovesdy CP. Epidemiology of chronic kidney disease: an update 2022. *Kidney Int Suppl* 2022;12(1):7–11.
6. Mills KT, Xu Y, Zhang W, Bundy JD, Chen CS, Kelly TN, et al. A systematic analysis of worldwide population-based data on the global burden of chronic kidney disease in 2010. *Kidney Int* 2015;88(5):950–7.
7. Global Prevalence of Chronic Kidney Disease – A Systematic Review and Meta-Analysis | PLOS ONE [Internet]. [cited 2024 Sep 6]; Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0158765>
8. Kaze AD, Ilori T, Jaar BG, Echouffo-Tcheugui JB. Burden of chronic kidney disease on the African continent: a systematic review and meta-analysis. *BMC Nephrol* 2018;19(1):125.
9. Matsha TE, Yako YY, Rensburg MA, Hassan MS, Kengne AP, Erasmus RT. Chronic kidney diseases in mixed ancestry south African populations: prevalence, determinants and concordance between kidney function estimators. *BMC Nephrol* 2013;14:75.
10. ElHafeez SA, Bolignano D, D'Arrigo G, Dounousi E, Tripepi G, Zoccali C. Prevalence and burden of chronic kidney disease among the general population and high-risk groups in Africa: a systematic review. *BMJ Open* 2018;8(1):e015069.
11. Matsha TE, Erasmus RT. Chronic kidney disease in sub-Saharan Africa. *Lancet Glob Health* 2019;7(12):e1587–8.

12. ISN-Global Kidney Health Atlas [Internet]. *Int. Soc. Nephrol.* [cited 2024 Sep 6]; Available from: <https://www.theisn.org/initiatives/global-kidney-health-atlas/>
13. Chukwuonye II, Ogah OS, Anyabolu EN, Ohagwu KA, Nwabuko OC, Onwuchekwa U, et al. Prevalence of chronic kidney disease in Nigeria: systematic review of population-based studies. *Int J Nephrol Renov Dis* 2018;11:165–72.
14. Ulasi II, Ijoma CK. The enormity of chronic kidney disease in Nigeria: the situation in a teaching hospital in South-East Nigeria. *J Trop Med* 2010;2010:501957.
15. Noncommunicable Diseases and Injuries in Latin America and the Caribbean: Time for Action | PLOS Medicine [Internet]. [cited 2024 Sep 7]; Available from: <https://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.0030344>
16. Kalima NA, Gabriel BK, Muhindo R, Musingo A. Chronic kidney disease in patients admitted to the medical ward of Mbarara Regional Referral Hospital in southwestern Uganda: Prevalence and associated factors. *Int J Med Biomed Res* 2015;4(2):107–16.
17. Global, regional, and national age-sex specific all-cause and cause-specific mortality for 240 causes of death, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2015;385(9963):117–71.
18. Rhee CM, Kovesdy CP. Spotlight on CKD deaths—increasing mortality worldwide. *Nat Rev Nephrol* 2015;11(4):199–200.
19. Ali E, Woldie M. Reasons and Outcomes of Admissions to the Medical Wards of Jimma University Specialized Hospital, Southwest Ethiopia. *Ethiop J Health Sci* 2010;20(2):113.
20. Rahim A, Khurshid N, Mukhtar MU. Pattern of diseases in medical wards of a tertiary care hospital of Lahore: a retrospective study. *Discov Med* 2024;1(1):1–6.
21. Adedapo ADA. Morbidity and mortality patterns of medical admissions in a Nigerian secondary health care hospital. *Afr J Med Med Sci* 2012;41(1):13–20.
22. Olanrewaju TO, Aderibigbe A, Popoola AA, Braimoh KT, Buhari MO, Adedoyin OT, et al. Prevalence of chronic kidney disease and risk factors in North-Central Nigeria: a population-based survey. *BMC Nephrol* 2020;21(1):467.
23. CKD-EPI Creatinine Equation (2021) | National Kidney Foundation [Internet]. [cited 2024 Sep 23]; Available from: <https://www.kidney.org/ckd-epi-creatinine-equation-2021>
24. Stevens PE, Ahmed SB, Carrero JJ, Foster B, Francis A, Hall RK, et al. KDIGO 2024 Clinical Practice Guideline for the Evaluation and Management of Chronic Kidney Disease. *Kidney Int* 2024;105(4):S117–314.
25. A systematic analysis of world-wide population-based data on the global burden of chronic kidney disease in 2010 - PMC [Internet]. [cited 2024 Sep 6]; Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4653075/>
26. O S, Jacob AO, Sharma S. Etiology of Chronic Kidney Disease in Nigeria and Management Challenges. *Indian J Clin Pract* 2023;33(9):10–4.

27. Adejumo OA, Akinbodewa AA, Okaka EI, Alli OE, Ibukun IF. Chronic kidney disease in Nigeria: Late presentation is still the norm. *Niger Med J J Niger Med Assoc* 2016;57(3):185–9.
28. AIA AE Ekrikpo UE, Udo. Changing Aetiologies of End Stage Kidney Disease in a Resource Poor Environment, What Is the Way Forward? [Internet]. *World J. Biomed. Res.* 2014 [cited 2024 Sep 10]; Available from: <https://wjbmr.org/articles/14/09/changing-aetiologies-of-end-stage-kidney-disease-in-a-resource-poor-environment-what-is-the-way-forward/>
29. Bollyky TJ, Templin T, Cohen M, Dieleman JL. Lower-Income Countries That Face The Most Rapid Shift In Noncommunicable Disease Burden Are Also The Least Prepared. *Health Aff Proj Hope* 2017;36(11):1866–75.
30. Rennert-May E, Leal J, Thanh NX, Lang E, Dowling S, Manns B, et al. The impact of COVID-19 on hospital admissions and emergency department visits: A population-based study. *PLoS ONE* 2021;16(6):e0252441.
31. Shehu M, Ihekaike M, Jimoh AO, Shehu H, Mava Y, Esegbe EE, et al. Impact of COVID-19 on Admission into the Department of Paediatric Bingham University Teaching Hospital (BHUTH) Jos. *J Adv Med Med Res* 2022;85–91.
32. Wachukwu CM, Emem-Chioma PC, Wokoma FS, Oko-Jaja RI. Pattern and outcome of renal admissions at the University of Port Harcourt Teaching Hospital, Nigeria: A 4 years review. *Ann Afr Med* 2016;15(2):63–8.
33. Nath JD, Kashem A. Etiology and Frequency of Hospital Admissions in Maintenance Hemodialysis Patients in Chronic Kidney Disease. *Saudi J Kidney Dis Transplant* 2019;30(2):508.
34. Ulasi II, Ijoma CK, Onodugo OD, Arodiwe EB, Ifebunandu NA, Okoye JU. Towards prevention of chronic kidney disease in Nigeria: a community-based study in Southeast Nigeria. *Kidney Int Suppl* 2013;3(2):195–201.
35. Anupama YJ, Sankarasubbaiyan S, Taduri G. Chronic Kidney Disease of Unknown Etiology: Case Definition for India – A Perspective. *Indian J Nephrol* 2020;30(4):236–40.
36. Correa-Rotter R, Wesseling C, Johnson RJ. CKD of unknown origin in Central America: the case for a Mesoamerican nephropathy. *Am J Kidney Dis Off J Natl Kidney Found* 2014;63(3):506–20.
37. Sulaiman MM, Shettima J, Ndahi K, Abdul H, Baba MM, Ummate I, et al. Chronic kidney disease of unknown origin in Northern Yobe, Nigeria: Experience from a regional tertiary hospital in northeastern Nigeria. *Borno Med J [Internet]* 2019 [cited 2024 Sep 12];16(1). Available from: <https://www.ajol.info/index.php/bomj/article/view/257065>
38. Das BR, Medhi AH, Das N, Kakoti G, Nath G, Sarkar AH. Admission Pattern and Outcome of Patients Attending a Tertiary Care Hospital. *Int J Curr Res Acad Rev* 2016;4(4):21–8.

39. Fadare J, Babatunde O, Olanrewaju T, Busari O. Discharge against medical advice: Experience from a rural Nigerian hospital. *Ann Niger Med* 2013;7:60–5.
40. Eze B, Agu, Nwosu. Discharge against medical advice at a tertiary center in southeastern Nigeria: sociodemographic and clinical dimensions. *Patient Intell* 2010;27.
41. Okaka EI, Adejumo OA, Akinbodewa AA. Late Referral and Associated Factors among Chronic Kidney Disease Outpatients in Southern Nigeria. *Ann Afr Med* 2020;19(1):47.
42. Chukuezi CO, Anelechi AB. Factors Associated with Delay in Seeking Medical Care among Educated Nigerians. 2009;
43. Weaver VM, Fadrowski JJ, Jaar BG. Global dimensions of chronic kidney disease of unknown etiology (CKDu): a modern era environmental and/or occupational nephropathy? *BMC Nephrol* 2015;16:145.
44. Akpan EE, Ekrikpo UE, Effa EE, Udo AIA, Umoh VA. Demographics, Cost, and Sustainability of Haemodialysis among End-Stage Kidney Disease Patients in Southern Nigeria: A Single-Center Study. *Niger Med J J Niger Med Assoc* 2020;61(6):307–11.

ABBREVIATIONS:

CGN-Chronic Glomerulonephritis
CKD- Chronic Kidney Disease
CIN – Chronic Interstitial Nephritis
CKDu- Chronic Kidney Disease of Unknown Origin
FSGS- Focal Segmental Glomerulosclerosis
LAMA – Left Against Medical Advice