

Determination of Astrocyte Reaction Using Glial Fibrillary Acidic Protein (GFAP) Following Aluminium Chloride Induced Hippocampus Damage Of Adult Wistar Male Rats Treated With Ethanolic Extracts Of *Carpolobialutea* At Different Doses

ABSTRACT

Background: Aluminium is a well-established neurotoxicant involved in the etiology of neurodegenerative diseases. *Carpolobia lutea*, commonly called cattle stick or poor man's candle belongs to the plant family polygalaceae. It is a small tree native to West and Central tropical Africa and it is one of the medicinal plants that play a major role in the health care system of developing countries such as Nigeria.

Aim of the Study: This study was aimed at investigating astrocyte reaction in aluminium chloride induced neurotoxicity in the hippocampus in adult male wistar rats treated with ethanolic extracts of *C.lutea* at different doses.

Materials and Methods: Thirty wistar rats weighing 180-200g were used for this study. The animals were randomized into five groups of six rats each. Group A rats received only animal feed. Groups B,C,D and E were given 100mg/kg bw of aluminium chloride intraperitoneally five times a week for three weeks to induce neurotoxicity. In order to reverse the neurotoxicity, Group C was treated with 10mg/kg bw of donepezil as standard drug while group D and E were treated with 200mg/kg and 400mg/kg of *Carpolobia lutea* respectively for 14 days.

Histopathological study was done on the hippocampus of the rat brain and thereafter hematoxylin and eosin staining were carried out. Immunohistochemical studies for GFAP was done using Novocasa™Novolink™polymer detection system and appropriate primary monoclonal antibodies. Image J cell counter tool was used to note the number of GFAP positive cells across all groups.

Results: The result from our histopathological study shows that lower dose(200mg/kg/bw) of ethanolic extract of the leaf of *Carpolobia lutea* have better protection on the cytoarchitecture of the CA₁, CA₃, and dentate gyrus of the hippocampus than 400mg/kg/bw of the plant extract. From our study, it is evident that 200mg/kg of *Carpolobia lutea* reduced the GFAP immunoreactivity in the hippocampus of AlCl₃ induced neurotoxicity. When we compared the result of 200mg/kg/bw of *C.lutea*with 10mg/kg per body weight of donepezil(standard drug),it was observed that *C.lutea* had a slight reduced GFAP immunoreactivity than the standard drug(donepezil). The lower dose, 200mg/kg of ethanolic extract of the leaf of *Carpolobia lutea*have a better neuroprotective benefit than the standard drug on the hippocampus

Conclusion: In conclusion, 200mg/kg of *Carpolobia lutea* have the tendencies of protecting the neurons in the hippocampus from degenerating as a result of aluminium chloride induced neurotoxicity compared to the standard drug (10mg/kg donepezil).

Keywords: Glial Fibrillary Acidic Protein, Hippocampal Damage, Wistar Male Rats, Ethanolic Extracts, *Carpolobia lutea*

1. INTRODUCTION

1.1 Background of the Study

“Medicinal plants play a major role in the health care system of developing countries such as Nigeria. Despite their high usage, most of these plants lack information on, active chemical constituent, quality, clinical studies, safety and efficacy of which *Carpolobia lutea* is one of them” (Odigie et al., 2003).

“*Carpolobia lutea*, commonly called cattle stick or poor man’s candle belongs to the plant family polygalaceae” (Akhondzadehand Abbas, 2006). “It is a small tree native to West and Central tropical Africa. It is common in rainforest and Guinea savannah of Sierra Leone and Cameroon. This shrub reaches up to 5m in height and occurs as a dense

overgrowth or an evergreen shrub or small tree. The common names, which the plant is known include cattle stick (English), AbekpokIbuhu (Eket), Ikpafum, Ndiyan, Nyayanga (Ibibio), Agba or Angalagala (Igbo) and Egbo oshunshun (Yoruba)” (Essien et al., 2011).

“Herbalists in Nigerian tribes use the essence of the root as an aphrodisiac and the treatment of genitourinary infections, gingivitis, and waist pains. The tree extract could pave the way for new drugs to tackle patient symptoms but without the unwanted side effects associated with some current treatments” (Ajiwhen and Bisong, 2013).

“Neurodegenerative diseases represent a huge health burden globally, placing pressure on health services and having a negative impact on the lives of patients and their families” (Ballard et al., 2011).

“Researchers and drug companies are racing to discover new treatments for these disorders and have begun looking to plant extracts as a potential source of novel drugs. In patients with Alzheimer’s disease and other diseases such as Parkinson’s disease and myasthenia gravis, the activity of the neurotransmitter acetylcholine, is increased, leading to problems with memory and attention” (Syad and Devi, 2014).

“Current drugs — called acetylcholinesterase inhibitors — reduce the normal breakdown of acetylcholine. Extensive research is underway to find new versions of these drugs but with additional beneficial properties” (Parihar and Hemnani, 2004).

“According to a study published in Pharmacognosy Review, It is an accepted and commonly utilized herbal booster of libido. It is used to cure male infertility and to boosts libido thereby augmenting male sexual functions or it is used to induce penile erection, and enhance male

virility (Tundis et al., 2016). The chewing stick prepared from the stem and root of *Carpolobia lutea* is patronized because it boosts male sexual performance” (Mesulam et al., 2002).

“The leaf essential oil contains a variety of terpenoids, while polyphenols and triterpenoid saponins have been isolated from the root and leaf extracts respectively. Other ethnomedicinal uses include curing of stomach ailments, rheumatism, fever, pains, insanity, dermal infection, venereal diseases; to promote childbirth; and as a taeniafuge and vermifuge” (Manoharan et al., 2007; Lockridge 2015).

“It has also been reported to possess other anti-inflammatory, anti-arthritic, antimicrobial, antimalarial, and analgesic properties. This could be particularly important in Alzheimer’s disease as there is more evidence emerging that Alzheimer’s patients have inflammation in the brain” (Duysen et al., 2007).

“Aluminium is a well-established neurotoxicant involved in the etiology of neurodegenerative diseases” (Sun et al., 2009). “It is an abundant metal on earth with easy access to the human body through agrochemicals, water, food additives, utensils, deodorants and drugs. In the brain, aluminum predominantly accumulates in the hippocampus and frontal cortex, regions known to be particularly susceptible in Alzheimer’s disease” (Flaten, 2001; Campbell, 2002). It induces misfolding of cytoskeleton proteins which leads to the formation of amyloid beta plaques and tau neurofibrillary tangles (El-Sebae et al., 1993) in the brain. “Aluminium supplementation causes neurodegeneration and apoptotic neuronal loss along with cognitive dysfunction, as it is a potent cholinotoxin. Normally, cholinergic activity is necessary for the acquisition and retrieval of learning and memory skills” (Domingo, 2006). “Hence patients with AD normally demonstrate impaired performance in various cognitive tasks. Various animal studies have also shown that

prolonged exposure to aluminium can cause neurochemical, neurobehavioral and neuropathological changes in the brain, which impair the learning ability of the rats” (Kaur and Gill, 2006; Prema et al., 2017).

1.2 Aim of the Study: This study was aimed at investigating astrocyte reaction in aluminium chloride induced neurotoxicity in the hippocampus in adult male wistar rats treated with ethanolic extracts of *C.lutea* at different doses

2. MATERIALS AND METHOD

2.1 Extraction Procedures

The leaves of *Carpolobia lutea* were removed from their stalks. The leaves were oven dried for 72 hours and pulverized. Dried sample was ground into powder mechanically using manual grinder.

Thereafter, 400g of the macerated plant powder was soaked in 1000ml of absolute alcohol and kept in a container for 48 hours. The mixture was shaken vigorously at intervals for another 2 hours, to allow complete extraction.

The resulting mixture was rapidly filtered through whatmann filter paper to obtain a homogenous filtrate.

This filtrate was concentrated in a vacuum at low temperature (37-40⁰C). The extract was later reconstituted in distilled water at a concentration of 1g/ml before administration. The extract was refrigerated until use.

2.2 Standard Drug

A standard drug known for its therapeutic management of neurotoxicity in the hippocampus, namely donepezil was used to ascertain and compare with the potency of ethanolic extract of *Carpolobia lutea* against aluminium chloride induced neurotoxicity . A tablet of this drug was dissolved in 20ml of distilled water and mixed thoroughly.

2.3 Recruitment of the Study Population

Thirty Wistar rats weighing between 180-200g were used for this study. The animals were obtained from the animal house of the Department of Biological Sciences of Rivers State University. They were housed in standard cages and left to acclimatize for 14 days under natural conditions in the animal house of the Department of Human Anatomy of Rivers State University before the commencement of the experiment. The animals were fed with vitakraft complete nutrition rat feed and water ad libitum.

2.4 Study Design

The animals were randomized into five groups (A ,B ,C, D, & E) of six rats each. Rats in group A received only animal feed and water. All rats in groups B, C, D, & E were injected with

aluminium chloride (AlCl_3) at 100mg/kg bw intraperitoneally five times a week for three weeks (Aljarari, 2023).

The ethanolic extract of *Carpolobia lutea* and donepezil were administered orally by gavage.

Group B received 100mg/kg bw of aluminium chloride (Aljarari, 2023).

Group C received 10mg/kg bw of donepezil (Shin et al., 2018).

Group D received 200mg/kg per body weight of ethanolic extract of *Carpolobia lutea*.

Group E received 400mg/kg per body weight of ethanolic extract of *Carpolobia lutea*.

The administration of *Carpolobialutea* and donepezil lasted for 14 days. On the 12th, 13th and 14th day, the rats were subjected to neuro-behavioural tests after which they were sacrificed by cervical dislocation on the 15th day.

2.5 Surgical procedures

At the end of behavioural studies, animals were euthanized by cervical dislocation. Blood was quickly obtained by cardiac puncture and processed for antioxidant studies. Also, their brains were rapidly excised. Whole brain tissues of two animals per group were fixed in 10% neutral buffer formalin. All tissues were processed for rapid routine paraffin wax embedding for histopathological and immunohistochemical studies.

2.6 Histopathological Studies

After sacrifice, the brains of two rats from each group were fixed whole in 10% formal saline. After 48 hours, hippocampus was excised from each rat brain for histological assessment using H&E.

The fixed tissues were dehydrated to remove water which is not miscible with paraffin (the embedding medium). The tissues were placed in ascending grades of alcohol (70%, 80%, 90%, and absolute). This was done to prevent water from rushing out of the tissue, which may distort and damage the cell structure. The time for each grade of alcohol was one hour and in absolute (100%) it was changed twice, one hour each time. After dehydration the tissues were cleared in 2 changes of xylene for 30 mins each to remove alcohol which removed water from tissue. This was done because paraffin used for impregnation and embedding is not miscible with alcohol.

The tissues were then passed through four changes of molten paraffin wax at constant temperatures of 46-68°C in an oven of paraffin bath. This was to prepare the tissue for the embedding medium and to confirm firmness to it. This will facilitate the process of sectioning. Metal blocks were taken and filed with paraffin wax and tissues were placed in it immediately with forceps, the face to be cut facing downward. When the paraffin cools, a thin scum of solid paraffin is formed on the bottom of the block which was now immersed in water to solidify and then removed for sectioning.

The solid paraffin blocks were then taken to the rotary microtome where excess paraffin wax was first trimmed off and the tissues cut at 5µm. Paraffin sections were taken to a water bath set at 45°C to straighten it. The side of the glass slide to receive the section was made sticky by rubbing with egg albumin. The paraffin sections were allowed to float in the water bath to straighten out the wrinkles. Water was drained off and the slide was put in an incubator for the sections to be completely fixed on the slide and allow to dry.

2.7 Staining: Hematoxylin and eosin staining technique

The procedure for H&E staining as described by Robert et al., (2014), Bancroft and Stevens, (2005) was adopted. The sections were dewaxed in xylene, two changes of xylene for 2 minutes each and afterwards rehydrated in descending grades of alcohol, 100%, 95%, 90%, 70%, 50% ethanol for 2 minutes each.

Tissues sections were then rinsed in distilled water and stained in haematoxylin for 10-15 minutes. Afterwards sections were rinsed in distilled water for 2-3 minutes and examined to confirm sufficient degree of staining. Excess stain was removed or differentiated in 1% HCL acid alcohol for a second or two as the acid breaks the mordant dye linkage.

The sections are again rinsed in distilled water for 2-3 minutes to regain the blue colour. Sections were then stained in 1% aqueous eosin for about 3-5 minutes. Surplus stain was washed off in distilled water.

Stained sections were then mounted in distrene plasticizer xylene (DPX) using clean glass cover slide and placed under the microscope for examination. Photomicrographs were taken using the photographic light microscope (National optical).

2.8 Immunohistochemical Studies

Immunohistochemical studies for GFAP was carried out using NovocastraTMNovolinkTM polymer detection system and appropriate primary monoclonal antibodies.

2.8.1 Immunohistochemistry Using ImmPRESSTM HRP Polymer System

The non-biotin, enzymatic, one-step detection kit, ImmPRESSTM Polymerized Reporter Enzyme Staining System (Vector® Labs, USA), provides very high sensitivity staining with very

minimal background interference in immunohistochemical applications. The ImmPRESS™ Reagent uses an innovative, exclusive approach to conjugate horseradish peroxidase (HRP) micropolymers to affinity-purified, extensively cross-adsorbed secondary antibodies.

2.8.2 Reagents supplied (#MP-7401; Vector® Labs, USA):

- ImmPRESS™ (Peroxidase) Polymer Anti-Rabbit IgG Reagent (made in horse, ready-to-use)
- 2.5% Normal Animal (Horse) Serum for blocking (ready- to-use)

2.8.3 ImmPRESS™ Immunohistochemical Detection:

The ImmPRESS™ Reagent is ready-to-use. It requires no mixing or titrating of the ImmPRESS™ reagent to obtain optimal immunohistochemical staining. The staining procedure is performed at room temperature. For optimal performance, the ImmPRESS™ Reagent is equilibrated to room temperature before use. Phosphate buffered saline (PBS) was used as wash buffer.

2.9 Staining Protocol:

The protocol was performed as previously described (Erukainure et al., 2019; Ijomone et al., 2018).

- Paraffin embedded sections were deparaffinized with xylene, and rehydrated through descending grades of ethanol (100%, 95%, 70 % ethanol) and taken to water.
- Heat-mediated antigen retrieval was performed using a citrate-based antigen unmasking solution, pH 6.0 (Vector®, Burlingame, CA, USA; #H3300) in a steamer for 30 mins.
- Wash sections in PBS for 2 mins

- Endogenous peroxidase blocking in 0.3 % hydrogen peroxide solution in PBS for 10 mins.
- Wash sections in PBS for 2 mins
- Sections were incubated in animal free blocker for 20 mins for protein blocking.
- Sections were then incubated at room temperature for 2 hours in primary rat antibodies: GFAP (ThermoFisher, USA; #16825-1-AP), Sections were incubated in ImmPRESS™ HRP Anti-Rabbit IgG (Peroxidase) Polymer Reagent, made in horse for 30 mins
- Wash in PBS for 5 mins x 2
- Colour is developed with DAB Peroxidase (HRP) Substrate Kit (Vector® Labs, USA)
- Sections were rinsed well in tap water
- Sections were counter-stained in hematoxylin
- Sections dehydrated through ascending grades of ethanol (70%, 95%, 100%), cleared in Xylene and mounted with Permount (Fischer Scientific, USA).

Sections without primary antibodies were similarly processed to control for immunohistochemistry procedures. No specific immunoreactivity was detected in control sections.

2.10 Photomicrography and Image Analysis

Stained sections were viewed under a Leica DM750 Digital Light microscope and digital photomicrographs of slides were taken by an attached Leica ICC50 camera. Image Analysis and Processing for Java (Image J), a public domain software sponsored by the National Institute of Health (USA), was used to analyze and quantify photomicrographs. Image J cell counter tool was used to identify and quantify number of intact and degenerating neurons in H&E stained sections. Also the Image J cell counter tool was used to note the number of GFAP positive cells.

2.11 Statistical Analysis

All other data obtained were analysed using one way ANOVA followed by student newman-keuls (SNK) for post test. All behavioural studies were analysed using two-way repeated measures (2-RM) ANOVA, with time as the repeated measures variable.

3. RESULTS

3.1 Haematoxylin And Oesin (H & E) Staining Results


The histological studies were carried out using the Haematoxylin and Oesin (H & E) method. Apart of the brain was studied, the hippocampus (CA1, CA3 and DG). The histology of the hippocampus, dentate gyrus and their components are presented in figures 1, 2 and 3.

The control group, show typical histological features of the hippocampal CA1 and CA3. There are three distinct layers: stratum radiatum, pyramidalis and oriens. The pyramidal layers are composed of large pyramidal neurons with large round nucleus and prominent nucleoli with glial cells interspaced within the neurons as can be seen in figures 1 and 2.

The negative control group and the treated groups were observed for features that characterize degenerating process in neurons, as can be seen in light microscopy study of H & E stained sections (Carless et al, 2007, German, 2011). These features include;


1. Prominent eosinophilic cytoplasm with or without shrunken nuclei
2. Pyknotic nuclei and
3. Neuron swelling and/ or vacuolation within the cytoplasm

The 100mg /kg of AlCl₃ group showed degenerative features characterized by pyknotic cells with deeply stained nuclei. There were obvious perinuclear spaces and vacuoles formed from pyknotic neurons and lost neurons respectively as can be seen in figures 1 and 2. The aforementioned degenerative features were also present, albeit mildly in the 400mg/kg of *Capolobia lutea* group when compared to control as seen in figures 1 and 2. Conversely, the 10mg/kg of donepezil and 200mg/kg of *C. lutea* showed no tissue alteration when compared to control.




Group of Figures 1: Histological changes in the hippocampal CA1 of experimental groups.

H&E 400x magnification. SR – stratum radiatum; SP – stratum pyramidalis; SO – stratum oriens; black arrows – intact pyramidal neurons; broken/dotted arrows – glial cells; red arrows – pyknotic neurons; red arrow heads – perinuclear spaces.




Groups of Figures 2: Histological changes in the hippocampal CA3 of experimental groups.

H&E 400x magnification. SR – stratum radiatum; SP – stratum pyramidalis; SO – stratum oriens; black arrows – intact pyramidal neurons; dotted/broken arrows – glial cells; red arrows – pyknotic neurons; red arrow heads – perinuclear spaces.



Control

100mg/kg AlCl₃



10mg/kg Donepezil

200mg/kg *C. lutea*

400mg/kg *C. lutea*

Group of Figures 3: Histological changes in the hippocampal DG of experimental groups.

H&E x400 magnification. PL – polymorphic layer; GL -granular layer; ML – molecular layer; black arrows – intact granular neurons; dotted/broken arrows – glial cells; red arrows – pyknotic neurons; red arrow heads – perinuclear spaces.

Dentate Gyrus as seen in figures 3, the control group exhibits typical histological characteristics; polymorphic, granular and molecular layers. The granular layer is made up of many neurons with small spherical nuclei and glial cells scattered among them.

The 100mg/kg of AlCl₃ group exhibit degenerative characteristic such as pyknotic cells with darkly coloured nuclei. Perinuclear gaps and vacuoles are clearly visible, generated by pyknotic neurons and lost neurons, respectively. The aforementioned degenerative characteristics are present, however weakly in the 400mg/kg *C. lutea* group just as it is in CA1 and CA3.

When compared to the control group, the 10mg/kg of donepezil and 200mg/kg *C. lutea* groups exhibit no tissue modification as can be seen in figure 3.

We can therefore infer that the lower dose of ethanolic extract of the leaf of *Carpolobia lutea* gave better protection on the cytoarchitecture of the hippocampus.


3.2 Results of Immunohistochemistry Studies

Using image J cell counter tool, Immunohistochemical analyses with one way ANOVA was used for this study.

3.2.1 GFAP – CA1

Immunohistochemical analysis with one-way ANOVA demonstrated significant changes in the number of GFAP-positive cells [$F(4, 47) = 8.249, p < 0.0001$] in the hippocampal CA1 region following AlCl₃ exposure in rats. Post-hoc analysis using Tukey's test revealed that rats subjected to 100mg/kg AlCl₃ ($p < 0.0001$) and 400mg/kg *C. lutea*, ($p < 0.01$) had considerably more GFAP-positive cells in hippocampal CA1 than in the control. In contrast, when compared

to 100mg/kg AlCl₃-exposed rats, the number of GFAP-expressing cells in the hippocampus CA1 of 10mg/kg donepezil- ($p < 0.01$) and 200mg/kg *C. lutea*-treated ($p < 0.0001$) rats was considerably lower. Further, there was an observable, but not significant reduction in the number of hippocampal CA1 GFAP-positive cells in the 200mg/kg *C. lutea*-treated group compared to 10mg/kg donepezil. Similar observation was made between 400mg/kg *C. lutea*-treated group and 100mg/kg AlCl₃-exposed rats.



Group of Figures 4: Immunohistochemical demonstration of GFAP in the hippocampal CA1 of rats exposed to experimental drugs. 400x magnification. Brown-stained cells are GFAP-expressing astrocytic cells.






Figure 5: Bar graphs depict the number of cells positive for GFAP in experimental rats. Each column represents mean \pm S.E.M. Data were analysed using one-way ANOVA followed by Tukey's post-test. ****p < .0001, **p < .01 versus Control; #####p < .0001, ##p < .01 versus 100mg/kg AlCl₃.

3.2.2 GFAP – CA3


Immunohistochemical analysis with one-way ANOVA demonstrated significant changes in the number of GFAP-positive cells [$F(4, 45) = 9.037, p < 0.0001$] in the hippocampal CA3 region following AlCl_3 exposure in rats. Post-hoc analysis with Tukey's test indicated that there was significantly more number of GFAP-positive cells in the hippocampal CA3 of rats exposed to 100mg/kg AlCl_3 ($p < 0.001$) and 400mg/kg *C. lutea* ($p < 0.01$) compared to the control. Conversely, in comparison to 100mg/kg AlCl_3 -exposed rats, there was significantly lower number of GFAP-expressing cells in the hippocampal CA3 of 10mg/kg donepezil- ($p < 0.05$) and 200mg/kg *C. lutea*-treated ($p < 0.001$) rats. Furthermore, there was an observable, albeit insignificant reduction in the number of hippocampal CA3 GFAP-positive cells in the 200mg/kg *C. lutea*-treated group compared to 10mg/kg donepezil. Similar observation was made between 400mg/kg *C. lutea*-treated group and 100mg/kg AlCl_3 -exposed rats (Figure 6)




Control




100mg/kg AlCl₃



10mg/kg Donepezil



200mg/kg *C. lutea*



400mg/kg *C. lutea*

Group of Figures 6: Immunohistochemical demonstration of GFAP in the hippocampal CA3 of rats exposed to experimental drugs. 400x magnification. Brown-stained cells are GFAP-expressing astrocytic cells.






Figure 7: Bar graphs depict the number of cells positive for GFAP in experimental rats. Each column represents mean \pm S.E.M. Data were analysed using one-way ANOVA followed by Tukey's post-test. *** $p < .001$, * $p < .05$ versus Control; ### $p < .001$, # $p < .05$ versus 100mg/kg AlCl₃.

3.2.3 GFAP – DG


Immunohistochemical analysis with one-way ANOVA demonstrated significant changes in the number of GFAP-positive cells [$F(4, 30) = 11.33, p < 0.0001$] in the hippocampal DG region following AlCl_3 exposure in rats. Post-hoc Tukey test revealed that rats subjected to 100mg/kg AlCl_3 ($p < 0.0001$) and 400mg/kg *C. lutea* ($p < 0.05$) had considerably more GFAP-positive cells in the hippocampus DG than in controls. In contrast to 100mg/kg AlCl_3 -exposed rats, 10mg/kg donepezil- ($p < 0.001$) and 200mg/kg *C. lutea*-treated ($p < 0.0001$) rats had a considerably lower number of GFAP-expressing cells in the hippocampal DG. Furthermore, there was an observable, albeit insignificant reduction in the number of GFAP-positive cells in the hippocampal DG of 200mg/kg *C. lutea*-treated group compared to 10mg/kg donepezil. Similar observation was made between 400mg/kg *C. lutea*-treated group and 100mg/kg AlCl_3 -exposed rats (Figure 8).




Control




100mg/kg AlCl₃



10mg/kg Donepezil



200mg/kg *C. lutea*



400mg/kg *C. lutea*

Group of Figures 8: Immunohistochemical demonstration of GFAP in the hippocampal DG of rats exposed to experimental drugs. 400x magnification. Brown-stained cells are GFAP-expressing astrocytic cells.




Figure 9: Bar graphs depict the number of cells positive for GFAP in experimental rats. Each column represents mean \pm S.E.M. Data were analysed using one-way ANOVA followed by Tukey's post-test. **** $p < .0001$, * $p < .05$ versus Control; ##### $p < .0001$, ### $p < .001$ versus 100mg/kg AlCl₃.




Figure 10: Bar graphs depict the number of cells positive for GFAP in experimental rats. Each column represents mean \pm S.E.M. Data were analysed using one-way ANOVA followed by Tukey's post-test. **** $p < .0001$, *** $p < .001$ versus Control; #### $p < .0001$, ### $p < .001$ versus 100mg/kg AlCl₃

4.DISCUSSION

Concerns continually exist globally on neurodegenerative diseases which sometimes could be traced to dangerous environmental contaminants like aluminium; its compounds and products (kawahara and kato-Negishi,2011). Aluminum is a well established neurotoxicant involved in the etiology of neurodegenerative diseases such as dialysis associated encephalopathy, Alzheimer's disease, Parkinson's disease, amyotrophic lateral sclerosis, multiple sclerosis and autism (kawahara and kato-Negishi, 2011 ; Arain et al., 2015; Jones et al., 2017; Mirza et al., 2017).

GFAP is a very sensitive marker for rapid astrocytic response to injury and disease (Zhang et al., 2014). In the present study, there was significant changes in the number of GFAP positive cells in the hippocampus CA₁ and CA₃ region and the dentate gyrus following AlCl₃ exposure in rats when compared to the control. There was an increased GFAP positive cells which is an indication of increased astrocytic reaction which occur as a result of neurotoxicity that may lead to neuroinflammation in the hippocampus. The hippocampus and dentate gyrus which are known for their role in learning and memory would invariably be impaired.

However, there was significantly reduced number of GFAP expressing cells in the hippocampal CA₁, CA₃ and dentate gyrus in the groups that received 10mg/kg donepezil and 200mg/kg *C. lutea*. Furthermore, it was observed that there was an insignificant reduction in the number of hippocampal CA₁, and CA₃ GFAP positive cells in the 200mg/kg *Carpolobia lutea* compared with the standard donepezil group. It may be suggested that neuroinflammation could be suppressed which may arise from neurotoxicity from aluminium chloride. The 400mg/kg of *C. lutea* group had an increased GFAP reactive cells with a corresponding decrease in the number

of hippocampal CA1 and CA3 GFAP positive cells though insignificant to the AlCl_3 . This implies that higher dose of *C. lutea* could trigger-astrocytic reactivity similar to AlCl_3 group.

It is noteworthy that the major functional phenotypes associated with astrocytes reactivity include neuroinflammation and impaired glutamate (Savory et al., 2003). Astrocytes reactivity is a hallmark of neuroinflammation that arises with Alzheimer disease (Mirza et al., 2017).

Glutamate is an excitatory neurotransmitter found throughout the central nervous system especially in the hippocampus. It maintains optimal levels within the extracellular space. As such it is important in memory and learning (Keshava et al., 2019).

Previous studies, revealed that exposure of co-cultured neurons and astrocytes to aluminium resulted in significant metal accumulation on both cells, whereas aluminium induced apoptosis was revealed only in astrocytes (suarez-fernandes et al., 1999). Jangra et al., (2015) indicated how oral administration of AlCl_3 (100mg/kg/b.w) for 42 days significantly elevated the levels of activity of cholinesterase and expression of amyloid precursor protein, amyloid beta in the hippocampus. Raj et al., (2021) asserted that aluminium chloride is a neurotoxin which generates neuropathologies and cognitive depletion resembling aging and Alzheimer disease. Also d-galactose + AlCl_3 led to neurotoxicity assessed on the basis of neuroinflammation, glial cell activation, neuronal damage in mice hippocampus (Zameer et al., 2021). Ezzat et al., 2022 indicated an upward glial fibrillary acidic protein (GFAP) immunoexpression in rat brain hippocampus. Chronic AlCl_3 administration raised the proinflammatory cytokines (interleukin-1 beta and tumor necrosis factor alpha level and increased acetylcholinesterase activity and reduced the BDNF content in the hippocampus (Kasbe et al., 2015). Zheng et al., (2019) suggested that exposure to AlCl_3 can cause neuroinflammation that may result in loss of spine in

the hippocampus and thereby leading to learning and memory deficits. Huo et al., (2023) posited that AlCl_3 induced mitochondrial damage, NLRP3 inflammasome activation and apoptosis. Another study demonstrated that Aluminium accelerates neuronal loss and astrogliosis in rain brain (Suryavanshi et al., (2022): Abdel et al.,(2022).

From the present study, it is evident that 200mg/kg of *C.lutea* reduced the GFAP immuno reactivity when compared to 10mg/kg of donepezil, the standard drug although statistically insignificant. It could be said that lower dose of *C.lutea* may have the tendencies of reversing impaired memory and learning and diseases associated with it. Our findings correlates with Noah et al., (2024). who suggested that *C. lutea* reverses the cardinal symptoms of ketamine-induced schizophrenia in a dose-dependent fashion by modulating the oxido-inflammatory and neurotransmitter-related mechanisms. Also our study is in consonance with Adeniran et al.,(2022) who suggested that *C. lutea* root extract improved steroidogenic activity in male wistar rats exposed to cadmium

Conclusion

In conclusion, 200mg/kg of *Carpolobia lutea* have the tendencies of protecting the neurons in the hippocampus from degenerating as a result of aluminium chloride induced neurotoxicity compared to the standard drug (10mg/kg donepezil).

Disclaimer

Author(s) hereby declare that NOgenerative AItechnologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators has been used during the writing or editing of this manuscript.

5.REFERENCES

1. Abdel-Aal RA, Hussein OA, Elsaady RG, Abdelzaher LA. Naproxen as a potential candidate for promoting rivastigmine anti-Alzheimer activity against aluminum chloride-prompted Alzheimer's-like disease in rats; neurogenesis and apoptosis modulation as a possible underlying mechanism. *Eur J Pharmacol.* 2022 Jan 15;915:174695. doi: 10.1016/j.ejphar.2021 PMID: 349
2. Akinola AO, Wahab OA, Raji Y. *Carpolobia lutea* Root Extract Improved Steroidogenic Activity in Male Wistar Rats Exposed to Cadmium. *Niger J Physiol Sci.* 2021 Dec 31;36(2):181-187. doi: 10.54548/njps.v36i2.6. PMID: 35947742.
3. Aljarari RM. Neuroprotective effects of a combination of *Boswellia papyrifera* and *Syzygium aromaticum* on AIC13 induced Alzheimer's disease in male albino rat. *Braz J Biol.* 2023 Oct 13;83:e272466. doi: 10.1590/1519-6984.272466. PMID: 37851769
4. Ajiwhen IO, Bisong SA. Effect of ethanolic extract of *Carpolobia lutea* G. Don (Polygalaceae) root on learning and memory in CD1 mice. *Niger J PhysiolSci.* 2013; 28:141–145.[PubMed] [Google Scholar]
5. AkhondzadehS, AbbasiSH. Herbal medicine in the treatment of Alzheimer's disease. *Am J Alzheimers Dis Other Demen,*2006;21:113–118.
6. Arain MS, Afridi HI, Kazi,TG, Talpur FN, Arain MB, Kazi A, Ali J. Correlation of aluminum and manganese concentrations in scalp hair samples of patients having neurological disorders. *Environ Monit Assess.*2015;187(2):10-15
7. Ballard C, Corbett A, Brayne C, Aarstand D, Jones E. Alzheimer's disease. *Lancet.*2001; 377:1019-1031.(Pubmed) (Google scholar).
8. Campbell A.(2002).”The potential role of aluminium in Alzheimer's disease”. *Nephrology Dialysis Transplantation.*2002; 17(2):17-20.

9. Carliss RD, Radovsky A, Chengelis, CP, O'Neill TP, Shuey DL. Oral administration of dextromethorphan does not produce neuronal vacuolation in the rat brain. *NeuroToxicol.* 2007; 28:813–818.

10. Domingo JL. (2006). Reproductive and developmental toxicity of aluminum: a review. *Neurotoxicology.* 2006;17(4), 515–521. [Google Scholar]

11. Duysen EG, Li B, Darvesh S, Lockridge O. Sensitivity of Butyrylcholinesterase knockout mice to (-)-huperzine A and donepezil suggest humans with butyrylcholinesterase deficiency may not tolerate Alzheimer's disease drugs and indicates butyrylcholinesterase function. *Toxicology.* 2007; 233:60-69.

12. El-Sebae, Y.D., Xu, W. U., Zhou, M. V. (2012). "Aluminum induces neurodegeneration and its toxicity arises from increased iron accumulation and reactive oxygen species (ROS) production," *Neurobiology of Aging.* 2012;3(1):199-205.

13. Essien EE, Ajibesin KK, Ekpo BA, Bala DN, Adesanya SA. Ethnobotanical survey of Southern States of Nigeria. *J Ethnopharmacol.* 2011. 115: 387-408.

14. Ezzat MI, Issa MY, Sallam IE, Zaafar D, Khalil HMA, Mousa MR, Sabry D, Gawish AY, Elghandour AH, Mohsen E. Impact of different processing methods on the phenolics and neuroprotective activity of *Fragaria ananassa* Duch. extracts in a D-galactose and aluminum chloride-induced rat model of aging. *Food Funct.* 2022 Jul 18;13(14):7794-7812. doi: 10.1039/d2fo00645f. PMID: 35766389

15. Flaten TP. Aluminium as a risk factor in Alzheimer's disease, with emphasis on drinking water. *Brain Science.* 2001; 55(2):187–196.

16. German HR. (2011). Histology of the central nervous system. *Toxicol Pathol* 39:22-35.

17. Huo S, Zhang X, Xu J, Zhang J, Du J, Li B, Song M, Shao B, Li Y, Xu F. Parkin-mediated mitophagy protects against aluminum trichloride-induced hippocampal apoptosis in mice via the mtROS-NLRP3 pathway. *Ecotoxicol Environ Saf.* 2023 Oct 1;264:115459. doi: 10.1016/j.ecoenv.2023.115459. Epub 2023 Sep 11. Erratum in: *Ecotoxicol Environ Saf.* 2024 Jan 1;269:115730. doi: 10.1016/j.ecoenv.2023.115730. PMID: 37703808

18. Ijomone OM, Olaibi OK, Obi AU, Alese MO. Immunohistochemistry staining protocols; Glial fibrillary acidic protein. *NeuroToxicol.* 2018;5(1):10-16.
19. Jangra A, Kasbe P, Pandey SN, Dwivedi S, Gurjar SS, Kwatra M, Mishra M, Venu AK, Sulakhiya K, Gogoi R, Sarma N, Bezbaruah BK, Lahkar M. Hesperidin and Silibinin Ameliorate Aluminum-Induced Neurotoxicity: Modulation of Antioxidants and Inflammatory Cytokines Level in Mice Hippocampus. *Biol Trace Elem Res.* 2015 Dec;168(2):462-71. doi: 10.1007/s12011-015-0375-7. PMID: 26018497
20. Jones K, Linhart C, Haekin C, Exley C. Urinary excretion of aluminium and silicon in secondary progressive multiple sclerosis. *EBioMedicine.* 2017; 26:60–67.
21. Kaur A, Joshi K, Minz RW, Gill KD. Neurofilament phosphorylation and disruption: a possible mechanism of chronic aluminium toxicity in Wistar rats. *Toxicology.* 2006; 219(3):1–10. [PubMed] [Google Scholar].
22. Kawahara M, Kato-Negishi M. (2011). Link between aluminum and the pathogenesis of Alzheimer's disease: The integration of the aluminum and amyloid cascade hypothesis. *Int J Alzheimer's Dis.* 2011;27:63-83.
23. Kasbe P, Jangra A, Lahkar M. Mangiferin ameliorates aluminium chloride-induced cognitive dysfunction via alleviation of hippocampal oxido-nitrosative stress, proinflammatory cytokines and acetylcholinesterase level. *J Trace Elem Med Biol.* 2015;31:107-12. doi: 10.1016/j.jtemb.2015.04.002. PMID: 26004900
24. Keshava R, Vazhayil V, Mitra R, Bhagavatula ID, Gope R. (2019). AlCl₃ causes Fas/Fas-L mediated cell death in the cortex and hippocampus of mouse brain. *Int. J. Biosci.* 2019; 12(3):21–35.
25. Lockridge O. Review of human butyrylcholinesterase structure, function, genetic variants, history of use in the clinic, and potential therapeutic uses. *Pharmacol Ther.* 2015; 148:34–46. [PubMed] [Google Scholar]
26. Manoharan I, Boopathy R, Darvesh S, Lockridge O. A medical health report on individuals with silent butyrylcholinesterase in the Vysya of India. *Clin Chim Acta.* 2007; 378:128–135. [PubMed] [Google Scholar]
27. Mirza A, King A, Troakes C, Exley C. Aluminium in brain tissue in familial Alzheimer's disease. *J. Trace Elem. Med. Biol.* 2017;40:30–36.

28. Mesrzzulam MM, Guillozet A, Shaw P, Levey A, Duysen EG, Lockridge O. Acetylcholinesterase knockouts establish central cholinergic pathway and can use butyrylcholinesterase to hydrolyze acetylcholine. *Neuroscience*. 2002; 110:627–639.
29. Odigie IP, Ettarh RR, Adigun SA. Chronic administration of extracts of *Carpolonia lutea* attenuates hypertension and reverses cardiac hypertrophy in 2K-1 C hypertensive rats. *Journal of Ethnopharmacology*. 2003; 86(12):181–185.
30. Omeiza NA, Bakre A, Ben-Azu B, Sowunmi AA, Abdulrahim HA, Chimezie J, Lawal SO, Adebayo OG, Alagbonsi AI, Akinola O, Abolaji AO, Aderibigbe AO. Mechanisms underpinning *Carpolonia lutea* G. Don ethanol extract's neurorestorative and antipsychotic-like activities in an NMDA receptor antagonist model of schizophrenia. *J Ethnopharmacol*. 2023 Jan 30;301:115767. doi: 10.1016/j.jep.2022.115767. PMID: 36206872.
31. Parihar MS, Hemnani T. Alzheimer's disease pathogenesis and therapeutic interventions. *J Clin Neurosci*. 2004;11:456–467.
32. Prema A, Justin TA, Manivasagam T, Mohamed EM, Guillemin GJ. Fenugreek seed powder attenuated aluminum chloride induced tau pathology, oxidative stress, and inflammation in a rat model of Alzheimer's disease. *J. Alzheimers*, 2017; 60(1):209–220.
33. Sun PI, Keen CL, Han B, Golub MS. Aluminum accumulation and neurotoxicity in Swiss-Webster mice after long-term dietary exposure to aluminum and citrate. *Journal of Neuroscience*. 2009;42:12961300.
34. Samaila, M., Chiroma, M., Aris, M., Moklas, C., Norma, M., Taib, M. D galactose and aluminium chloride induced rat model with cognitive impairments. *Biomedicine & Pharmacotherapy*. 2018;103:1602-1608.
35. Savory J, Herman MM, Ghribi O. Intracellular mechanisms underlying aluminum-induced apoptosis in rabbit brain. *J. Inorg. Biochem* 2003; 97(1),151– 154. [PubMed] [Google Scholar].
36. Suárez-Fernández MB, Soldado AB, Sanz-Medel A, Vega JA, Novelli A,

- Fernández-Sánchez MT, (1999). Aluminum-induced degeneration of astrocytes occurs via apoptosis and results in neuronal death. *Brain*. 1999; 23:10-15
37. Syad AN, Devi KP (2014). Botanicals: a potential source of new therapies for Alzheimer's disease. *Botanicals: Targets Therapy*. 2014; 4:11–26.
38. Tundis R, Bonesi M, Menichini F, Loizzo MR. Recent knowledge on medicinal plants as source of cholinesterase inhibitors for the treatment of dementia. *Mini Rev Med Chem*. 2016;16:605–618.
39. Suryavanshi J, Prakash C, Sharma D. Asiatic acid attenuates aluminium chloride-induced behavioral changes, neuronal loss and astrocyte activation in rats. *Metab Brain Dis*. 2022 Aug;37(6):1773-1785. doi: 10.1007/s11011-022-00998-3. PMID: 35554794.
40. Shin CY, Kim HS, Cha KH, Won DH, Lee JY, Jang SW, Sohn UD. The Effects of Donepezil, an Acetylcholinesterase Inhibitor, on Impaired Learning and Memory in Rodents. *Biomol Ther (Seoul)*. 2018 May 1;26(3):274-281. doi: 10.4062/biomolther.2017.189. PMID: 29463072; PMCID: PMC5933894
41. Raj K, Gupta GD, Singh S. Spermine protects aluminium chloride and iron-induced neurotoxicity in rat model of Alzheimer's disease via attenuation of tau phosphorylation, Amyloid- β (1-42) and NF- κ B pathway. *Inflammopharmacology*. 2021 Dec;29(6):1777-1793. doi: 10.1007/s10787-021-00883-y. Nov 2. PMID: 34727278.
42. Zameer S, Hussain S, Vohora D, Kalam Najmi A, Ali J, Akhtar M. Alendronate reduces the cognitive and neurological disturbances induced by combined doses of d-galactose and aluminum chloride in mice. *J Appl Toxicol*. 2021 Nov;41(11):1779-1793. doi: 10.1002/jat.4160. *E J Appl Toxicol*. 2023 Jun;43(6):951-954. doi: 10.1002/jat.4454. PMID: 33694194
43. Zhang Z, Rydel RE, Drzewiecki GJ, Fuson K, Wright S, Wogulis M, Audia JE, May PC, Hyslop PA. Amyloid β -mediated oxidative and metabolic stress in rat cortical neurons: No direct evidence for a role for H_2O_2 generation. *J. Neurochem*. 2014;67:1595–1606.
44. Zhang J, Huang W, Xu F, Cao Z, Jia F, Li Y. Iron Dyshomeostasis Participated in Rat Hippocampus Toxicity Caused by Aluminum Chloride. *Biol Trace Elem Res*. 2020 Oct;197(2):580-590. doi: 10.1007/s12011-019-02008-7 PMID: 31848921
14971