

Growth analysis, NPK uptake and Crude Protein Content in diversified Cropping System under Natural Farming

Commented [C1]: What did you do under natural farming? Be specific

ABSTRACT

A field experiment was conducted during *rabi Rabi* 2020-21 to *kharif* 2021 at Research Farm, Agronomy, CSKHPKV, Palampur (H.P.) to study growth analysis, NPK uptake and crude protein content in diversified cropping system under natural farming. The experiment was consisted of 9 treatments (C₁-Maize - wheat, C₂-Black gram - wheat + gram, C₃- Soybean - wheat + lentil, C₄-Cowpea - wheat + sarson, C₅-Okra - wheat + pea, C₆- Maize + black gram - gram, C₇-Maize + soybean - lentil, C₈-Maize + cowpea - sarson and C₉-Maize + okra - pea), replicated three time in Randomized Block Design. Legume-based systems resulted in significantly higher CGR and RGR among the cropping systems. During *rabi* 2020-21, the highest CGR was observed in the C₃ (soybean - wheat + lentil) of 5.53 g m⁻² day⁻¹, followed by C₁ (maize - wheat) at 5.29 g m⁻² day⁻¹. In *kharif* 2021, C₄ (cowpea - wheat + sarson) showed the highest CGR (10.22 g m⁻² day⁻¹). Higher RGR was found in C₇ system (maize + soybean - lentil) at 8.89 mg g⁻¹ day⁻¹, in *rabi* 2020-21. While, the system C₂ (black gram - wheat + gram) resulted in higher RGR at 13.76 mg g⁻¹ day⁻¹ before harvest in *kharif* 2021. Legumes enhances nutrient uptake by improving soil fertility and also resulted in improved the dry matter accumulation during mid-to-late crop stages under natural farming. A significant variation was observed higher nutrient uptake in both seed and by-products with legume-based systems under natural farming. Enhanced crude protein content was found in the legume-based systems, such as C₃ (maize + cowpea - sarson) with 75.50 % in *rabi* 2020-21.

Commented [C2]: What natural farming practises you have applied in whole experimental unit irrespective of different treatments. ? Mention here

Key words: Natural farming, cropping system, crop growth rate, relative growth rate, NPK uptake, crude protein content

Commented [C3]: Arrange all key words alphabetically

INTRODUCTION

In 1960's the Green Revolution started, which led to use of chemical pesticides, fertilizers, and high yielding varieties, resulted in increased agricultural productivity. Although it addressed food security, extensive long-term use of chemicals resulted in water pollution, reduction in biodiversity and soil degradation (Pingali 2012). Introduction of the natural farming in the recent decades which prioritize environmentally friendly, chemical-free techniques. The methods which are used in Natural Farming like Crop rotation, green manure, and animal integration to preserve soil fertility and manage pests. Also known as Subhash Palekar Natural Farming (SPNF), no use of artificial/synthetic chemicals, inputs the inputs used in natural farming results in improving soil health, by utilizing locally available resources like jeevamrit (a microbially rich biofertilizer) (Palekar 2016). In short, compared to conventional farming, the natural farming, reduces the environmental impact of agriculture, while enhancing soil organic matter and biodiversity. Natural farming is becoming more and more known as sustainable method of farming, as health concerns about chemical residues in food and the environment progress under conventional farming Devarinti (2016).

Commented [C4]: Include hypothesis, need of crop diversification. Also write the inclusion of legume's advantages for crop growth, soil health and sustainability.

MATERIALS AND METHODS

A field experiment was carried out during *rabi* 2020-21 to *kharif* 2021 at the research farm, Department of Agronomy, CSK HPKV, Palampur, Kangra, Himachal Pradesh.

Commented [C5]: You should mention meteorological data, soil pH, EC, organic carbon, NPK available in experimental plot. Also write varieties of all the crops diversified. Write seed rate, and plant geometry as well.

Methods of field preparation must be incorporated.

The experiment was carried out in Randomised Block Design with three replication and nine cropping system. The soil type that was silty-clay loam had low amounts of potassium and nitrogen, moderate levels of phosphorus, and an acidic content. The treatments comprised of C₁-Maize - wheat, C₂-Black gram - wheat + gram, C₃-Soybean - wheat + lentil, C₄-Cowpea - wheat + sarson, C₅-Okra - wheat + pea, C₆-Maize + black gram - gram, C₇-Maize + soybean - lentil, C₈-Maize + cowpea - sarson and C₉-Maize + okra – pea. In the *rabi* season the crops was sown in replacement series in intercropping system, while in *kharif* the additive series.

Crop Growth Rate (g m⁻² day⁻¹)

Crop growth rate (CGR) expresses the gain in dry matter production of the crop per unit land area per unit time and is expressed as gram per meter square per day (g m⁻² day⁻¹). It is calculated according to the formula given by Watson (1952).

$$\text{CGR} = \frac{1}{p} \times \frac{W_2 - W_1}{T_2 - T_1}$$

Where, W₂ and W₁ are dry weights at two sampling times T₂ and T₁ respectively.

Relative Growth Rate (mg g⁻¹ day⁻¹)

The relative growth rate (RGR) represents the rate of increase in dry weight per unit of plant dry weight and is expressed as mg m⁻² day⁻¹ (Blackman 1919).

$$\text{RGR} = \frac{\ln W_2 - \ln W_1}{t_2 - t_1}$$

NPK Uptake (kg ha⁻¹)

After the harvesting, the plant sample and seeds both were dried and converted to fine powder by using grinder total N, P and K were determined. Total N modified kjeldahl method (Jackson 1967); total phosphorus by using vanado-molybdate phosphoric acid yellow colour method (Jackson 1967) and total potassium diacid digestion method (Black 1965). Nutrient uptake is calculated by multiplying the nutrient concentration by the dry matter content and dividing by 100.

$$\text{Nutrient uptake} = \frac{\text{nutrient concentration (\%)} \times \text{dry matter content (kg ha}^{-1}\text{)}}{100}$$

Crude Protein

By multiplying the N content of grains by a factor of 6.25, the crude protein of grains was determined (Jones 1941).

Results and discussion

Crop Growth Rate (g m⁻² day⁻¹)

The variation in Crop Growth Rate (CGR) was observed in the cropping system after sowing to till harvest during *rabi* 2020-21 and *kharif* 2021 (Fig.1 and Fig. 2). At 30 DAS the CGR, in the cropping system ranging from 0.15 g m⁻² day⁻¹ in C₆ (maize + black gram - gram) to 2.46 g m⁻² day⁻¹ in C₈ (maize + cowpea - sarson) system in *rabi* 2020-21. At 120-150 DAS the CGR ranges between 0.39 g m⁻² day⁻¹ in C₆ (maize + black gram – gram) to 5.53 g m⁻² day⁻¹ in C₃ (soybean - wheat + lentil) system. The highest CGR was observed in C₃ (soybean - wheat + lentil) system with

Commented [C6]: Check whole section again. Support your findings with latest research , 2017 onwards

Commented [C7]: Where is yield data, and other growth parameters like NAR, LAI, LAD, SLA etc.

5.53 g m⁻² day⁻¹, followed by the C₁ (maize – wheat) system at 5.29 g m⁻² day⁻¹. Lowest CGR was observed in C₆ (maize + black gram - gram) system at 0.39 g m⁻² day⁻¹. Among intercrops at 120-150 DAS, the highest CGR was observed in C₃ (soybean - wheat + lentil) system with 1.77 g m⁻² day⁻¹, followed by the C₅ (okra - wheat + pea) at 1.70 g m⁻² day⁻¹. Lowest CGR was observed in C₂ (black gram - wheat + gram) system at 0.41 g m⁻² day⁻¹. In subsequent, *kharif* 2021, initially at 30 DAS, CGR among the cropping system ranging from 0.09 g m⁻² day⁻¹ in C₂ (black gram - wheat + gram) to 1.82 g m⁻² day⁻¹ in C₇ (maize + soybean - lentil) system. At 60-90 DAS the CGR ranges between 0.56 g m⁻² day⁻¹ in C₂ (black gram - wheat + gram) to 10.22 g m⁻² day⁻¹ in C₄ (cowpea - wheat + sarson). The highest CGR was observed in C₄ (cowpea - wheat + sarson) system with 10.22 g m⁻² day⁻¹, followed by the C₅ (okra - wheat + pea) at 3.57 g m⁻² day⁻¹. Lowest CGR was observed in C₂ (black gram - wheat + gram) system at 0.56 g m⁻² day⁻¹. Among intercrops at 60-90 DAS, the highest CGR was observed in C₈ (maize + cowpea - sarson) system with 10.18 g m⁻² day⁻¹, followed by the C₉ (maize + okra - pea) at 3.77 g m⁻² day⁻¹. The lowest CGR was observed in C₆ (maize + black gram - gram) at 0.49 g m⁻² day⁻¹. In the wheat - legume rotation, the CGR resulted in higher than in monocropping or continuous maize - wheat systems, during vegetative stage till harvest of crop growth (Choruma et al., 2022; Jensen et al., 2012). The legume-based rotations enhance CGR; also contribute carbon sequestration and enhanced nutrient cycling (Dwivedi et al., 2015; Kirkegaard and Ryan, 2014).




Fig.1 Effect of cropping system on crop growth rate (g m⁻² day⁻¹) under natural farming during *rabi* 2020-21




Fig.2 Effect of cropping system on crop growth rate ($\text{g m}^{-2} \text{day}^{-1}$) under natural farming during *kharif* 2020-21

Relative Growth Rate ($\text{mg g}^{-1} \text{day}^{-1}$)

The Relative Growth Rate (RGR) was observed in the cropping system after sowing till harvest during *rabi* 2020-21 and *kharif* 2021 (Fig. 3 and Fig. 4). Initially at 30 DAS the RGR ranges from $9.06 \text{ mg g}^{-1} \text{day}^{-1}$ in C₉ (maize + okra - pea) to $18.02 \text{ mg g}^{-1} \text{day}^{-1}$ in C₇ (maize + soybean - lentil) system in *rabi* 2020-21. At 120-150 DAS the RGR ranges between $0.50 \text{ mg g}^{-1} \text{day}^{-1}$ in C₈ (maize + cowpea - sarson) to $8.89 \text{ mg g}^{-1} \text{day}^{-1}$ in C₇ (maize + soybean - lentil) system. The highest RGR was observed in C₇ (maize + soybean - lentil) system with $8.89 \text{ mg g}^{-1} \text{day}^{-1}$, followed by the C₃ (soybean - wheat + lentil) system at $6.80 \text{ mg g}^{-1} \text{day}^{-1}$. Lowest RGR was observed in C₈ (maize + cowpea - sarson) system at $0.50 \text{ mg g}^{-1} \text{day}^{-1}$. Among intercrops at 120-150 DAS, the highest RGR was observed in C₃ (soybean - wheat + lentil) system with $4.26 \text{ mg g}^{-1} \text{day}^{-1}$, followed by the C₂ (black gram - wheat + gram) at $3.85 \text{ mg g}^{-1} \text{day}^{-1}$. Lowest RGR was observed in C₄ (cowpea - wheat + sarson) system at $1.69 \text{ mg g}^{-1} \text{day}^{-1}$. In subsequent, *kharif* 2021, initially at 30 DAS, RGR among the cropping system ranging from $9.71 \text{ mg g}^{-1} \text{day}^{-1}$ in C₃ (soybean - wheat + lentil) to $45.83 \text{ mg g}^{-1} \text{day}^{-1}$ in C₄ (cowpea - wheat + sarson) system. At 60-90 DAS the RGR ranges between $2.19 \text{ mg g}^{-1} \text{day}^{-1}$ in C₇ (maize + soybean - lentil) to $13.76 \text{ mg g}^{-1} \text{day}^{-1}$ in C₂ (black gram - wheat + gram). The highest RGR was observed in C₂ (black gram - wheat + gram) system with $13.76 \text{ mg g}^{-1} \text{day}^{-1}$, followed by the C₄ (cowpea - wheat + sarson) at $11.72 \text{ mg g}^{-1} \text{day}^{-1}$. Lowest RGR was observed in C₇ (maize + soybean - lentil) system at $2.19 \text{ mg g}^{-1} \text{day}^{-1}$. Among intercrops at 60-90 DAS, the highest RGR was observed C₆ (maize + black gram - gram) system with $16.61 \text{ mg g}^{-1} \text{day}^{-1}$, followed by the C₈ (maize + cowpea - sarson) at $13.84 \text{ mg g}^{-1} \text{day}^{-1}$. Lowest RGR was observed in C₉ (maize + okra - pea) system at $6.46 \text{ mg g}^{-1} \text{day}^{-1}$. Dwivedi et al. (2015) reported that the maize-legume intercropping systems led to significant improvements in RGR; also, systems increased overall biomass production due to enhanced nutrient availability from legumes. The inclusion of legumes resulted in

improved nutrient status also enhanced growth rates (Jensen et al. 2012). Crop rotation with legumes enhances growth rates and yield in multiple cropping system (Kirkegaard and Ryan 2014). In wheat and maize system, addition of legumes resulted in higher RGR by improving soil health and nutrient availability (Choruma et al. 2022).




Fig.3 Effect of cropping system on relative growth rate ($\text{mg g}^{-1} \text{ day}^{-1}$) under natural farming during *rabi* 2020-21




Fig.4 Effect of cropping system on relative growth rate ($\text{mg g}^{-1} \text{day}^{-1}$) under natural farming during *kharif* 2021

NPK Uptake (kg ha^{-1})

The data for NPK uptake in the seed and by-product for *rabi* 2020-21 and *kharif* 2021 presented in the Fig. 5, Fig.6 and Fig.7. A variation for the NPK uptake in seed and by-product was observed during both the crop season under natural farming.

N Uptake (seed and by-product)

In the *rabi* 2020-21, the highest seed N uptake was observed in the C₉ (maize + okra - pea) system at 67.52 kg ha^{-1} , followed by C₇ (maize + soybean - lentil) at 64.92 kg ha^{-1} . The Lowest seed N uptake was observed in the C₁ (maize - wheat) system with 30.96 kg ha^{-1} . In subsequent *kharif* 2021, the highest seed N uptake was observed in the C₅ (okra - wheat + pea) system at 56.84 kg ha^{-1} , followed by C₉ (maize + okra - pea) at 53.00 kg ha^{-1} . The lowest seed N uptake was observed in the C₂ (black gram - wheat + gram) system with 24.26 kg ha^{-1} . Overall system's, the highest seed N uptake was observed in the C₉ (maize + okra - pea) system at $120.52 \text{ kg ha}^{-1}$, followed by C₅ (okra - wheat + pea) at $107.69 \text{ kg ha}^{-1}$. The Lowest seed N uptake was observed in the C₂ (black gram - wheat + gram) system with 57.66 kg ha^{-1} . In the *rabi* 2020-21, the highest by-product N uptake was observed in the C₇ (maize + soybean - lentil) system at 18.43 kg ha^{-1} , followed by C₈ (maize + cowpea - sarson) at 16.14 kg ha^{-1} . The Lowest by-product N uptake was observed in the C₉ (maize + okra - pea) system with 6.66 kg ha^{-1} . In subsequent *kharif* 2021, the highest by-product N uptake was observed in the C₂ (black gram - wheat + gram) system at 34.10 kg ha^{-1} , followed by C₈ (maize + cowpea - sarson) at 28.33 kg ha^{-1} . The lowest by-product N uptake was observed in the C₅ (okra - wheat + pea) system with 3.50 kg ha^{-1} . Overall system's, the highest by-product N uptake was observed in the C₄ (cowpea - wheat + sarson) system at 50.43 kg ha^{-1} , followed by C₂ (black gram - wheat + gram) at 46.85 kg ha^{-1} . The Lowest by-product N uptake was observed in the C₅ (okra - wheat + pea) system with 11.55 kg ha^{-1} .




Fig.5 (a) N uptake in seed during *rabi* 2020-21, (b) N uptake in seed during *kharif* 2021, (c) N uptake in straw during *rabi* 2020-21 and (d) N uptake in straw during *kharif* 2021

Adu-Gyamfi et al. (2007) observed higher nitrogen uptake under soybean-maize intercropping systems compared to monocropping system. Rao and Mathuva (2000) found significantly higher nitrogen accumulation in the cowpea and subsequent cereal crop under cowpea - maize intercropping system. In lentil-based intercropping with cereals like wheat and barley resulted in improved N uptake due to biological nitrogen fixation by lentils and cowpea (Stagnari et al. 2017; Sharma T et al. 2023). Rotation with legumes such as soybeans under wheat-based system resulted in enhanced nitrogen content in both the seed and straw of subsequent cereal crops due to residual nitrogen from the legumes (Peoples et al.1995)




Fig.6 (a) P uptake in seed during *rabi* 2020-21, (b) P uptake in seed during *kharif* 2021, (c) P uptake in straw during *rabi* 2020-21 and (d) P uptake in straw during *kharif* 2021

P Uptake (seed and by-product)

In the *rabi* 2020-21, the highest seed P uptake was observed in the system C₉ (maize + okra - pea) at 41.29 kg ha⁻¹, followed by C₅ (okra - wheat + pea) at 22.54 kg ha⁻¹. The Lowest seed P uptake was observed in the system C₇ (maize + soybean - lentil) with 3.74 kg ha⁻¹. In subsequent *kharif* 2021, the highest seed P uptake was observed in the C₅ (okra - wheat + pea) system at 14.30 kg ha⁻¹, followed by C₉ (maize + okra - pea) at 11.54 kg ha⁻¹. The lowest seed P uptake was observed in the C₂ (black gram - wheat + gram) system with 1.94 kg ha⁻¹. Overall system's, the highest seed P uptake was observed in the C₉ (maize + okra - pea) system at 52.83 kg ha⁻¹, followed by C₅ (okra - wheat + pea) at 36.84 kg ha⁻¹. The Lowest seed P uptake was observed in the C₂ (black gram - wheat + gram) system with 7.06 kg ha⁻¹. In the *rabi* 2020-21,

The highest by-product P uptake was observed in the system C₉ (maize + okra - pea) at 10.97 kg ha⁻¹, followed by C₈ (maize + cowpea - sarson) at 8.51 kg ha⁻¹. The Lowest by-product P uptake was observed in the system C₆ (maize + black gram - gram) with 2.27 kg ha⁻¹. In subsequent *kharif* 2021, the highest by-product P uptake was observed in the C₂ (black gram - wheat + gram) system at 6.05 kg ha⁻¹, followed by C₈ (maize + cowpea - sarson) at 5.85 kg ha⁻¹. The lowest by-product P uptake was observed in the C₅ (okra - wheat + pea) system with 0.81 kg ha⁻¹. Overall system's, the highest by-product P uptake was observed in the C₄ (cowpea - wheat + sarson) system at 14.49 kg ha⁻¹, followed by C₈ (maize + cowpea - sarson) at 14.36 kg ha⁻¹. The Lowest by-product P uptake was observed in the C₆ (maize + black gram - gram) system with 4.43 kg ha⁻¹. Singh et al. (2015) reported that in intercropping with legumes resulted in enhanced phosphorus uptake compared to monocropping. (Naseem et al. 2020) reported that significantly enhanced phosphorus use efficiency, resulting in higher seed and biomass phosphorus uptake, when maize intercropped legumes. The leguminous crops resulted in crucial role in the phosphorus cycling and uptake due to their symbiotic relationships with mycorrhizal fungi (Dutta et al. 2016).

K Uptake (seed and by-product)

In the *rabi* 2020-21, the highest seed K uptake was observed in the C₉ (maize + okra - pea) system at 45.71 kg ha⁻¹, followed by C₅ (okra - wheat + pea) at 3.32 kg ha⁻¹. The Lowest seed K uptake was observed in the C₇ (maize + soybean - lentil) system with 3.34 kg ha⁻¹. In subsequent *kharif* 2021, the highest seed K uptake was observed in the C₅ (okra - wheat + pea) system at 24.52 kg ha⁻¹, followed by C₉ (maize + okra - pea) at 16.63 kg ha⁻¹. The lowest seed K uptake was observed in the C₂ (black gram - wheat + gram) system with 4.68 kg ha⁻¹. Overall system's, the highest seed K uptake was observed in the C₉ (maize + okra - pea) system at 31.21 kg ha⁻¹, followed by C₅ (okra - wheat + pea) at 24.84 kg ha⁻¹. The Lowest seed K uptake was observed in the C₂ (black gram - wheat + gram) system with 5.84 kg ha⁻¹. In the *rabi* 2020-21, the highest by-product K uptake was observed in the C₁ (maize - wheat) system at 28.77 kg ha⁻¹, followed by C₂ (black gram - wheat + gram) at 28.16 kg ha⁻¹. The Lowest by-product K uptake was observed in the C₉ (maize + okra - pea) system with 7.52 kg ha⁻¹.




Fig.7 (a) K uptake in seed during *rabi* 2020-21, (b) (f) K uptake in seed during *kharif* 2021, (c) K uptake in straw during *rabi* 2020-21 and (d) K uptake in straw during *kharif* 2021

In subsequent *kharif* 2021, the highest by-product K uptake was observed in the C₈ (maize + cowpea - sarson) system at 73.63 kg ha⁻¹, followed by C₇ (maize + soybean - lentil) at 55.49 kg ha⁻¹. The lowest by-product K uptake was observed in the C₅ (okra - wheat + pea) system with 4.86 kg ha⁻¹. Overall system's, the highest by-product K uptake was observed in the C₈ (maize + cowpea - sarson) system at 85.38 kg ha⁻¹, followed by C₁ (maize - wheat) at 72.52 kg ha⁻¹. The Lowest by-product K uptake was observed in the C₅ (okra - wheat + pea) system with 25.03 kg ha⁻¹. The combination of legumes and cereals resulted in enhanced K uptake compared to sole cropping, overall increased yields and nutrient content (Ghosh and Hossain 2019). Overall improved nutrient profile in intercropping legumes with cereals, including potassium uptake, particularly in systems like maize-legume intercropping (Adu-Gyamfiet al. 2007).

Crude Protein

Crude protein content for the cropping system in *rabi* 2020-21 and *kharif* 2021, presented in the Table 1. In *rabi* 2020-21, the highest crude protein was found in the C₈ (maize + cowpea - sarson) system at 75.50 %, followed by C₆ (maize + black gram - gram) at 71.48 %. The lowest crude protein was found in the C₄ (cowpea - wheat + sarson) with 8.58 %. The crude protein among the intercrops was found in the order C₄ (cowpea - wheat + sarson) > C₂ (black gram - wheat + gram) > C₃ (soybean - wheat + lentil) > C₅ (okra - wheat + pea). In *rabi* 2020-21, the highest crude protein was found in C₂ (black gram - wheat + gram) system at 22.63 %, followed by C₃ (soybean - wheat + lentil) at 18.06 %. The lowest crude protein was found in the C₁ (maize - wheat) with 7.46 %. The crude protein among the intercrops was found in the order C₆ (maize + black gram - gram) > C₇ (maize + soybean - lentil) > C₈ (maize + cowpea - sarson) > C₉ (maize + okra - pea). The legumes significantly enhanced the crude protein content of the main crops due to their

nitrogen-fixing ability (Ghosh et al. 2019). Ali et al. (2017) observed that intercropping legumes with cereals improved protein content in both crops compared to sole. The addition of legumes as intercrop (*viz.* black gram and cowpea) resulted in increased crude protein content in main crops (Sahni et al. 2021). Legume as an intercrop improves main crops nutritional quality especially the crude protein content (Choudhary et al. 2020).

Table 1. Effect of cropping systems on crude protein in grains during crop season 2020-21 under natural farming

Cropping system	Crude Protein			
	Rabi 2020-21	Intercrop	Kharif 2021	Intercrop
C ₁ Maize - wheat	8.69	-	7.46	-
C ₂ Black gram - wheat + gram	8.75	66.81	22.63	-
C ₃ Soybean - wheat + lentil	8.75	50.75	18.06	-
C ₄ Cowpea - wheat + sarson	8.58	75.06	13.81	-
C ₅ Okra - wheat + pea	8.88	10.06	6.96	-
C ₆ Maize + black gram - gram	71.48	-	7.73	20.13
C ₇ Maize + soybean - lentil	52.13	-	7.69	16.00
C ₈ Maize + cowpea - sarson	78.50	-	7.79	13.13
C ₉ Maize + okra - pea	10.33	-	8.31	5.81

Commented [C8]: Where is CD value?

Conclusion

The study evaluated Crop Growth Rate (CGR), Relative Growth Rate (RGR), NPK uptake and crude protein content across various cropping systems during *rabi* 2020-21 and *kharif* 2021. It was found that the significant variations in CGR and RGR among systems, with legume-based systems (e.g., C₃: soybean - wheat + lentil and C₈: maize + cowpea - sarson). At 120-150 DAS, the highest CGR (5.53 g m⁻² day⁻¹) was observed in the C₃ (soybean - wheat + lentil) system, while C₄ (cowpea - wheat + sarson) and C₈ systems showed superior performance in subsequent *kharif* 2021. Similarly, the inclusion legume in cropping system resulted in higher RGR, with C₇ (maize + soybean - lentil) and C₃ (soybean - wheat + lentil) system maximum rates at later stages. Enhanced NPK uptake in seeds and by-products was found significantly higher in legume-based cropping system compared to C₁ (maize - wheat) system under natural farming. The crude protein content was also found to be higher in legume-based systems (*viz.* C₈ and C₂) under natural farming. In addition, legume-based systems not only improve soil fertility, but also resulted in increase in dry matter accumulation, enhanced nutrient uptake and crude protein content among the cropping system under natural farming.

Commented [C9]: Conclusion must be short, explaining findings and recommendation to the farmer

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

References

- Adu-Gyamfi JJ, Myaka FA, Sakala WD, Odgaard R, Vesterager JM and Høgh-Jensen H. Biological nitrogen fixation and nitrogen transfer in intercropping systems. *Plant and Soil*, 2007;296(1-2), 31-41.
- Ali MH and Rahman MZ. Nutritional benefits of intercropping legumes with cereals. *International Journal of Plant Production*, 2017;11(3), 309-318.
- Black CA. *Methods of soil analysis. Part II. Chemical and mineralogical properties.* American Society of Agronomy, Madison, Wisconsin, USA, 1965.
- Blackman VH. The compound interest law and plant growth. *Annals of botany*, 191933(131) 353-360.
- Choruma DJ, Senzanje A, Chagonda I and Mpepereki S. Legumes improve wheat and maize productivity. *Agronomy for Sustainable Development*. 2022;42(1), 6.
- Choudhary V, Tiwari S and Dey S. Nutritional advantages of intercropping legumes with cereals. *Journal of Agricultural Science and Technology*. 2020;22(5), 1234-1245.
- Devarinti SR. Natural farming: eco-friendly and sustainable. *Agrotechnology*. 2016;5(2), p.1000147.
- Dutta S, Bhowmik A and Mondal MK. Role of legumes in phosphorus acquisition in intercropping systems. *Agricultural Sciences*. 2016;7(7), 989-999.
- Dwivedi A, Reddy S and Bansal A. Potential role of maize-legume intercropping in agro-ecosystem sustainability. *International Journal of Life Sciences Biotechnology and Pharma Research*. 2015;4(3), 154-158.
- Ghosh P, Choudhury S and Dutta D. Effects of intercropping on the protein content of food crops. *Agronomy for Sustainable Development*. 2019;39(3), 32.
- Ghosh S and Hossain MI. The effect of intercropping on the nutrient uptake and yield of crops in the cereal-legume systems. *Journal of Plant Nutrition*, 2019;42(5), 545-555.
- Jackson M. *Soil chemical analysis prentice.* Hall of India Private Limited, New Delhi. 1967;498(1).
- Jensen ES, Peoples MB, Boddey RM, Gresshoff PM, Hauggaard-Nielsen H and Alves BJR. Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries: A review. *Agronomy for Sustainable Development*. 2012;32(2), 329–364.
- Jones DB. Factors for converting percentages of nitrogen in foods and feeds into percentages of protein. *United States Department of Agriculture Circular*, No. 183. Washington, DC: USDA, 1941.
- Kirkegaard JA and Ryan MH. Magnitude and mechanisms of persistent crop sequence effects on wheat. *Field Crops Research*. 2014;164, 154-165.
- Naseem M, Shah AA and Raza SH. Phosphorus efficiency in intercropped legumes and cereals. *Nutrient Cycling in Agroecosystems*. 2020;118(2), 191-203.

Commented [C10]: Check all the references, author name, year, issue, volume, name of journal, publisher, title of manuscript and page number. You can also write DOI in each references

Palekar S. Zero Budget Natural Farming. Subhash Palekar Natural Farming. Pingali P 2012."Green Revolution: Impacts, limits, and the path ahead." Proceedings of the National Academy of Sciences. 2016;109(31),12302–12308.

Peoples MB, Ladha JK and Herridge DF. Enhancing legume N₂ fixation through selection and breeding. *Agronomy Journal*. 1995;87(1), 1-8.

Rao MR and Mathuva M. Legumes for improving maize yields and nitrogen balance in a humid tropical environment. *Field Crops Research*. 2000;68(3), 181-190.

Sahni N, Singh G and Kumar V. Intercropping of legumes with cereals enhances protein content: A review. *Journal of Crop Science and Technology*. 2021;25(2), 117-129.

Singh P, Kumari M, Yadav H, and Singh R. Phosphorus dynamics in intercropped systems: A meta-analysis. *Plant and Soil*. 2015;392(1-2), 117-134.

Stagnari F, Ramazzotti S and Pisante M. Conservation agriculture in Europe: An approach to sustainable crop production by enhancing biodiversity and soil fertility. *Agronomy*. 2017;7(7), 63.

Sharma T, Singh J, Singh A, Sharma R, Chauhan G. Effect of organic nutrient sources on the yield, nutrient uptake and nodulation in Cowpea (*Vigna unguiculata*) under mid-hill conditions of Western Himalayas. *Environment Conservation Journal*. 2023 7;24(2):250-6.

Watson DJ. The physiological basis of variation in yield. *Advances in agronomy*. 1952;4,101-145.