

## Growth, NPK uptake and Crude Protein Content in diversified Cropping System under Natural Farming

### ABSTRACT

A field experiment was conducted during *Rabi* 2020-21 to *Kharif* 2021 at Research Farm, Agronomy, CSKHPKV, Palampur (H.P.) to study growth analysis, NPK uptake and crude protein content in diversified cropping system under natural farming. The experiment was consisted of 9 treatments (C<sub>1</sub>-Maize - wheat, C<sub>2</sub>-Black gram - wheat + gram, C<sub>3</sub>-Soybean - wheat + lentil, C<sub>4</sub>-Cowpea - wheat + sarson, C<sub>5</sub>-Okra - wheat + pea, C<sub>6</sub>-Maize + black gram - gram, C<sub>7</sub>-Maize + soybean - lentil, C<sub>8</sub>-Maize + cowpea - sarson and C<sub>9</sub>-Maize + okra - pea), replicated three time in Randomized Block Design. The natural farming inputs includes *beejamrit* (used for seed treatment @ 1 litre/10 kg seed), *jeevamrit* (5 drenching application at 21 days interval; total used 450 litres), *ghanjeevamrit* (soil applied during sowing @ 500 kg ha<sup>-1</sup>) and mulching (applied after germination @ 10000 kg ha<sup>-1</sup>). Legume-based systems resulted in significantly higher CGR and RGR among the cropping systems. During *Rabi* 2020-21, the highest CGR was observed in the C<sub>3</sub> (soybean - wheat + lentil) of 5.53 g m<sup>-2</sup> day<sup>-1</sup>, followed by C<sub>1</sub> (maize - wheat) at 5.29 g m<sup>-2</sup> day<sup>-1</sup>. In *Kharif* 2021, C<sub>4</sub> (cowpea - wheat + sarson) showed the highest CGR (10.22 g m<sup>-2</sup> day<sup>-1</sup>). Higher RGR was found in C<sub>7</sub> system (maize + soybean - lentil) at 8.89 mg g<sup>-1</sup> day<sup>-1</sup>, in *Rabi* 2020-21. While, the system C<sub>2</sub> (black gram - wheat + gram) resulted in higher RGR at 13.76 mg g<sup>-1</sup> day<sup>-1</sup> before harvest in *Kharif* 2021. Legumes enhances nutrient uptake by improving soil fertility and also resulted in improved the dry matter accumulation during mid-to-late crop stages under natural farming. A significant variation was observed higher nutrient uptake in both seed and by-products with legume-based systems under natural farming. Enhanced crude protein content was found in the legume-based systems, such as C<sub>8</sub> (maize + cowpea - sarson) with 75.50 % in *Rabi* 2020-21.

**Key words:** Crop Growth Rate, Cropping System, Crude Protein Content, Natural Farming, NPK Uptake, Relative Growth Rate

### INTRODUCTION

In 1960's the Green Revolution started, which led to use of chemical pesticides, fertilizers, and high yielding varieties, resulted in increased agricultural productivity. Although it addressed food security, extensive long-term use of chemicals resulted in water pollution, reduction in biodiversity and soil degradation (Pingali 2012). For sustainable agriculture, it is important to conserve natural resources by reducing degradation and dependency on synthetic chemicals used as fertilizers, pest control, and weed management. The inclusion of legumes, vegetables, and other crops in the existing non-leguminous systems enhances overall production and profitability. Also, inclusion of legumes in the cropping system as intercrop/sole crop resulted in soil health by enhancing soil fertility, increasing organic matter and nutrient cycling, results in sustainable and productive agricultural systems (Jena et al. 2022). Introduction of the natural farming in the recent decades which prioritize environmentally friendly, chemical-free techniques. The methods which are used in Natural Farming like Crop rotation, green manure, and animal integration to preserve soil fertility and manage pests. Also known as Subhash Palekar Natural Farming

(SPNF), no use of artificial/synthetic chemicals, inputs the inputs used in natural farming results in improving soil health, by utilizing locally available resources like jeevamrit (a microbially rich biofertilizer) (Palekar 2016). In short, compared to conventional farming, the natural farming, reduces the environmental impact of agriculture, while enhancing soil organic matter and biodiversity. Natural farming principles advocate that the growing of two or more than two crops in a year result in reduced pest attacks, optimizes land use and maintain ecological balance, without affecting the soil resources. Natural farming is becoming more and more known as sustainable method of farming, as health concerns about chemical residues in food and the environment progress under conventional farming Devarinti (2016).

## MATERIALS AND METHODS

A field experiment was carried out during *Rabi* 2020-21 to *Kharif* 2021 at the research farm, Department of Agronomy, CSK HPKV, Palampur, Kangra, Himachal Pradesh. The experiment was carried out in Randomised Block Design with three replication and nine cropping system. The soil type that was silty-clay loam had low amounts of potassium and nitrogen, moderate levels of phosphorus, and an acidic content. The treatments comprised of C<sub>1</sub> -Maize - wheat, C<sub>2</sub>-Black gram - wheat + gram, C<sub>3</sub>-Soybean - wheat + lentil, C<sub>4</sub>-Cowpea - wheat + sarson, C<sub>5</sub>-Okra - wheat + pea, C<sub>6</sub>-Maize + black gram - gram, C<sub>7</sub>-Maize + soybean - lentil, C<sub>8</sub>-Maize + cowpea - sarson and C<sub>9</sub>-Maize + okra – pea. In the *Rabi* season the crops was sown in replacement series in intercropping system, while in *Kharif* the additive series. The initial status of the before the start of the experiment presented in the table 1. The natural farming inputs includes *beejamrit* (used for seed treatment @ 1 litre/10 kg seed), *jeevamrit* (5 drenching application at 21 days interval; total used 450 litres), *ghanjeevamrit* (soil applied during sowing @ 500 kg ha<sup>-1</sup>) and mulching (applied after germination @ 10000 kg ha<sup>-1</sup>). The details of the varieties used, seed rates and row spacing presented in the table 2.

**Table 1. Initial soil physical, chemical and biological parameters of the experimental site**

Particulars	Content in soil	Analytical method employed
<b>A. Physical properties</b>		
i. Sand (%)	20.5	International pipette method
ii. Silt (%)	43.6	(Piper 1966)
iii. Clay (%)	35.7	
iv. Texture	Silty clay loam	
v. Bulk density (g cm <sup>-3</sup> )	1.33	Core sampler method (Blake 1965)
<b>B. Chemical properties</b>		
i. pH	5.01	Glass electrode pH meter (Jackson 1973)
ii. Electrical Conductivity (EC) (µS m <sup>-1</sup> )	110.01	Suspension with EC meter (Jackson 1973)
iii. Organic carbon (%)	0.72	Wet digestion method (Walkley and Black 1934)
iv. Available nutrient (kg ha <sup>-1</sup> )		

a. Available nitrogen	258.45	Alkaline permanganate method (Subbiah and Asija 1956)
b. Available phosphorus	24.89	Ammonium molybdate blue colour method (Olsen et al. 1954)
c. Available potassium	167.46	Ammonium acetate extraction method (AOAC 1970)

**Table 2. Details of the varieties used, seed rates and row spacing**

Crop	Variety	Seed rate (kg ha <sup>-1</sup> )	Spacing (cm)
<b>Rabi</b>			
Wheat	HPW-368	100	22.5
Gram	Him-Chana 1	45	25
Lentil	HPLO-1	30	25
Pea	PB-89	75	45 x 10
Sarson	Sheetal	6	30
<b>Kharif</b>			
Maize	Girija	20	60 x 20
Soybean	HIMCO-1685	75	45 x 10
Black Gram	Him Mash 1	20	30
Cowpea	CO-519	20	45
Okra	P8	20	45 x 15

### Crop Growth Rate (g m<sup>-2</sup> day<sup>-1</sup>)

Crop growth rate (CGR) expresses the gain in dry matter production of the crop per unit land area per unit time and is expressed as gram per meter square per day (g m<sup>-2</sup> day<sup>-1</sup>). It is calculated according to the formula given by Watson (1952).

$$\text{CGR} = \frac{1}{P} \times \frac{W_2 - W_1}{T_2 - T_1}$$

Where,  $W_2$  and  $W_1$  are dry weights at two sampling times  $T_2$  and  $T_1$  respectively.

### Relative Growth Rate (mg g<sup>-1</sup> day<sup>-1</sup>)

The relative growth rate (RGR) represents the rate of increase in dry weight per unit of plant dry weight and is expressed as mg m<sup>-2</sup> day<sup>-1</sup> (Blackman 1919).

$$\text{RGR} = \frac{\ln W_2 - \ln W_1}{t_2 - t_1}$$

### NPK Uptake (kg ha<sup>-1</sup>)

After the harvesting, the plant sample and seeds both were dried and converted to fine powder by using grinder total N, P and K were determined. Total N modified kjeldahl method (Jackson 1967); total phosphorus by using vanado-molybdate phosphoric acid yellow colour method (Jackson 1967) and total potassium diacid digestion method (Black 1965). Nutrient uptake is calculated by multiplying the nutrient concentration by the dry matter content and dividing by 100.

$$\text{Nutrient uptake (Kg ha}^{-1}\text{)} = \frac{\text{nutrient concentration (\%)} \times \text{dry matter content (kg ha}^{-1}\text{)}}{100}$$

## Crude Protein

By multiplying the N content of grains by a factor of 6.25, the crude protein of grains was determined (Jones 1941).

## Results and discussion

### Crop Growth Rate ( $\text{g m}^{-2} \text{ day}^{-1}$ )

The variation in Crop Growth Rate (CGR) was observed in the cropping system after sowing to till harvest during **Rabi** 2020-21 and **Kharif** 2021 (Fig.1 and Fig. 2). At 30 DAS the CGR, in the cropping system ranging from  $0.15 \text{ g m}^{-2} \text{ day}^{-1}$  in C<sub>6</sub> (maize + black gram - gram) to  $2.46 \text{ g m}^{-2} \text{ day}^{-1}$  in C<sub>8</sub> (maize + cowpea - sarson) system in **Rabi** 2020-21. At 120-150 DAS the CGR ranges between  $0.39 \text{ g m}^{-2} \text{ day}^{-1}$  in C<sub>6</sub> (maize + black gram - gram) to  $5.53 \text{ g m}^{-2} \text{ day}^{-1}$  in C<sub>3</sub> (soybean - wheat + lentil) system. The highest CGR was observed in C<sub>3</sub> (soybean - wheat + lentil) system with  $5.53 \text{ g m}^{-2} \text{ day}^{-1}$ , followed by the C<sub>1</sub> (maize - wheat) system at  $5.29 \text{ g m}^{-2} \text{ day}^{-1}$ . Lowest CGR was observed in C<sub>6</sub> (maize + black gram - gram) system at  $0.39 \text{ g m}^{-2} \text{ day}^{-1}$ . Among intercrops at 120-150 DAS, the highest CGR was observed in C<sub>3</sub> (soybean - wheat + lentil) system with  $1.77 \text{ g m}^{-2} \text{ day}^{-1}$ , followed by the C<sub>5</sub> (okra - wheat + pea) at  $1.70 \text{ g m}^{-2} \text{ day}^{-1}$ . Lowest CGR was observed in C<sub>2</sub> (black gram - wheat + gram) system at  $0.41 \text{ g m}^{-2} \text{ day}^{-1}$ . In subsequent, **Kharif** 2021, initially at 30 DAS, CGR among the cropping system ranging from  $0.09 \text{ g m}^{-2} \text{ day}^{-1}$  in C<sub>2</sub> (black gram - wheat + gram) to  $1.82 \text{ g m}^{-2} \text{ day}^{-1}$  in C<sub>7</sub> (maize + soybean - lentil) system. At 60-90 DAS the CGR ranges between  $0.56 \text{ g m}^{-2} \text{ day}^{-1}$  in C<sub>2</sub> (black gram - wheat + gram) to  $10.22 \text{ g m}^{-2} \text{ day}^{-1}$  in C<sub>4</sub> (cowpea - wheat + sarson). The highest CGR was observed in C<sub>4</sub> (cowpea - wheat + sarson) system with  $10.22 \text{ g m}^{-2} \text{ day}^{-1}$ , followed by the C<sub>5</sub> (okra - wheat + pea) at  $3.57 \text{ g m}^{-2} \text{ day}^{-1}$ . Lowest CGR was observed in C<sub>2</sub> (black gram - wheat + gram) system at  $0.56 \text{ g m}^{-2} \text{ day}^{-1}$ . Among intercrops at 60-90 DAS, the highest CGR was observed in C<sub>8</sub> (maize + cowpea - sarson) system with  $10.18 \text{ g m}^{-2} \text{ day}^{-1}$ , followed by the C<sub>9</sub> (maize + okra - pea) at  $3.77 \text{ g m}^{-2} \text{ day}^{-1}$ . The lowest CGR was observed in C<sub>6</sub> (maize + black gram - gram) at  $0.49 \text{ g m}^{-2} \text{ day}^{-1}$ . Among the systems intercrop like Lentil, cowpea resulted in higher N fixation compared to other legumes; also, the decomposition of the mulching provides the sufficient nutrients in the soil, resulted in enhanced dry matter accumulation. The application of Jeevamrit enhances soil microbial activity, leading to enhanced decomposition of ghanjeevamrit and mulching improves soil moisture, resulted in higher dry matter accumulation among the intercrop (Prasad and Srivastava 2019; Saharan et al. 2023). Similarly, Abdel-Wahab et al. (2019) and Manhas et al. (2021) found that the wheat - legume rotation, the CGR resulted in higher than in monocropping or continuous maize - wheat systems, during vegetative stage till harvest of crop growth.




Fig.1 Effect of cropping system on crop growth rate (g m<sup>-2</sup> day<sup>-1</sup>) under natural farming during *Rabi* 2020-21





Fig.2 Effect of cropping system on crop growth rate (g m<sup>-2</sup> day<sup>-1</sup>) under natural farming during *Kharif* 2020-21


### Relative Growth Rate (mg g<sup>-1</sup> day<sup>-1</sup>)

The Relative Growth Rate (RGR) was observed in the cropping system after sowing to till harvest during *Rabi* 2020-21 and *Kharif* 2021 (Fig. 3 and Fig. 4). Initially at 30 DAS the RGR ranges from 9.06 mg g<sup>-1</sup> day<sup>-1</sup> in C<sub>9</sub> (maize + okra - pea) to 18.02 mg

g<sup>-1</sup> day<sup>-1</sup> in C<sub>7</sub> (maize + soybean - lentil) system in **Rabi** 2020-21. At 120-150 DAS the RGR ranges between 0.50 mg g<sup>-1</sup> day<sup>-1</sup> in C<sub>8</sub> (maize + cowpea - sarson) to 8.89 mg g<sup>-1</sup> day<sup>-1</sup> in C<sub>7</sub> (maize + soybean - lentil) system. The highest RGR was observed in C<sub>7</sub> (maize + soybean - lentil) system with 8.89 mg g<sup>-1</sup> day<sup>-1</sup>, followed by the C<sub>3</sub> (soybean - wheat + lentil) system at 6.80 mg g<sup>-1</sup> day<sup>-1</sup>. Lowest RGR was observed in C<sub>8</sub> (maize + cowpea - sarson) system at 0.50 mg g<sup>-1</sup> day<sup>-1</sup>. Among intercrops at 120-150 DAS, the highest RGR was observed in C<sub>3</sub> (soybean - wheat + lentil) system with 4.26 mg g<sup>-1</sup> day<sup>-1</sup>, followed by the C<sub>2</sub> (black gram - wheat + gram) at 3.85 mg g<sup>-1</sup> day<sup>-1</sup>. Lowest RGR was observed in C<sub>4</sub> (cowpea - wheat + sarson) system at 1.69 mg g<sup>-1</sup> day<sup>-1</sup>. In subsequent, **Kharif** 2021, initially at 30 DAS, RGR among the cropping system ranging from 9.71 mg g<sup>-1</sup> day<sup>-1</sup> in C<sub>3</sub> (soybean - wheat + lentil) to 45.83 mg g<sup>-1</sup> day<sup>-1</sup> in C<sub>4</sub> (cowpea - wheat + sarson) system. At 60-90 DAS the RGR ranges between 2.19 mg g<sup>-1</sup> day<sup>-1</sup> in C<sub>7</sub> (maize + soybean - lentil) to 13.76 mg g<sup>-1</sup> day<sup>-1</sup> in C<sub>2</sub> (black gram - wheat + gram). The highest RGR was observed in C<sub>2</sub> (black gram - wheat + gram) system with 13.76 mg g<sup>-1</sup> day<sup>-1</sup>, followed by the C<sub>4</sub> (cowpea - wheat + sarson) at 11.72 mg g<sup>-1</sup> day<sup>-1</sup>. Lowest RGR was observed in C<sub>7</sub> (maize + soybean - lentil) system at 2.19 mg g<sup>-1</sup> day<sup>-1</sup>. Among intercrops at 60-90 DAS, the highest RGR was observed C<sub>6</sub> (maize + black gram - gram) system with 16.61 mg g<sup>-1</sup> day<sup>-1</sup>, followed by the C<sub>8</sub> (maize + cowpea - sarson) at 13.84 mg g<sup>-1</sup> day<sup>-1</sup>. Lowest RGR was observed in C<sub>9</sub> (maize + okra - pea) system at 6.46 mg g<sup>-1</sup> day<sup>-1</sup>. **This can be attributed to improved soil physio-chemical and biological properties by the application of jeevamrit, ghanjeevamrit and mulching with legume-based system.** The inclusion of legumes resulted in improved nutrient status also enhanced dry matter accumulation improved CGR and RGR (**Manhas et al. 2021**). In wheat and maize system, addition of legumes resulted in higher RGR by improving soil health and nutrient availability (**Abdel-Wahab et al. 2019**).



**Fig.3 Effect of cropping system on relative growth rate (mg g<sup>-1</sup> day<sup>-1</sup>) under natural farming during *Rabi* 2020-21**



**Fig.4 Effect of cropping system on relative growth rate (mg g<sup>-1</sup> day<sup>-1</sup>) under natural farming during *Kharif* 2021**


#### **NPK Uptake (kg ha<sup>-1</sup>)**

The data for NPK uptake in the seed and by-product for *Rabi* 2020-21 and *Kharif* 2021 presented in the Fig. 5, Fig.6 and Fig.7. A variation for the NPK uptake in seed and by-product was observed during both the crop season under natural farming.

#### **N Uptake (seed and by-product)**


In the *Rabi* 2020-21, the highest seed N uptake was observed in the C<sub>9</sub> (maize + okra - pea) system at 67.52 kg ha<sup>-1</sup>, followed by C<sub>7</sub> (maize + soybean - lentil) at 64.92 kg ha<sup>-1</sup>. The Lowest seed N uptake was observed in the C<sub>1</sub> (maize - wheat) system with 30.96 kg ha<sup>-1</sup>. In subsequent *Kharif* 2021, the highest seed N uptake was observed in the C<sub>5</sub> (okra - wheat + pea) system at 56.84 kg ha<sup>-1</sup>, followed by C<sub>9</sub> (maize + okra - pea) at 53.00 kg ha<sup>-1</sup>. The lowest seed N uptake was observed in the C<sub>2</sub> (black gram - wheat + gram) system with 24.26 kg ha<sup>-1</sup>. Overall system's, the highest seed N uptake was observed in the C<sub>9</sub> (maize + okra - pea) system at 120.52 kg ha<sup>-1</sup>, followed by C<sub>5</sub> (okra - wheat + pea) at 107.69 kg ha<sup>-1</sup>. The Lowest seed N uptake was observed in the C<sub>2</sub> (black gram - wheat + gram) system with 57.66 kg ha<sup>-1</sup>. In the *Rabi* 2020-21, the highest by-product N uptake was observed in the C<sub>7</sub> (maize + soybean - lentil) system at 18.43 kg ha<sup>-1</sup>, followed by C<sub>8</sub> (maize + cowpea - sarson) at 16.14 kg ha<sup>-1</sup>. The Lowest by-product N uptake was observed in the C<sub>9</sub> (maize + okra - pea) system with 6.66 kg ha<sup>-1</sup>. In subsequent *Kharif* 2021, the highest by-product N uptake was observed in the C<sub>2</sub> (black gram - wheat + gram) system at 34.10 kg ha<sup>-1</sup>, followed by C<sub>8</sub> (maize + cowpea - sarson) at 28.33 kg ha<sup>-1</sup>. The lowest by-product N uptake was observed in the C<sub>5</sub> (okra - wheat + pea) system with 3.50 kg ha<sup>-1</sup>. Overall system's, the highest by-product N uptake was observed in the C<sub>4</sub> (cowpea - wheat + sarson) system at 50.43 kg ha<sup>-1</sup>, followed by C<sub>2</sub> (black

gram - wheat + gram) at 46.85 kg ha<sup>-1</sup>. The Lowest by-product N uptake was observed in the C<sub>5</sub> (okra - wheat + pea) system with 11.55 kg ha<sup>-1</sup>.



**Fig.5 (a) N uptake in seed during *Rabi* 2020-21, (b) N uptake in seed during *Kharif* 2021, (c) N uptake in straw during *Rabi* 2020-21 and (d) N uptake in straw during *Kharif* 2021**

Natural farming inputs viz. jeevamrit, ghanjeevamrit and mulching resulted in improved soil microbial activity leads to N mineralisation and N fixation by legumes attributed in increased N availability; resulted in higher N uptake by plants. Mogale et al. (2023) found significantly higher nitrogen accumulation in the cowpea and subsequent cereal crop under cowpea - maize intercropping system. In lentil-based intercropping with cereals like wheat and barley resulted in improved in N uptake due to biological nitrogen fixation by lentils and cowpea (Kebede 2021 and Sharma T et al. 2023). Rotation with legumes such as soybeans under wheat-based system resulted in enhanced nitrogen content in both the seed and straw of subsequent cereal crops due to residual nitrogen from the legumes (Mesfin et al. 2023).



**Fig.6 (a) P uptake in seed during *Rabi* 2020-21, (b) P uptake in seed during *Kharif* 2021, (c) P uptake in straw during *Rabi* 2020-21 and (d) P uptake in straw during *Kharif* 2021**


### P Uptake (seed and by-product)

In the *Rabi* 2020-21, the highest seed P uptake was observed in the system C<sub>9</sub> (maize + okra - pea) at 41.29 kg ha<sup>-1</sup>, followed by C<sub>5</sub> (okra - wheat + pea) at 22.54 kg ha<sup>-1</sup>. The Lowest seed P uptake was observed in the system C<sub>7</sub> (maize + soybean - lentil) with 3.74 kg ha<sup>-1</sup>. In subsequent *Kharif* 2021, the highest seed P uptake was observed in the C<sub>5</sub> (okra - wheat + pea) system at 14.30 kg ha<sup>-1</sup>, followed by C<sub>9</sub> (maize + okra - pea) at 11.54 kg ha<sup>-1</sup>. The lowest seed P uptake was observed in the C<sub>2</sub> (black gram - wheat + gram) system with 1.94 kg ha<sup>-1</sup>. Overall system's, the highest seed P uptake was observed in the C<sub>9</sub> (maize + okra - pea) system at 52.83 kg ha<sup>-1</sup>, followed by C<sub>5</sub> (okra - wheat + pea) at 36.84 kg ha<sup>-1</sup>. The Lowest seed P uptake was observed in the C<sub>2</sub> (black gram - wheat + gram) system with 7.06 kg ha<sup>-1</sup>. In the *Rabi* 2020-21, the highest by-product P uptake was observed in the system C<sub>9</sub> (maize + okra - pea) at 10.97 kg ha<sup>-1</sup>, followed by C<sub>8</sub> (maize + cowpea - sarson) at 8.51 kg ha<sup>-1</sup>. The Lowest by-product P uptake was observed in the system C<sub>6</sub> (maize + black gram - gram) with 2.27 kg ha<sup>-1</sup>. In subsequent *Kharif* 2021, the highest by-product P uptake was observed in the C<sub>2</sub> (black gram - wheat + gram) system at 6.05 kg ha<sup>-1</sup>, followed by C<sub>8</sub> (maize + cowpea - sarson) at 5.85 kg ha<sup>-1</sup>. The lowest by-product P uptake was observed in the C<sub>5</sub> (okra - wheat + pea) system with 0.81 kg ha<sup>-1</sup>. Overall system's, the highest by-product P uptake was observed in the C<sub>4</sub> (cowpea - wheat + sarson) system at 14.49 kg ha<sup>-1</sup>, followed by C<sub>8</sub> (maize + cowpea - sarson) at 14.36 kg ha<sup>-1</sup>. The Lowest by-product P uptake was observed in the C<sub>6</sub> (maize + black gram - gram) system with 4.43 kg ha<sup>-1</sup>. Natural farming inputs viz. jeevamrit, ghanjeevamrit and mulching resulted in improved soil microbial activity attributed to enhanced P mineralization of native P and decomposition of ghanjeevamrit and mulching; resulted in improved P uptake. Tang et al. (2021) and Sharma et al. (2024) reported that significantly enhanced phosphorus use efficiency,

resulting in higher seed and biomass phosphorus uptake, when cereals intercropped legumes.

### K Uptake (seed and by-product)

In the **Rabi** 2020-21, the highest seed K uptake was observed in the C<sub>9</sub> (maize + okra - pea) system at 45.71 kg ha<sup>-1</sup>, followed by C<sub>5</sub> (okra - wheat + pea) at 3.32 kg ha<sup>-1</sup>. The Lowest seed K uptake was observed in the C<sub>7</sub> (maize + soybean - lentil) system with 3.34 kg ha<sup>-1</sup>. In subsequent **Kharif** 2021, the highest seed K uptake was observed in the C<sub>5</sub> (okra - wheat + pea) system at 24.52 kg ha<sup>-1</sup>, followed by C<sub>9</sub> (maize + okra - pea) at 16.63 kg ha<sup>-1</sup>. The lowest seed K uptake was observed in the C<sub>2</sub> (black gram - wheat + gram) system with 4.68 kg ha<sup>-1</sup>. Overall system's, the highest seed K uptake was observed in the C<sub>9</sub> (maize + okra - pea) system at 31.21 kg ha<sup>-1</sup>, followed by C<sub>5</sub> (okra - wheat + pea) at 24.84 kg ha<sup>-1</sup>. The Lowest seed K uptake was observed in the C<sub>2</sub> (black gram - wheat + gram) system with 5.84 kg ha<sup>-1</sup>. In the **Rabi** 2020-21, the highest by-product K uptake was observed in the C<sub>1</sub> (maize – wheat) system at 28.77 kg ha<sup>-1</sup>, followed by C<sub>2</sub> (black gram - wheat + gram) at 28.16 kg ha<sup>-1</sup>. The Lowest by-product K uptake was observed in the C<sub>9</sub> (maize + okra - pea) system with 7.52 kg ha<sup>-1</sup>.



**Fig.7 (a) K uptake in seed during Rabi 2020-21, (b) (f) K uptake in seed during Kharif 2021, (c) K uptake in straw during Rabi 2020-21 and (d) K uptake in straw during Kharif 2021**

In subsequent **Kharif** 2021, the highest by-product K uptake was observed in the C<sub>8</sub> (maize + cowpea - sarson) system at 73.63 kg ha<sup>-1</sup>, followed by C<sub>7</sub> (maize + soybean - lentil) at 55.49 kg ha<sup>-1</sup>. The lowest by-product K uptake was observed in the C<sub>5</sub> (okra - wheat + pea) system with 4.86 kg ha<sup>-1</sup>. Overall system's, the highest by-product K uptake was observed in the C<sub>8</sub> (maize + cowpea - sarson) system at 85.38 kg ha<sup>-1</sup>, followed by C<sub>1</sub> (maize – wheat) at 72.52 kg ha<sup>-1</sup>. The Lowest by-product K uptake was observed in the C<sub>5</sub> (okra - wheat + pea) system with 25.03 kg ha<sup>-1</sup>. **As nutrient uptake is dependent on the nutrient concentration and dry matter yield of the**

plant. Jeevamrit, ghanjeevamrit, mulching and legumes resulted in improved soil microbial activity attributed to enhanced dry matter by the increased availability and solubility of K in soil. While the low K uptake in crops due to low K content or less yield compared to other crops. The combination of legumes and cereals resulted in enhanced K uptake compared to sole cropping, overall increased yields and nutrient content (Chamkhi et al. 2022). Overall improved nutrient profile in intercropping legumes with cereals, including potassium uptake, particularly in systems like maize-legume intercropping (Namatsheve et al. 2020).

### Crude Protein

Crude protein content for the cropping system in **Rabi** 2020-21 and **Kharif** 2021, presented in the Table 5. In **Rabi** 2020-21, the highest crude protein was found in the C<sub>8</sub> (maize + cowpea - sarson) system at 75.50 %, followed by C<sub>6</sub> (maize + black gram - gram) at 71.48 %. The lowest crude protein was found in the C<sub>4</sub> (cowpea - wheat + sarson) with 8.58 %. The crude protein among the intercrops was found in the order C<sub>4</sub> (cowpea - wheat + sarson) > C<sub>2</sub> (black gram - wheat + gram) > C<sub>3</sub> (soybean - wheat + lentil) > C<sub>5</sub> (okra - wheat + pea). In **Rabi** 2020-21, the highest crude protein was found in C<sub>2</sub> (black gram - wheat + gram) system at 22.63 %, followed by C<sub>3</sub> (soybean - wheat + lentil) at 18.06 %. The lowest crude protein was found in the C<sub>1</sub> (maize - wheat) with 7.46 %. The crude protein among the intercrops was found in the order C<sub>6</sub> (maize + black gram - gram) > C<sub>7</sub> (maize + soybean - lentil) > C<sub>8</sub> (maize + cowpea - sarson) > C<sub>9</sub> (maize + okra - pea). The crude protein dependent on the dry matter accumulation, nutrient uptake as jeevamrit, ghanjeevamrit, mulching and legumes attributed to improved soil-physio-chemical and biological properties leads to higher nutrient uptake, resulted in enhanced crude protein content in the plant. The legumes significantly enhanced the crude protein content of the main crop due to their nitrogen-fixing ability (Li et al. 2023; Raza et al. 2023). The addition of legumes as intercrop (*viz.* black gram and cowpea) resulted in increased crude protein content in main crops (Varatharajan et al. 2022; Nyande et al. 2023 and Kumar et al. 2024). Legume as an intercrop improves main crops nutritional quality especially the crude protein content (Jensen et al. 2020).

**Table 5. Effect of cropping systems on crude protein in grains during crop season 2020-21 under natural farming**

Cropping system	Crude Protein			
	<i>Rabi</i> 2020-21	Intercrop	<i>Kharif</i> 2021	Intercrop
<b>C<sub>1</sub> Maize - wheat</b>	8.69	-	7.46	-
<b>C<sub>2</sub> Black gram - wheat + gram</b>	8.75	66.81	22.63	-
<b>C<sub>3</sub> Soybean - wheat + lentil</b>	8.75	50.75	18.06	-
<b>C<sub>4</sub> Cowpea - wheat + sarson</b>	8.58	75.06	13.81	-
<b>C<sub>5</sub> Okra - wheat + pea</b>	8.88	10.06	6.96	-
<b>C<sub>6</sub> Maize + black gram - gram</b>	71.48	-	7.73	20.13
<b>C<sub>7</sub> Maize + soybean - lentil</b>	52.13	-	7.69	16.00
<b>C<sub>8</sub> Maize + cowpea - sarson</b>	78.50	-	7.79	13.13
<b>C<sub>9</sub> Maize + okra - pea</b>	10.33	-	8.31	5.81

## Conclusion

The study evaluated Crop Growth Rate (CGR), Relative Growth Rate (RGR), NPK uptake and crude protein content across various cropping systems during *Rabi* 2020-21 and *Kharif* 2021. It was found that the significant variations in CGR and RGR among systems, with legume-based systems (e.g., C<sub>3</sub>: soybean - wheat + lentil, C<sub>4</sub>: cowpea - wheat + sarson, with C<sub>7</sub>: maize + soybean – lentil and C<sub>8</sub>: maize + cowpea - sarson). Enhanced NPK uptake in seeds and by-products was found significantly higher in legume-based cropping system compared to C<sub>1</sub> system under natural farming. The crude protein content was also found to be higher in legume-based systems (*viz.* C<sub>8</sub> and C<sub>2</sub>) under natural farming. The legume-based systems not only improve soil fertility, but also resulted in increase in dry matter accumulation, enhanced nutrient uptake and crude protein content under natural farming.

## DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declares that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

## References

- Abdel-Wahab TI, Abdel-Wahab SI, and Abdel-Wahab EI. Benefits of intercropping legumes with cereals. International Journal of Conference Proceedings, 2019, March;10, 225-234.
- AOAC. Methods of Analysis. Association of Official Agriculture Chemists, Washington DC 1970.
- Black CA. Methods of soil analysis. Part II. Chemical and mineralogical properties. American Society of Agronomy, Madison, Wisconsin, USA, 1965.
- Blackman VH. The compound interest law and plant growth. Annals of botany, 1919;33(131) 353-360.
- Blake GR. Bulk density. Methods of Soil Analysis: Part 1 Physical and Mineralogical Properties, Including Statistics of Measurement and Sampling, 1965;9, 374-390.
- Chamkhi, I, Cheto S, Geistlinger J, Zeroual Y, Kouisni L, Bargaz A and Ghoulam C. Legume-based intercropping systems promote beneficial rhizobacterial community and crop yield under stressing conditions. Industrial Crops and Products, 2022;183, p.114958.
- Devarinti SR. Natural farming: eco-friendly and sustainable. Agrotechnology. 2016;5(2), p.1000147.
- Jackson M. Soil chemical analysis prentice. Hall of India Private Limited, New Delhi. 1967;498(1).
- Jackson ML. Soil Chemical Analysis. Prentice Hall Inc. Englewood Cliffs, New Jersey, USA, 1973.
- Jena J, Maitra S, Hossain A, Pramanick B, Gitari HI, Praharaj S, Shankar T, Palai JB, Rathore A, Mandal TK, Jatav HS. Role of legumes in cropping system for soil

ecosystem improvement. *Ecosystem Services: Types, Management and Benefits*. Nova Science Publishers, Inc. 2022;415.

Jensen ES, Carlsson G and Hauggaard-Nielsen H. 2020. Intercropping of grain legumes and cereals improves the use of soil N resources and reduces the requirement for synthetic fertilizer N: A global-scale analysis. *Agronomy for sustainable development*, 2020;40(1), p.5.

Jones DB. Factors for converting percentages of nitrogen in foods and feeds into percentages of protein. United States Department of Agriculture Circular, No. 183. Washington, DC: USDA, 1941.

Kebede E. Contribution, utilization, and improvement of legumes-driven biological nitrogen fixation in agricultural systems. *Frontiers in Sustainable Food Systems*, 2021;5, p.767998.

Kumar N, Singh R, Agrawal RK, Sharma GD, Singh A, Sharma T and Rana RS. Optimizing forage harvest and the nutritive value of Italian ryegrass-based mixed forage cropping under northwestern Himalayan conditions. *Frontiers in Plant Science*, 2024;15, 1346936.

Li C, Stomph TJ, Makowski D, Li H, Zhang C, Zhang F and van der Werf W. The productive performance of intercropping. *Proceedings of the National Academy of Sciences*, 2023;120(2), p.e2201886120.

Manhas S, Singh J, Saini A, Sharma T and Parita K. Effect of tillage and fertilizer doses on growth and growth indices of soybean (*Glycine max* L.) under conservation tillage systems. *Environment Conservation Journal*, 2021;22(3), 181-186.

Mesfin S, Gebresamuel G, Haile M and Zenebe A. Potentials of legumes rotation on yield and nitrogen uptake of subsequent wheat crop in northern Ethiopia. *Heliyon*, 2023;9(6).

Mogale ET, Ayisi KK, Munjonji L and Kifle YG. Biological nitrogen fixation of cowpea in a No-till intercrop under contrasting rainfed agro-ecological environments. *Sustainability*, 2023;15(3), p.2244.

Namatsheve T, Cardinael R, Corbeels M and Chikowo R. Productivity and biological N<sub>2</sub>-fixation in cereal-cowpea intercropping systems in sub-Saharan Africa. A review. *Agronomy for sustainable development*, 2020;40(4), p.30.

Nyande A, George MS, Kassoh, FA and Bah AM. Effect of Intercropping and Crop Arrangement on Yield and Yield Components of Late Season Maize and Cowpea in the Upland of Njala Soil Series Southern Sierra Leone. *Journal of Experimental Agriculture International*, 2023;45(11), pp.178-189.

Olsen RS, Cole CV, Watanabe FS and Dean LA. Estimation of available P in soil by extraction with sodium bicarbonate; Circular United State Development of Agriculture 1954;19: 939.

Palekar S. Zero Budget Natural Farming. Subhash Palekar Natural Farming. Pingali P 2012."Green Revolution: Impacts, limits, and the path ahead." *Proceedings of the National Academy of Sciences*. 2016;109(31),12302–12308.

Piper CS. Soil and plant analysis. University of Adelaide, Australia. Hans publishers. Bombay. 1966; 223-237.

Raza MA, Zhiqi W, Yasin HS, Gul H, Qin R, Rehman SU, Mahmood A, Iqbal Z, Ahmed Z, Luo S and Juan C. Effect of crop combination on yield performance, nutrient uptake, and land use advantage of cereal/legume intercropping systems. *Field Crops Research*, 2023;304, p.109144.

Saharan BS, Tyagi S, Kumar R, Vijay Om, H, Mandal BS and Duhan JS. Application of Jeevamrit improves soil properties in zero budget natural farming fields. *Agriculture*, 2023;13(1), p.196.

Sharma T, Singh J, Madaik S, Kuma, P, Singh A, Rana B B, and Chauhan G. Organic input incorporation for enhancing sustainability and economic viability of cowpea in the North-Western Himalayan region. *Frontiers in Agronomy*, 2024;6, Article 1458603.

Sharma T, Singh J, Singh A, Sharma R, Chauhan G. Effect of organic nutrient sources on the yield, nutrient uptake and nodulation in Cowpea (*Vigna unguiculata*) under mid-hill conditions of Western Himalayas. *Environment Conservation Journal*. 2023 7;24(2):250-6.

Subbiah BV and Asija GL. A rapid procedure for the determination of available N in soils. *Current Science* 1956;25: 259-260.

Tang X, Zhang C, Yu Y, Shen J, van der Werf W and Zhang F. Intercropping legumes and cereals increases phosphorus use efficiency; a meta-analysis. *Plant and Soil*, 2021;460, pp.89-104.

Varatharajan T, Dass A, Choudhary AK, Sudhishri S, Pooniya V, Das TK, Rajanna GA, Prasad S, Swarnalakshmi K, Harish MN and Dhar S. Integrated management enhances crop physiology and final yield in maize intercropped with blackgram in semiarid South Asia. *Frontiers in Plant Science*, 2022;13, p.975569.

Walkley A and Black CA. An estimation of the method for determination of soil organic matter and a proposed modification of chromic acid titration method. *Soil Science* 1934;37: 29-39.

Watson DJ. The physiological basis of variation in yield. *Advances in agronomy*. 1952;4,101-145.