

Studies on path coefficient analysis and correlation to evaluate the relationships between yield and its components in Fenugreek (*Trigonella foenum-graecum* L.)

ABSTRACT

The present investigation was carried out for correlation studies and path coefficient analysis in Fenugreek (*Trigonella foenum-graecum* L.) The correlation of different traits with seed yield was examined both at phenotypic and genotypic levels and the study showed that genotypic correlation coefficient values were higher than the phenotypic values. This indicated that how much of phenotypic correlation coefficient are influenced by environment. The seed yield per plant was positive and significant with biological yield per plant, test weight and number of branches per plant at genotypic and phenotypic levels. The maximum direct positive effect on seed yield per plant was observed in biological yield per plant, harvest index, days to 50 per cent flowering, number of seed per pod, number of branches per plant and number of pods per plant. In the present investigation, genotypes viz., AFG-4 and RMT-303 were superior not only for seed yield per plant along with other but also related traits like biological yield, test weight and number of pods per plant. These diverse genotypes can be used in future breeding programme of fenugreek.

Keywords: Correlation analysis, Path coefficient analysis, Fenugreek, Seed yield, genotypes.

INTRODUCTION

Fenugreek (*Trigonella foenum-graecum* L.; $2n=16$) belongs to the family Fabaceae and is a multiuse and commercially important spice crop grown for its seeds, tender shoots and fresh leaves. It is an annual and self-pollinated seed spice that originated from Western Asia and South East Europe. The genus *Trigonella* consists of 50 species, most of which have an oriental origin in the Iranian/Indian region; 11 species occur in India, out of which *Trigonella foenum-graecum* L. (fenugreek) and *Trigonella corniculata* L. (kasurimethi /champaramethi) are commercially cultivated in India (Singhania *et al.*, 2006).

Fenugreek seeds and herbs are well known for their distinct aroma and slightly bitter taste. In the recent past, the use of fenugreek has been well recognized and demand is increasing due to its multifarious uses as food, spice, forage, medicinal plant, cosmetics and a source for natural diosgenin and galactomannan in the pharmaceutical and steroid industries (Rajagopalan, 2001; Moyer *et al.*, 2003 and Acharya, *et al.*, 2007). Dried seeds are used in many Indian dishes such as pickles and curries to bring special flavor and aroma. Chopped leaves are used as a flavoring agent, sprouts as salad and pot herbs as decoration. The Diosgenin content found in the leaves is of great medicinal value and is being used in contraceptive pills. Normally fenugreek seeds contain 0.4-1.26 percent of diosgenin. The seeds also contain saponins and fibers. Apart from its use as a seed spice, several medicinal uses of fenugreek have been also reported. From ancient times, it has been used to cure flatulence, dysentery, diarrhea, dyspepsia with lack of appetite and severe cough. Fibers present in the seed help lower blood sugar and thus helps in curing diabetes. It is also helpful in gastric problems, effective in regulating the digestive system and provides relief in joint pain of old age people. Seed is also used as a concentrated mixture to feed cattle which increases milk flow. Seeds of fenugreek contain 48% carbohydrates, 25.5% protein, 20% mucilaginous matter, 7.9% fat and 4.8% saponins. Apart from these properties, fenugreek seeds are rich in riboflavin (Vit B₂), niacin (Vit B₃), thiamin (B₁), folic acid, vitamin A, B₆ (pyridoxine), and vitamin C. Important mineral nutrients such as copper, potassium, calcium, iron, selenium, zinc, manganese and magnesium are also present in sufficient amount. Seeds taste a little bitter due to the presence of an alkaloid, called, —*Trigonellin*. Though seed spices are being used for a long time and their importance is known from even ancient times but still there is a lack of effort for the improvement of the genetic potential of these crops.

Comment [L1]: How the genotypes were stated that they are diverse without statistical analysis? Which statistical analysis says the genotypes were diverse?

The germplasm is used as the most valuable reservoir to provide variability for various traits that are a prerequisite for a successful breeding programme. Yield is a complex quantitative character highly influenced by environmental conditions, resulting in difficulty to determine whether the observed variability is heritable or not, for this, heritability is a suitable measure to estimate the heritable genetic portion of the total variability. The estimation of heritability along with genetic advances is generally more useful to provide a clear prediction of the breeding value. Correlation reveals only the direction and magnitude of association between any two traits however the path coefficient analysis partition the correlation into direct and indirect effects of various components on yield and serves as a powerful tool in the selection procedure accordingly.

Comment [L2]: Introduction should contain the importance of the analysis that included in the study. But heritability and genetic advance were not included in the study?

MATERIALS AND METHODS

The present investigation on fenugreek for correlation and path analysis was conducted at 12 genotypes, raised in randomized block design with three replications during the Rabi season 2023-2024. Observations were recorded on plot as well as single plant basis. Observations on plot basis were recorded for days to 50 per cent flowering, days to maturity, which observations based on single plant were recorded for plant height (cm), number of branches per plant, number of pods per plant, number of seed per pod, pod length, test weight (g), biological yield per plant (g), harvest index and seed yield per plant. For recording single plant observations five competitive plants were randomly selected from each plot. Average of these five plants with respect to plant height (cm), number of branches per plant, number of pods per plant, number of seed per pod, pod length, test weight (g), biological yield per plant (g), harvest index and seed yield per plant was used for statistical analysis. The genotypic co-variance was calculated as per Johnson et al., (1955). To estimate a cause and effect relationship the partitioning of genotypic and phenotypic correlation coefficient was done into direct and indirect effects by path analysis as suggested by Dewey and Lu (1959) and developed by Wright (1921).

Comment [L3]: What are those genotypes? Are they varieties or germplasm. Please mention. Add a table giving the details of the materials used.

Comment [L4]: Where does the experiment was conducted?

Comment [L5]: What is competitive mean? Do u mean random plants?

RESULTS AND DISCUSSION

Correlation coefficient

The study of correlation coefficient indicated the degree of association between the dependent variable i.e. yield and its contributing component traits. In the present study, thirteen characters were studied to find out the genotypic and phenotypic correlations with seed yield. It was found that the value of the genotypic correlation coefficient was higher than the phenotypic correlation coefficient value for all the characters under study. This indicated how much of the phenotypic correlation coefficient is influenced by the environment. The seed yield per plant had a positive and significant association with biological yield per plant, test weight and number of branches per plant. Whereas, negative and significant correlation expressed with days to 50 per cent flowering. These results were found in agreement with an earlier report by Chandra *et al.* (2000), Ayanoglu *et al.* (2004), Kole and Mishra (2006), Sharma & Sastry (2008), Singh & Pramila (2009), Kole and saha (2012), Pushpa *et al.* (2012), Yadav *et al.* (2013), Patil (2014), Gurjar *et al.* (2015), Wojoet *et al.* (2016), Mamatha *et al.* (2017), Kumar *et al.* (2018) and Singh *et al.* (2019).

Comment [L6]: Give the table or diagram of correlation and path co-efficient analysis, without which the results part is incomplete.

Comment [L7]: Discussion has only reports of earlier similar experiments. But should also contain your evaluation of the analysis results of the experiment.

Path analysis

The results of path coefficient based on genotypic coefficient taking seed yield per plant as dependent variable indicated that the characters biological yield per plant, harvest index, days to 50 per cent flowering, number of seed per pod, number of branches per plant and number of pods per plant showed positive direct effect on seed yield per plant while traits like test weight, plant height, pod

length and days to maturity exhibited negative direct effect on seed yield per plant. Similar finding also reported by Chandra and Sastry (2000), Balai *et al.* (2006), Sharma and Sastry (2008), Singh and Pramila (2009), Prajapati *et al.* (2010), Dashora *et al.* (2011), Fikreselassie *et al.* (2012), Yadav *et al.* (2013), Pathak *et al.* (2014), Kumari *et al.* (2015), Lodhi *et al.* (2015), Singh *et al.* (2016) and Kumar *et al.* (2018).

CONCLUSION

In the present investigation, genotypes *viz.*, AFG-4 and RMT-303 were superior not only for seed yield per plant but also related traits like biological yield, test weight and number of pods per plant. These diverse genotypes can be used in future breeding programme of fenugreek.

REFERENCES

- Acharya, S. N., J. E. Thomas, and S. K. Basu. (2007). Improvement in the medicinal and nutritional properties of fenugreek. In *Advances in medicinal plant research*, ed. S. N. Acharya and J. E. Thomas. Trivandrum, Kerala, India: Research Signpost.
- Ayanoglu, F.; Arslan, M. and Mert, A. (2004) Correlation and path analysis of the relationship between yield and yield components in fenugreek (*Trigonella foeniculum-graecum* L.). *Turkish Journal of Field Crops*. 9(1):11-15.
- Chandra, K. and Sastry, E.V.D. (2000). Genetic variation and character association of seed yield and its component characters in fenugreek. *Agriculture Science Digest*, 20(2): 93-95.
- Dewey, D.R. and Lu, K.H. 1959. A correlation and path coefficient analysis of components of Wheat grass seed production. *Agron J.*, 51: 515-518.
- Gurjar, M.; Naruka, I. S. and Shaktawat, R. P. S. (2015). Variability and correlation analysis in fenugreek (*Trigonella foenum-graecum* L.). *Legume Research*. 39(3):459-465.
- Johnson, H. W., H. F. Robinson and Comstock, R. E. (1955). Genotypic and phenotypic correlation in soybean and their implication in selection. *Agronomy Journal*. 47: 477-483.
- Kole, P. C. and Saha, A. (2012) Genetic divergence in fenugreek in rich and poor environments. *Journal of New Seeds*. 10(2):138-147
- Kole, P.C. and Mishra, A.K. (2006). Pattern of variability and associations among quantitative characters in fenugreek. *Indian Agriculturist*. 50(3/4): 93-96.
- Kumar, A., Pandey, V.P., Maurya, V.K., Tiwari, D. and Sriom. (2018). Genetic variability, heritability & genetic advance in fenugreek (*Trigonella foenumgraecum* L.). *International Journal of Chemical Studies*, 6: 153-156.
- Mamatha, N.C.; Tehlan, S.K.; Srikanth, M.; Ravikumar, T.; Batra, V.K.; Reddy, K.P. and Kumar, M., (2017). Variability Studies for Yield and Its Attributing Traits in Fenugreek (*Trigonella foenum-graecum* L.) Genotypes. *International Journal of Pure and Applied Bio science*, 5(3): 1075- 1079.

Moyer, J. R., S. N. Acharya, Z. Mir, and R. C. Doran. (2003). Weed management in irrigated fenugreek grown for forage in rotation with other annual crops. Canadian Journal of Plant Science, 83: 181-188.

Patil, J.; Vijayapadma, S.S. and Kopad, S. (2014). Genetic variability studies in fenugreek (*Trigonella foenum-graecum* L.). Research in Environment and Life Sciences, 9(12): 1482-1483.

Pushpa, T.N.; Chandregowda, M.; Srikantaprasad, D. and Gowda, A.P.M. (2012). Evaluation of fenugreek (*Trigonella foenum-graecum* L.) genotypes for growth and seed yield. Crop Research, 43(1, 2&3): 238-244.

Rajagopalan, M. S. (2001). Fenugreek is a savory medicinal. Supplement Industry Executive, 5(6): 43-44.

Sharma, K. C. and Sastry, E. V. D. (2008). Path analysis for seed yield and its component characters in fenugreek (*Trigonella foenum-graecum* L.). Journal of Spices and Aromatic Crops. 17(2):69-74.

Singh A. K., Singh D. R., Singh A., Maurya J. K., Pandey V. P. and Sriom (2019). Studies on character association and path analysis of yield with important yield contributing traits in Fenugreek (*Trigonella foenum-graecum* L.). Journal of Pharmacognosy and Phytochemistry 2019, 8(3): 4616-4619.

Singh, S. P. and Pramila (2009) Correlation and path analysis in fenugreek (*Trigonella foenum-graecum* L.). Asian Journal of Horticulture. 4(1):105- 107.

Wojo, A. A.; Alamerew, S.; Nebiyu, A. and Menamo, T. (2016). Genotype and phenotype variability studies in fenugreek (*Trigonella foenum-graecum* L.). Journal of Spices and Aromatic Crops. 25(2):159-168.

Wright, S. 1921. Correlation and causation J. Agric. Re., 20: 557-565.

Yadav, Y.; Yadava, P. S.; Pandey, V. P. and Kumar A. (2013) Genetic variability, correlation and path coefficient analysis studies in fenugreek (*Trigonella foenum-graecum* L.). Asian Journal of Horticulture. 8(2):456- 459.