

Groundnut (*Arachis hypogaea* L.) Response to Nitrogen Fertilizer and Earthing Up Practices in the Tigray Region of Ethiopia

ABSTRACT

To increase groundnut yields, integrated agronomic techniques must be implemented. The current study focuses on using starter nitrogen fertilizer in conjunction with earthing up for groundnut cultivation. Three replications and a factorial combination were used in the complete randomized block design of the experiment. Two types of earthing up (no earthing up and earthing up) and four levels of nitrogen fertilizer (0, 15, 30, and 45 kg ha⁻¹) were used in the treatment combinations. For each of the Hamedo 1 and Hamedo 2 experimental sites, two undisturbed and two disturbed composite soil samples were collected at a depth of 20 cm before sowing. The data analysis showed that neither the combination application of N fertilizer and earthing up nor the individual independent applications of N fertilizer and earthing up had a significant effect on groundnut flowering and maturity. Groundnut plant height, number of branches per plant, number of pods per plant, number of seeds per plant, seed weight per plant, and haulm yields were not significantly affected by earthing up alone or by the co-application of nitrogen fertilizer and earthing up. On the other hand, the exclusive use of nitrogen fertilizer significantly affected groundnut plant height, the number of branches per plant, the number of pods per plant, the number of seeds per plant, seed weight per plant, and haulm yield. Applying 45 kg ha⁻¹ of nitrogen fertilizer to the Hamedo 1 and Hamedo 2 trial sites enhanced the haulm yield by 70.53% and 78.76%, respectively, compared to the unfertilized plots. Individual applications of nitrogen fertilizer, earthing up, or a combination of both had a significant effect on groundnut pod and kernel yields. In comparison to plots that received a combined application of 45 kg ha⁻¹ of nitrogen fertilizer and earthing up, the Hamedo 1 and Hamedo 2 trial sites exhibited increases in pod yields of 57.73% and 55.97%, respectively, when earthing up was combined with 15 kg ha⁻¹ of nitrogen fertilizer. In the same way, applying 15 kg ha⁻¹ of nitrogen fertilizer to the earthing-up soil in the Hamedo 1 and Hamedo 2 experimental sites boosted the kernel yields by 70.05% and 75.43%, respectively, in comparison to soil earthing-up plots that received 45 kg ha⁻¹ nitrogen. The findings of the economic and agronomic analyses showed that applying nitrogen fertilizer and earthing up together greatly increases groundnut yields and revenues. Therefore, to increase groundnut yields and receive more financial advantages, farmers can consider the combined application of 15 kg ha⁻¹ of nitrogen fertilizer and earthing-up.

Keywords: Earthing-up, Groundnut, Starter nitrogen fertilizer

1. INTRODUCTION

In the tropics and subtropics, groundnuts (*Arachis hypogaea* L.) rank third and fourth, respectively, as sources of protein and oil [1, 2]. A versatile oil-seed legume, groundnuts have numerous health advantages. In addition to the nutritional benefits of peanuts, groundnut crops are crucial for soil replenishment. Through a process called biological nitrogen fixation (BNF), groundnuts fix atmospheric nitrogen and make it available to the host and other crops [3]. Globally, groundnut yield was estimated to be an average of 1.70 t ha⁻¹; in Africa, however, average yields are significantly lower, at 1.1 t ha⁻¹ [4]. Ethiopia's average yield was 1.70 t ha⁻¹, quite comparable to the global average yield [5]. In the Tigray region, where the specific study was conducted, groundnut is predominantly grown in warm lowland soils with sandy and sandy loam textures [6,7]. Such soils are ideal because they have loose, friable characteristics that allow for easier root and peg penetration, better water infiltration, and easier harvesting of pods. Under research farming in the Tigray region, groundnut yields reached about 2.66 t ha⁻¹ [7], which is significantly higher compared to the national production [5]. Low groundnut production at the national level is attributed to farmers' poor agronomic management practices.

Optimum groundnut yield is achieved when the soil has optimum nutrients [7]. More importantly, nitrogen (N) fertilizer is a widely required nutrient for growing all crops, due to its important role in the biochemical and physiological processes of the plant [8-10]. If the soil intended for groundnut planting has less than 15.5 kg N ha⁻¹, it can be supplemented with starter N for improved groundnut yield and quality [7]. Because legume crops can produce their own N through biological N fixation from the atmosphere, they rely less on synthetic N fertilizers during early growth [11,12]. Therefore, for improved kernel yield and protein production, 15 to 30 kg ha⁻¹ of starter N fertilizer is sufficient [7, 13].

Apart from fertilizer application, another important agronomic practice to improve groundnut yield is earthing up [14-16]. "Soil earthing up" is the process of placing soil around the base of the plant to cover the roots [16, 17]. The practice of soil earthing up creates a favorable environment for groundnut pegs to easily penetrate the soil, protects pods from sun exposure, hinders weed growth, allows water to seep into the soil, and promotes good air movement in the soil, thereby increasing groundnut productivity. Research reports suggest that earthing up the soil

for groundnut crops five weeks after sowing [16] and seven weeks after sowing [17] can boost groundnut yields.

The high protein content of groundnut crops, which are a major food source and help restore soil fertility, has piqued the interest of farmers in the Mereb Leke district, where this particular study was carried out. In addition to using groundnuts to disrupt pest and disease cycles, farmers can improve soil fertility by rotating groundnut with cereal crops [7].

However, due to a lack of information about the importance of N fertilizer in groundnut production, farmers grow groundnut without using N fertilizer. No research has been conducted on the significance of soil earthing up practices in enhancing groundnut productivity. Therefore, the objective of this study was to evaluate the contribution of soil earthing up in combination with N fertilizer in enhancing groundnut productivity.

2. MATERIALS AND METHODS

2.1. Site Description

During the 2019–2020 study period, two experimental sites (Hamedo 1 and Hamedo 2) in Medhin Tabia, Mereb Leke, Tigray, Ethiopia, were used for irrigation (Figure 1). The research area is located at an elevation of 1390 meters above sea level. Crop and livestock production are usually involved in mixed farming, which is a popular practice. Sorghum, finger millet, groundnut, and tef are the most common crops in the region, and they are grown in a rotation system. In order to produce fruits and vegetables, farmers in the research area also regularly use irrigation techniques. The research region is characterized by a semi-arid agroclimatic zone with well-drained sandy loam and loam textures as well as a distinctive luvisol soil type. The area is characterized by unimodal rainfall. The yearly rainfall was 636 mm, and the average monthly temperature was 25.3 °C [18].




Figure 1. Study area ('tabia' is the smallest administration unit).

2.2. Experimental Design, Treatments Set up and Management

Two farmers' fields were chosen for the experiment. Four levels of N (0, 15, 30, and 45 kg ha⁻¹) and two levels of earthing up (no earthing up coded as NEU and earthing up coded as EU) represented the eight combinations of treatments that were used. A randomized complete block design (RCBD) with three replications was used to lay out the treatments in a 2x4 factorial combination. The control treatment was the one that had neither earthing up nor N fertilizer applied at any level. Thirty days after sowing, before flowering, and seventy days after sowing, when flowering stopped, soil earthing up was employed. At planting, a full rate of N fertilizer was applied using a urea source. For sowing, a landrace "Sedi" variety with a seed rate of 80 kg ha⁻¹ was used. The planting plot was 3.6 m × 4 m (14.4 m²) with a net harvestable area of 9.6 m². The groundnut crop's blocks, plots, rows, and plants were spaced 1.5 m apart, 1 m apart, 0.6 m apart, and 0.1 m apart, respectively. There were five rows and 150 groundnut plants in a single groundnut plot. One border row was eliminated from either side of the plot in order to harvest groundnut. As advised by the surrounding area, agronomic management techniques were consistently implemented across all experimental plots, except for the treatments. Every plot was

given the same amount of irrigation water in line with the farmers' recommended irrigation practices.

2.3. Soil Sampling and Analyzed Parameters

Before planting, two samples of disturbed composite soil, with depths ranging from 0 to 20 cm, were collected for the experimental sites. These samples were permitted to pass through a 0.5 mm sieve for the measurement of organic carbon and total nitrogen and a 2 mm sieve for general analysis after being air-dried and milled. Two undisturbed soil samples were also collected using a core sampler for estimating soil bulk density. Texture, pH, electrical conductivity (EC), organic carbon (OC), total nitrogen (TN), available phosphorus (Av. P), cation exchangeable capacity (CEC), exchangeable potassium (Exc. K), exchangeable sodium (Exc. Na), exchangeable calcium (Exc. Ca), and exchangeable magnesium (Exc. Mg) were all measured on the collected samples. After oxidizing soil organic carbon, the Bouyoucos hydrometer method was used to measure the texture of the soil [19]. Using a pH meter, the pH level of the soil was determined in the supernatant suspension of a 1:2.5 soil-to-water ratio [20]. An electrical conductivity meter was used to test the electrical conductivity of the soil in the saturated extract of a 1:2.5 soil-to-water ratio [21]. The Walkley-Black chromic acid wet oxidation method was employed to determine the value of organic carbon [22]. The Kjeldahl approach was applied to estimate the TN of the soil [23]. Available P was estimated using the Olsen method [24]. Using an ammonium acetate solution for leaching, the soil's exchangeable base and cation exchange capacity were determined [25]. Table 1 provides a summary of the physical and chemical characteristics of soil.

Table 1. Soil physiochemical properties at the study sites before planting.

Soil properties	Experimental sites			
	Hamedo 1	Hamedo 2	Ratings	References
%Clay	8	11	NA	
%Silt	39	39	NA	
%Sand	53	50	NA	
Textural class	Sandy loam	Loam	NA	
Bulk density (g cm ⁻³)	1.44	1.41	Moderate	[26]
pH	5.62	5.71	Moderately acid	[26]
ECe (dS m ⁻¹)	3.02	2.98	Slightly saline	[26]

SOC (%)	0.63	0.66	Low	[26]
TN (%)	0.04	0.05	Very low	[26]
Av. P (ppm)	7.02	7.23	Low	[26]
Exc. K (cmol (+) kg ⁻¹)	0.25	0.23	Low	[26]
Exc. Na (cmol (+) kg ⁻¹)	0.18	0.17	Low	[26]
Exc. Ca (cmol (+) kg ⁻¹)	6.34	9.45	Moderate	[26]
Exc. Mg (cmol (+) kg ⁻¹)	7.41	7.05	High	[26]
CEC (cmol (+) kg ⁻¹)	15.33	18.37	Moderate	[26]

2.4. Agronomic Data Collected

2.4.1. Phenological Traits

Days to 50% flowering were measured from the days after planting when 50% of the groundnut plant in each experimental plot produced flowers. Days to 90% physiological maturity were measured from the days after planting when 90% of the groundnut plant in each planting plot turned their leaves color to yellow.

2.4.2. Vegetative Traits

A graduated tape meter was used to measure ten groundnut plant heights (cm) from the base of the plant to the top of the shoot once the plant achieved maturity. In order to analyze the plant height data, the mean of the 10 plant samples was then calculated. The average number of branches per plant for each plot was determined by dividing the total number of plant branches counted from the samples by the 10 sampled plants chosen for each plot.

2.4.3. Yield and Yield Components

After harvesting groundnut plants, ten sample plants from each plot had their pod counted and totaled. The total number of pods was then divided by ten plant samples to get the average number of pods per plant. Following the removal of the husks from the pods, a count of all the seeds in the 10 plant samples was made. The average number of seeds per plant was then calculated by dividing the total number of seeds counted in the 10 plant samples by the total number of plant samples. Each of the ten plant samples selected for each plot had its seed weight weighed. The weight of the ten plant samples was then totaled, and the mean seed weight per plant was determined by dividing the total seed weight by the ten plant samples. During harvesting stage, the above ground total fresh weight of groundnut biomass was measured from a

9.6 m² area of the experimental plot. A 200 g fresh weight sub-sample was taken from the fresh groundnut stover biomass, weighed, and oven-dried at 65°C for 72 hours to attain constant weight. The sub-samples were reweighed to ascertain the dry weight after they had dried in the oven. Using formula 1, the stover yield was computed from the obtained sub-sample fresh weight and dry weight values.

$$\text{Haulm yield} = \frac{\text{Total haulm fresh weight (g)} * \text{Sub-sample dry haulm weight (g)} * 10}{\text{Sub-sample haulm fresh weight (g)} * \text{net harea harvested (9.6 m}^2)}, \text{ Formula 1}$$

At harvesting, all mature pods from the plants uprooted from a 9.6 m² area were separated from the haulms, weighed and recorded. This was total pod fresh weight. A sub-sample of 200 g fresh pod weight was taken from the total quantity of the pods and then oven dried at 65°C for 72 hours and was used to estimate stable weight of pod and grain yields and adjusted to 10% moisture content and adjusted last measured weight of pod yield and grain yield was determined at Formula 2 and Formula 4, respectively.

$$\text{Pod yield} = \frac{\text{Total pod fresh weight (g)} * \text{Sub-sample dry pod weight (g)} * 10}{\text{Sub-sample pod fresh weight (g)} * \text{Net harea harvested (9.6 m}^2)}, \text{ Formula 2}$$

Shelling percentage as a yield component was calculated per experimental plot. The oven dried pod sub samples taken from the entire harvest of each experimental plots was weighed (pod dry weight) and then shelled. The shelled grain was weighed and then shelling percentage was estimated (Formula 2).

$$\text{Shelling percentage} = \frac{\text{Weight of shelled groundnut} * 100}{\text{Weight of unshelled groundnut}}, \text{ Formula 3}$$

From the measured shelling percentage, grain yield was estimated basing on the formula 4.

$$\text{Grain yield} = \frac{\text{Shelling percentage} * \text{Pod yield}}{100}, \text{ Formula 4}$$

2.5. Partial Budget Analysis

To assess how beneficial earthing up and fertilizer application were, a partial budget analysis following a procedure of CIMMYT [27] was used. Accordingly, price N fertilizer, N transport cost, N application and labor cost for earthing up were considered as total variable cost, whereas land preparation, weeding, harvesting and irrigation practice remained the same between the treatment. Before calculating gross revenue, groundnut kernel yield obtained from each experimental plot, were adjusted down by 10% to reflect farmers' practices. To estimate the prices of groundnut kernel yield, farm gate price was considered. Gross benefit was estimated by

multiplying the farm gate price that farmers receive for the kernel yield of groundnut when they sell it. Net benefit was calculated by subtracting the total costs from the gross benefits. The marginal rate of return (%) was calculated by dividing the change in net benefit by the change in cost and multiplied by 100.

2.6. Data Analysis

Finally, the collected agronomic data were subjected to statistical analysis of a two-way analysis of variance (ANOVA). Before data analysis, normality and homogeneity the data were checked across separately levels and combination of earthing up and N fertilizer treatments. Analysis of variance was performed using SAS software version 9.1. Means were separated using Fisher's protected least significance difference (LSD) test at 95% level of probability. Pearson's correlation coefficient was used to decide the relationship between the various plant phenology, growth, yield and yield component of groundnut.

3. RESULTS AND DISCUSSION

3.1. Phenological Attributes

Results indicated that each separate type of earthing up and separate individual levels of N fertilizer or combination of both agronomic managements have not brought significant effect on phenological traits of groundnut, namely time of flowering and maturity (Table 2). This is attributed to the innate behavior that groundnut could be produced satisfactory N through biological N fixation for commencement of flowering and maturation [7]. Instead of management practices, phenological parameters are more influenced by genetic and adaptation plant species to environmental stresses [28, 29]. There is a minimal growth prerequisite for plants to germinate, bloom, and mature within their normal timeframes [30].

Table 2. Effect of earthing up and N fertilizer on 50% flowering and 90% maturity of groundnut.

Earthing up	N (kg ha ⁻¹)	Days to flowering (50%)		Days to maturity (90%)	
		Hamedo site 1	Hamedo site 2	Hamedo site 1	Hamedo site 2
NEU	0	58.00	58.33	111.67	109.00
NEU	15	58.67	60.00	109.67	113.00
NEU	30	59.00	60.33	109.67	114.00
NEU	45	58.00	61.67	110.00	111.33
EU	0	56.00	59.33	113.33	108.67

EU	15	56.67	58.33	108.67	110.00
EU	30	59.00	59.33	112.67	110.67
EU	45	58.33	61.67	113.33	111.67
CV (%)		2.51	3.56	2.41	3.62
P value		0.37	0.72	0.51	0.81
LSD		NS	NS	NS	NS
Earthing up					
NEU		58.41	60.08	110.25	111.83
EU		57.50	59.67	112.00	110.25
CV (%)		2.51	3.56	2.41	3.62
P value		0.14	0.64	0.13	0.35
LSD		NS	NS	NS	NS
N (kg ha⁻¹)					
0		57.00	58.83	112.50	108.83
15		57.67	59.17	109.17	111.50
30		59.00	59.83	111.17	112.33
45		58.17	61.67	111.67	111.50
CV (%)		2.51	3.56	2.41	3.62
P value		0.15	0.14	0.21	0.48
LSD		NS	NS	NS	NS

3.2. Growth Attributes

3.2.1. Plant Height

For the sites of Hamedo1 and Hamedo2, plant height was significantly affected by sole N fertilizer, and there were no significant differences in plant height under individual separate use of earthing up and integrated application of earthing up and N fertilizer rates (Table 3). At the highest level of N (45 kg ha⁻¹), plant height increased by 29.11% for Hamedo 1 and 22.88% for Hamedo 2 compared to plant height from the unfertilized plot. For both sites, a consistent increment trend of groundnut plant height was counted from a consistent increment of N fertilizer. The correlation study showed a significant association between haulm yield and plant height and branch count. This demonstrated that as one parameter increased, the other one followed the trend, and vice versa (Table7). The increase in plant height was attributed to more available nutrient sources for a prolonged time from the higher nitrogen rate. N fertilizer is the most important plant nutrient in increasing vegetative growth as it enhances cell division and elongation[31]. Vegetative growth usually increases when fertilizer is applied regularly enough [32]. In line with the current result, plant height was improved after applying an increased rate of N fertilizer [7]. Since plant height and growth rate are indicators of plant health, enough amounts of nitrogen along with soil that is balanced in terms of P, K, and S are essential for plant growth

[33]. Additionally, it is thought that sufficient soil calcium and magnesium levels are essential for vegetative growth (Table 1).

3.2.2. Number of Branches per Plant

At both sites of Hamedo1 and Hamedo 2, number of branches per plant was significantly affected by sole N fertilizer, but not in the case of separate using of earthing up and integrated application of earthing up and N fertilizer rates (Table 3). At Hamedo1 and Hamedo2 sites, number of branches per plant were increased by 37.01% and 31.02%, respectively, when plots fertilized with 45 kg N ha⁻¹ relative to no fertilized plots. Number of branches per plant increased with N doses up to 45 kg ha⁻¹ in both sites. Since N fertilizer promotes cell division and elongation, it is the most crucial nutrient for plants to boost vegetative growth [31]. Vegetative growth usually increases when fertilizer is supplied adequately [32]. Similar to the current findings, the number of branches per plant increased after applying a higher rate of N fertilizer [7].

Table 3. Plant height and number of branches per plant in relation to earthing up and N fertilizer.

Earthing up	N (kg ha ⁻¹)	Plant height (cm)		Number of branches per plant	
		Hamedo 1	Hamedo 2	Hamedo 1	Hamedo 2
NEU	0	27.40	31.60	11.73	11.67
NEU	15	30.80	36.00	12.53	13.67
NEU	30	33.93	37.07	15.67	15.00
NEU	45	35.73	39.10	16.27	15.33
EU	0	28.27	32.40	11.67	12.00
EU	15	32.67	36.47	13.40	13.67
EU	30	34.00	37.00	15.60	15.67
EU	45	36.13	39.53	15.80	15.67
CV (%)		4.43	5.38	11.16	12.55
P value		0.72	0.98	0.90	0.99
LSD		NS	NS	NS	NS
Earthing up					
NEU		31.97	35.94	14.05	13.92
EU		32.77	36.35	14.12	14.25
CV (%)		4.43	5.38	11.16	12.55
P value		0.19	0.61	0.92	0.65
LSD		NS	NS	NS	NS
N (kg ha ⁻¹)					
0		27.83 ^D	32.00 ^C	11.70 ^B	11.83 ^B
15		31.73 ^C	36.23 ^B	12.97 ^B	13.67 ^{AB}
30		33.97 ^B	37.03 ^{AB}	15.63 ^A	15.33 ^A
45		35.93 ^A	39.32 ^A	16.03 ^A	15.50 ^A
CV (%)		4.43	5.38	11.16	12.55

P value	< .0001	< .0001	0.0004	0.008
LSD	1.75	2.38	1.92	2.16

3.3. Yield and Yield Components

3.3.1. Number of Pods per Plant

There was no discernible change in the number of pods in either the Hamedo 1 or Hamedo 2 study areas when soil earthing was applied alone or in conjunction with N fertilizer rates; however, there was a significant variance when N fertilizer rates were applied alone (Table 4). Plots of the planting sites of Hamedo1 and Hamedo2 have resulted in producing better number of pods per plant after fertilized with 15 kg N ha⁻¹, but decreasing trend was shown as the rates increased beyond. At Hamedo1 and Hamedo2, the average large number of pods of 29.97 and 31.47 were obtained from 15 kg N ha⁻¹, but it was insignificant when compared to the number of pods obtained from 30 kg N ha⁻¹. For each site, the minimum number of pods were achieved without N fertilizer application. Similar to the present findings, Getachew Gebrehana and Abeble Dagnaw [34] stated that application of starter N fertilizer at the rate of 18 kg ha⁻¹ resulted in higher number of pods per plant.

3.3.2. Number of Seeds per Plant

At the Hamedo 1 and Hamedo 2 research sites, the number of seeds per plant was significantly impacted by the sole application of N fertilizer; however, the number of seeds was not significantly changed by earthing up either alone or in combination with N fertilizer (Table 4). The highest number of seed per plant was obtained from the application of 15 kg ha⁻¹ N fertilizer but no significantly different seed numbers was obtained from the plots receiving 30 kg ha⁻¹ N fertilizer. The low number of seeds obtained at both study sites was from unfertilized plots and plots receiving 45 kg ha⁻¹ N fertilizer. Compared to unfertilized plot, 15 kg N ha⁻¹ resulted in 16.48% and 12.92% higher yield in Hamedo 1 and Hamedo 2, respectively. Therefore, from the present results, 15 kg ha⁻¹ N played a crucial role in improving healthy seeds. The higher seed numbers at the plots receiving 15 kg ha⁻¹ N fertilizer is due to the fact that legume crops use 15 kg ha⁻¹ N fertilizer as a starter to produce their own N. According to the research work of Tekulu et al. [7], they obtained better groundnut yields using 15 kg ha⁻¹ N fertilizer.

3.3.3. Weight of Seeds per Plant

The weight of seeds per plant changed significantly in the Hamedo 1 and Hamedo 2 research sites when N fertilizer was applied alone, but there was no discernible change in seed weight when earthing up was applied alone or in conjunction with N fertilizer (Table 5). Accordingly, 15 kg N ha⁻¹ produces better seed weights compared to plots received 45 kg N ha⁻¹ and plots without fertilizer. However, there was no significant difference in seed weight between the plots receiving 15 kg N ha⁻¹ and 30 kg N ha⁻¹. This might be due to groundnut crop fertilized with 15 kg N ha⁻¹ or 30 kg N ha⁻¹ had better seed filling than the plots received 45 kg N ha⁻¹ and unfertilized plots. It is known that most legume crops require N fertilizer in the early stages of growth until they produce their own N from the atmosphere. Except for the early legume growth, the main source of their food is the N they have produced from the atmosphere through their nodules. All the yield components such as number of pods per plant, number of seeds per plant and weight of seeds per plant improved at the same rate of 15 kg N ha⁻¹. According to the correlation result, there is a significant association between the number of pods per plant, the number of seeds per plant, the weight of seeds per plant, and the yields of pods and kernels. This demonstrated that as one parameter got higher, the other one followed the trend, and vice versa (Table 7).

Table 4. Number of pods per plant and number of seeds per plant under earthing up and N fertilizer.

Earthing up	N (kg ha ⁻¹)	Number of pods per plant		Number of seeds per plant	
		Hamedo 1	Hamedo 2	Hamedo 1	Hamedo 2
NEU	0	24.40	27.47	61.00	68.67
NEU	15	28.87	30.93	72.17	77.33
NEU	30	28.33	30.53	70.83	76.33
NEU	45	26.27	25.13	65.67	62.83
EU	0	27.07	28.27	67.67	70.67
EU	15	31.07	32.00	77.67	80.00
EU	30	29.30	30.80	73.25	77.00
EU	45	26.80	27.30	67.00	68.25
CV (%)		7.00	8.37	7.00	8.37
P value		0.75	0.92	0.75	0.92
LSD		NS	NS	NS	NS
Earthing up					
NEU		26.97	28.52	67.42	71.29
EU		28.56	29.59	71.40	73.98
CV (%)		7	8.37	7.00	8.37
P value		0.06	0.29	0.06	0.29
LSD		NS	NS	NS	NS
N (kg ha ⁻¹)					

0	25.73 ^C	27.87 ^{BC}	64.33 ^C	69.67 ^{BC}
15	29.97 ^A	31.47 ^A	74.90 ^A	78.67 ^A
30	28.82 ^{AB}	30.67 ^{AB}	72.04 ^{AB}	76.67 ^{AB}
45	26.53 ^{BC}	26.22 ^C	66.33 ^{BC}	65.54 ^C
CV (%)	7.00	8.37	7.00	8.37
P value	0.0056	0.0059	0.0056	0.0059
LSD	2.38	2.98	5.95	7.44

Table 5. Response of weight of seeds per plant and haulm yield to earthing up and N fertilizer.

Earthing up	N (kg ha ⁻¹)	Weight of seeds per plant (gram)		Haulm yield (t ha ⁻¹)	
		Hamedo 1	Hamedo 2	Hamedo 1	Hamedo 2
NEU	0	18.91	21.29	4.25	4.15
NEU	15	22.37	23.97	4.98	5.08
NEU	30	21.96	23.67	6.77	6.48
NEU	45	20.36	19.48	7.15	7.33
EU	0	20.98	21.91	4.36	4.23
EU	15	24.08	24.80	4.77	5.19
EU	30	22.71	23.87	6.55	5.80
EU	45	20.77	21.16	7.54	7.65
CV (%)		7.00	8.36	13.55	19.50
P value		0.75	0.92	0.89	0.87
LSD		NS	NS	NS	NS
Earthing up					
NEU		20.90	22.10	5.79	5.76
EUA		22.13	22.94	5.81	5.72
CV (%)		7.00	8.36	13.55	19.50
P value		0.06	0.29	0.95	0.92
LSD		NS	NS	NS	NS
N (kg ha⁻¹)					
0		19.95 ^C	21.60 ^{BC}	4.31 ^B	4.19 ^C
15		23.23 ^A	24.39 ^A	4.88 ^B	5.14 ^{BC}
30		22.33 ^{AB}	23.77 ^{AB}	6.66 ^A	6.14 ^{AB}
45		20.57 ^{BC}	20.32 ^C	7.35 ^A	7.49 ^A
CV (%)		7.00	8.36	13.55	19.50
P value		0.0056	0.0059	< .0001	0.0008
LSD		1.84	2.30	0.96	

3.3.4. Haulm Yield

The haulm yield was significantly impacted by the application of N fertilizer alone in the Hamedo 1 and Hamedo 2 sites, but not by the application of earthing up alone or in conjunction with N fertilizer (Table 5). In both research sites, the best haulm yield was obtained from the application of 45 kg N ha⁻¹ but this did not give any different yield than that at 30 kg N ha⁻¹. In both sites, the lowest yields were obtained from the plots that were not fertilized with N fertilizer. In Hamedo 1 and Hamedo 2, application of 45 kg ha⁻¹ N fertilizer increased yield

by 70.53% and 78.76% over unfertilized plots, respectively. Continuous increase in N fertilizer rates at both research sites showed continuous increase in haulm yield. High N fertilizer application is a need to achieve an improved fodder cowpea yields [35]. Similarly, for the present work, Tekulu et al. [7] have found that the use of high amounts of N fertilizer promotes plant vegetative growth and accordingly increases haulm yield. The reason why earthing up did not increase haulm yield is that earthing up practice mostly plays a major role in the creating suitable environment for the pegs that produces pods and thereby increases pod yield and kernel yield.

3.3.5. Pod Yield

For both experimental sites of Hamedo 1 and Hamedo 2, the application of N fertilizer and earthing up technique separately as well as in combination had significant effects on pod yield (Table 6). Accordingly, earthing up in combination with 15 kg ha⁻¹ N fertilizer provided an inspiring pod yield. In research site Hamedo 2, there was no significant difference in pod yield between the combined application of earthing up with 15 kg ha⁻¹ N fertilizer and earthing up with 30 kg ha⁻¹ N fertilizer. The lowest yields were obtained from plots that did not receive earthing up and N fertilizer. In Hamedo 1 and Hamedo 2 trial sites, application of earthing up with 15 kg ha⁻¹ N fertilizer increased yields by 57.73% and 55.97%, respectively, than plots received combined application of earthing up and 45 kg ha⁻¹ N fertilizer. Earthing up increases pod yield by combining with starter N. It is known that most legume crops require N fertilizer in the early stages of growth until they produce their own N from the atmosphere. Except for the early legume growth, the main source of their food is the N they have produced from the atmosphere through their nodules. Earthing up prevented the pods from being attacked by insects, prevented them from being exposed to sunlight, and allowed easily peg penetration, and thus resulted in an improved groundnut pod yield. Groundnut yield can increase with timely application of earthing up [36]. In support of the present findings, the work reported by Wakjira et al. [37] revealed better anchote yields using earthing up agronomic practice. According to the correlation result, there is a significant association between the number of pods per plant, the number of seeds per plant, the weight of seeds per plant, and the yields of pods and kernels. This demonstrated that as one parameter got higher, the other one followed the trend, and vice versa (Table 7).

Table 6. Response of pod yield and kernel yield to earthing up and N fertilizer.

Earthing up types	N (kg ha ⁻¹)	Pod yield (t ha ⁻¹)		Kernel yield (t ha ⁻¹)	
		Hamedo 1	Hamedo 2	Hamedo 1	Hamedo 2
NEU	0	2.61 ^{CD}	2.70 ^D	1.59 ^E	1.81 ^C
NEU	15	3.39 ^B	3.55 ^B	2.13 ^{BC}	2.35 ^B
NEU	30	2.81 ^C	3.04 ^C	1.88 ^{CD}	2.13 ^{BC}
NEU	45	2.54 ^D	2.67 ^D	1.54 ^E	1.84 ^C
EU	0	2.82 ^C	2.95 ^{CD}	1.64 ^{DE}	1.90 ^C
EU	15	3.84 ^A	4.18 ^A	2.60 ^A	3.07 ^A
EU	30	3.34 ^B	3.98 ^A	2.24 ^B	2.88 ^A
EU	45	2.45 ^D	2.68 ^D	1.52 ^E	1.75 ^C
CV (%)		4.32	5.80	8.09	10.87
P value		0.003	0.03	0.036	0.014
LSD		0.22	0.32	0.26	0.42
Earthing up types					
NEU		2.84 ^B	2.99 ^B	1.78 ^A	2.03 ^A
EU		3.11 ^A	3.45 ^A	2.00 ^B	2.40 ^B
CV (%)		4.32	5.80	8.09	10.87
P value		< .0001	< .0001	0.003	0.002
LSD		0.11	0.16	0.13	0.21
N (kg ha⁻¹)					
0		2.71 ^C	2.83 ^C	1.61 ^C	1.85 ^B
15		3.62 ^A	3.87 ^A	2.36 ^A	2.71 ^A
30		3.08 ^B	3.51 ^B	2.06 ^B	2.50 ^A
45		2.50 ^D	2.68 ^C	1.53 ^C	1.79 ^B
CV (%)		4.32	5.80	8.09	10.87
P value		< .0001	< .0001	< .0001	< .0001
LSD		0.16	0.23	0.19	0.29

3.3.6. Kernel Yield

Separate treatments of N fertilizer and earthing up, as well as their combined applications, had a substantial impact on groundnut kernel yields in the trial plots (Hamedo 1 and Hamed 2) (Table 6). Average highest kernel yields of 2.60 t ha⁻¹ for Hamedo 1 and 3.07 t ha⁻¹ for Hamedo 2 were obtained in the plots with earthing up and 15 kg N ha⁻¹. The lowest average kernel yields, with 1.52 t ha⁻¹ for Hamedo 1 and 1.75 t ha⁻¹ for Hamedo 2, were obtained from the earthing up plots that received application of 45 kg N ha⁻¹, without significant yield difference with plots with no earthing up practice and no fertilizer. In the Hamedo 1 and Hamedo 2 test areas, applying 15 kg ha⁻¹ N fertilizer to the earthing up soil increased the yield by 70.05% and 75.43%, respectively, relative to earthing up plots that received 45 kg N ha⁻¹. This finding revealed that earthing up with starter N fertilizer increases kernel yield. It is known that most legume plants require N fertilization during the initial growth period until they can produce N from the

atmosphere. Most of the legumes' N needs come from the atmosphere via N fixation through legume nodules. Additionally, earthing up allows the pegs to penetrate the soil and provides the right environment for pod development, protecting the pods from insects and sunlight and ultimately ensuring high grain yields. Therefore, effective use of the interaction of starter N fertilizer with earthing up had a significant effect on improving groundnut yields. Coinciding with current findings, Rima [38] reported improvement in carrot yield from integrated application of earthing up and 120 kg ha⁻¹ potassium (K) fertilizer. Application of combined use of natural manures and man-made fertilizers along with earthing up played a major role in increasing potato yield [39]. For groundnut crop cultivated in Hamedo 1 and Hamedo 2 sites, integrated application of soil earthing up and starter N fertilizer has resulted in positive and strong correlations between number of pods per plant, number of seeds per plant, weight of seeds per plant, pod yield, and kernel yield (Table 6). According to the correlation result, there is a significant association between the number of pods per plant, the number of seeds per plant, the weight of seeds per plant, pod yield, and kernel yield. This demonstrated that as one parameter got higher, the other one followed the trend, and vice versa (Table 7). Because kernel yield is measured from the cumulative effect of seeds produced in the individual plants, the positive and strong relationship between yield components is critical.

Separate application of earthing up also brought a significant effect on the kernel yield of groundnut (Table 6). Groundnut yield from earthing up plots had higher kernel yield relative to no earthing up plots. Confirming the present results, Ahmad et al. [15] found improved yield when earthing up is applied for different groundnut varieties. Application of earthing up, which does not overlap with the flowering period of groundnut, increases kernel yield [17]. In support of the present findings, the work reported by Wakjira et al. [37] observed better anchote yields using earthing up agronomic practice. Another similar work by Kugedera et al. [39] and Doboeh et al. [340] demonstrated that earthing up practices increased the marketable yield of potato tubers. Covering tuber crops with soil protects them from insect attack and exposure to sun and thus increases yield [37]. Contrary to the current findings, Desmae et al. [41] found that using soil earthing up did not increase groundnut yield but incurred unwanted expenses and reduced the returned profits.

Similar to the earthing up practices, the kernel yield of groundnut is also significantly affected by the separate application of N fertilizer (Table 6). Nitrogen fertilizer at the dose of 15 kg N ha⁻¹ was found to have an optimum rate for producing an inspiring groundnut kernel yield. According to the research work of Tekulu et al. [7], they obtained better groundnut yield using 15 kg ha⁻¹ N fertilizer. A small application of N, which can be called starter N, significantly increases cowpea yield [35]. Application of large amounts of N to soybean plants reduces N fixation capacity and thereby reduces yield [42]. This study was in good agreement with those reported by Getachew Gebrehana and AbebleDagnaw [34], Amante et al. [43], and Dabessa et al. [44] who showed that soybean production is affected by starter nitrogen use. In a study by Scott Tubbs et al. [45], in contrast to the current findings, application of starter N fertilizer did not cause any significant changes in groundnut yield.

3.4. Economic Feasibility

Under A partial and marginal analysis of adjusted groundnut grain yield with respect to soil earthing up and N fertilizer rates is shown in Table 8. Economic studies on groundnut crop yields show that each treatment has a different net benefit and marginal rate of return. As a result, soil earthing up plots treated with 15 kg ha⁻¹ N fertilizer produced the highest net benefit and marginal rate of return. In Hamedo site 1, the highest net benefit was 1740.40US\$ ha⁻¹, with an acceptable marginal rate of return (MRR) of 340%; in Hamedo site 2, the highest net benefit was 2158.73US\$ ha⁻¹, with an MRR of 6740%. Experimental plots without earthing up soils that received 15 kg ha⁻¹ N fertilizer were shown to have the best marginal rate of return and optimum net benefits, similar to the plots with earthing up soils that received 15 kg ha⁻¹ N fertilizer. In cases where two treatments exhibit an acceptable rate of return, the treatment offering the greatest net benefit is ultimately a recommendation. Therefore, it is recommended that farmers cultivate groundnuts using a combination of soil earthing up and 15 kg ha⁻¹ of starter N fertilizer.

UNDER PEER REVIEW

Hamedo 1										
	DF	DM	PH	NB	NP	NS	WS	HY	PY	KY
DF	1									
DM	-0.05 ^{NS}	1								
PH	0.22 ^{NS}	-0.13 ^{NS}	1							
NB	0.34 ^{NS}	0.22 ^{NS}	0.79***	1						
NP	0.16 ^{NS}	-0.17 ^{NS}	0.20 ^{NS}	0.17 ^{NS}	1					
NS	0.16 ^{NS}	-0.17 ^{NS}	0.20 ^{NS}	0.17 ^{NS}	1***	1				
WS	0.16 ^{NS}	-0.17 ^{NS}	0.20 ^{NS}	0.17 ^{NS}	1***	1***	1			
HY	0.21 ^{NS}	0.13 ^{NS}	0.81***	0.74***	0.04 ^{NS}	0.04 ^{NS}	0.04 ^{NS}	1		
PY	-0.11 ^{NS}	-0.33 ^{NS}	-0.11 ^{NS}	-0.23 ^{NS}	0.64***	0.64***	0.64***	-0.37 ^{NS}	1	
KY	-0.08 ^{NS}	-0.36 ^{NS}	0.09 ^{NS}	-0.09 ^{NS}	0.71***	0.71***	0.71***	-0.15 ^{NS}	0.90***	1
Hamedo 2										
	DF	DM	PH	NB	NP	NS	WS	HY	PY	KY
DF	1									
DM	0.35 ^{NS}	1								
PH	0.56**	0.29 ^{NS}	1							
NB	-0.01 ^{NS}	0.15 ^{NS}	0.63***	1						
NP	-0.36 ^{NS}	0.20 ^{NS}	-0.24 ^{NS}	-0.06 ^{NS}	1					
NS	-0.36 ^{NS}	0.20 ^{NS}	-0.24 ^{NS}	-0.06 ^{NS}	1***	1				
WS	-0.36 ^{NS}	0.20 ^{NS}	-0.24 ^{NS}	-0.06 ^{NS}	1***	1***	1			
HY	0.29 ^{NS}	0.17 ^{NS}	0.65***	0.57**	-0.13 ^{NS}	-0.13 ^{NS}	-0.13 ^{NS}	1		
PY	-0.40 ^{NS}	-0.06 ^{NS}	0.04 ^{NS}	0.15 ^{NS}	0.54**	0.54**	0.54***	-0.23 ^{NS}	1	
KY	-0.30 ^{NS}	0.02 ^{NS}	0.13 ^{NS}	0.07 ^{NS}	0.59**	0.59***	0.59***	-0.09 ^{NS}	0.87***	1

Table 7. Pearson's correlation coefficients (r) for various groundnut parameters.

Stars indicate significant effects *p (probability level) ≤ 0.05 , **p ≤ 0.01 , ***p ≤ 0.001 , NS = non-significant, DF = days to 50% flowering, DM = days to 90% maturity, PH = plant height, NB = number of branches per plant, NP = number of pods per plant, NS = number of seeds per plant, WS = weight of seed per plant, HY = haulm yield, PY = pod yield, KY = kernel yield and SH = shelling percentage

Table 8. Partial and marginal analysis for Kernel yield of groundnut under earthing up N fertilizer at the Mereb Leke district (Hamedo 1 and Hamedo 2 trial sites).

Hamedo 1							
TRT	TVC (US\$ ha ⁻¹)	KY (t ha ⁻¹)	AKY (t ha ⁻¹)	GB (US\$ ha ⁻¹)	NB (US\$ ha ⁻¹)	DA	MRR (%)
NEU + No Fertilizer	0	1.59	1.43	1118.04	1118.04	-	-
NEU + 15 kg N Fertilizer	29.62	2.13	1.92	1497.75	1485.03	11.82	1182
NEU + 30 kg N Fertilizer	42.34	1.88	1.69	1321.96	1296.52	D	
NEU + 45 kg N Fertilizer	55.06	1.54	1.39	1082.88	1044.72	D	
EU + No Fertilizer	75.12	1.64	1.48	1153.20	1078.08	D	
EU + 15 kg N Fertilizer	104.74	2.6	2.34	1828.24	1740.40	3.40	340
EU + 30 kg N Fertilizer	117.34	2.24	2.02	1575.10	1474.66	D	
EU + 45 kg N Fertilizer	130.18	1.52	1.37	1068.82	955.54	D	
Hamedo 2							
TRT	TVC (US\$ ha ⁻¹)	KY (t ha ⁻¹)	AKY (t ha ⁻¹)	GB (US\$ ha ⁻¹)	NB (US\$ ha ⁻¹)	DA	MRR (%)
NEU + No Fertilizer	0	1.81	1.63	1118.04	1272.74	-	-
NEU + 15 kg N Fertilizer	29.62	2.35	2.12	1497.75	1652.45	12.82	1282
NEU + 30 kg N Fertilizer	42.34	2.13	1.92	1321.96	1497.75	D	
NEU + 45 kg N Fertilizer	55.06	1.84	1.66	1082.88	1293.83	D	
EU + No Fertilizer	75.12	1.9	1.71	1153.20	1336.02	D	
EU + 15 kg N Fertilizer	104.74	3.07	2.76	1828.24	2158.73	6.74	674
EU + 30 kg N Fertilizer	117.34	2.88	2.59	1575.10	2025.13	D	
EU + 45 kg N Fertilizer	130.18	1.75	1.58	1068.82	1230.55	D	

Trt = treatment, TVC = total variable cost, KY = kernel yield, AKY = adjusted kernel yield, GB = gross benefit, NB = net benefit, DA = dominance analysis, D = dominated, MRR = marginal rate of return, urea fertilizer price = 39 US\$ 100 kg⁻¹, fertilizer transport cost = 1.25 US\$ 100 kg⁻¹, fertilizer application cost = 15.65US\$ ha⁻¹, groundnut kernel yield price = 781.3US\$ t⁻¹, labor cost = 3.13US\$ person⁻¹ day⁻¹, one US\$ = 32 Ethiopian birr.

4. CONCLUSION

The study found that nitrogen fertilizer significantly impacts plant height, number of branches, pods, seeds, seed weight, and haulm yield in groundnut plants. The use of nitrogen fertilizer alone led to increased haulm yields, plant height, and number of branches per plant in plots treated with higher nitrogen rates. Soil earthing up, combined with nitrogen fertilizer application, also enhanced groundnut productivity by protecting pods from pests and allowing pegs to penetrate the soil. The combination of 15 kg ha⁻¹ nitrogen fertilizer and earthing up significantly increased groundnut yields and profits, making it a viable option for farmers to increase groundnut production and gain financial advantages. To properly assist and advise small-scale groundnut farmers, additional demonstration trials of treatments in diverse agroecological zones are required.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

The authors declared that no generative AI tools, such as text-to-image generators or large language models (ChatGPT, COPILOT, etc.), were used in the writing or editing of the paper.

DATA AVAILABILITY

The available datasets used and analyzed during the study will be made available by the corresponding author upon reasonable request.

ACKNOWLEDGEMENTS

The authors express their thanks to the Tigray Agricultural Research Institute (TARI) for funding this work. Additionally, the Shire Soil Research Center provided lab facilities for soil investigation, for which we are thankful. Lastly, we would want to express our gratitude to the local farmers for their labor and cooperation in providing the trial field.

COMPETING INTERESTS

The authors declare that there aren't any competing interests.

REFERENCES

1. Taru VB, Kyagya IZ, Mshelia SI, Adebayo EF. Economic efficiency of resource use in groundnut production in Adamawa State of Nigeria. *World Journal of Agricultural Sciences*. 2008;4(5):896-900.
2. Ojiewo CO, Janila P, Bhatnagar-Mathur P, Pandey MK, Desmae H, Okori P, Mwololo J, Ajeigbe H, Njuguna-Mungai E, Muricho G, Akpo E. Advances in crop improvement and delivery research for nutritional quality and health benefits of groundnut (*Arachis hypogaea* L.). *Frontiers in Plant Science*. 2020;11:29.
3. Akram NA, Shafiq F, Ashraf M. Peanut (*Arachis hypogaea* L.): A prospective legume crop to offer multiple health benefits under changing climate. *Comprehensive reviews in food science and food safety*. 2018;17(5):1325-1338.
4. FAOSTAT. FAOSTAT statistical database. Publisher: FAO (Food and Agriculture Organization of the United Nations), Rome, Italy. 2017.
5. Central Statistical Authority (CSA). Federal Democratic Republic of Ethiopia: Central Statistical Agency: Agricultural Sample Survey. 2019
6. Sibhatu B, Tekle G, Harfe M. Response of groundnut (*Arachis hypogaea* L.) to different rates of phosphorus fertilizer at Tanqua-Abergelle District, Northern Ethiopia. *Basic Res. J. Agric. Sci. Rev.* 2016;5(1):24-29.
7. Tekulu K, Taye G, Assefa D. Effect of starter nitrogen and phosphorus fertilizer rates on yield and yield components, grain protein content of groundnut (*Arachis Hypogaea* L.) and residual soil nitrogen content in a semiarid north Ethiopia. *Heliyon*. 2020;6(10):1-12.
8. Mohapatra AK, Dixit L. Integrated nutrient management in rainy season groundnut (*Arachis hypogaea*). *Indian Journal of Agronomy*. 2010;55(2):123-127.
9. Xie KaiYun XK, Li XiangLin LX, He Feng HF, Zhang YingJun ZY, Wan LiQiang WL, David BH, Wang Dong WD, Qin Yan QY, Gamal MA. Effect of nitrogen fertilization on yield, N content, and nitrogen fixation of alfalfa and smooth bromegrass grown alone or in mixture in greenhouse pots. *J. Integr. Agric.* 2015;14:1864–1876.
10. Leghari SJ, Wahocho NA, Laghari GM, HafeezLaghari A, MustafaBhabhan G, HussainTalpur K, Bhutto TA, Wahocho SA, Lashari AA. Role of nitrogen for plant growth and development: A review. *Advances in Environmental Biology*. 2016;10(9):209-219.
11. Osborne SL, Riedell WE. Impact of low rates of nitrogen applied at planting on soybean nitrogen fixation. *Journal of plant nutrition*. 2011;34(3):436-448.
12. Ohyama T, Minagawa R, Ishikawa S, Yamamoto M, Hung NV, Ohtake N, Sueyoshi K, Sato T, Nagumo Y, Takahashi Y. Soybean seed production and nitrogen nutrition. A comprehensive survey of international soybean research-Genetics, physiology, agronomy and nitrogen relationships. 2013:115-157.
13. Argaw A. Integrating inorganic NP application and Bradyrhizobium inoculation to minimize production cost of peanut (*Arachis hypogaea* L.) in eastern Ethiopia. *Agriculture & Food Security*. 2018;7:1-9.
14. Ouédraogo M, Zagre M'bi B, Jørgensen ST, Liu F. Effect of mounding times on yield of Bambara groundnut (*Vigna subterranea* (L.) Verdc.) landraces in Sahel-Burkina Faso. *African Journal of Agricultural Research*. 2012;7(32):4505-11.
15. Ahmad N, Zada A, Ali A, Junaid M. Effect of earthing up procedure on enhancement in yield of different groundnut varieties planted under agro-climatic conditions of Malakand division. *Journal of Agriculture and Environmental Sciences*. 2015 ;4(1):181-184.
16. Thilini SP, Pradheeban L, Nishanthan K. Effect of Different Time of Earthing Up on

- Growth and Yield Performances of Groundnut (*Arachis hypogea* L.) Varieties. *J. Dry Zone Agri.* 2019; 5: 16–25.
17. Ouedraogo M, Zagre M'bi B, Liu F, Ortiz R, Jørgensen ST. Timing of mounding for bambara groundnut affects crop development and yield in a rainfed tropical environment. *Acta Agriculturae Scandinavica, Section B–Soil & Plant Science.* 2013;63(4):370-5.
 18. National Meteorological Agency of Federal Democratic Republic of Ethiopia (NMA). Mekelle; Tigray; Ethiopia; 2019.
 19. Bouyoucos GJ. Hydrometer method improved for making particle size analyses *Agronomy Journal.* soil. 962;54(5):464-5.
 20. Rhoades JD, Chanduvi F, Lesch SM. Soil salinity assessment: Methods and interpretation of electrical conductivity measurements. Food & Agriculture Org.; 1999.
 21. Richards LA, editor. Diagnosis and improvement of saline and alkali soils. US Government Printing Office; 1954.
 22. Nelson DW, Sommers LE. Total carbon, organic carbon, and organic matter. *Methods of soil analysis: Part 3 Chemical methods.* 1996 Jan 1;5:961-1010.
 23. Bremner JM, Mulvaney CS. Nitrogen—total. *Methods of soil analysis: part 2 chemical and microbiological properties.* 1982;9:595-624.
 24. Olsen SR. Estimation of available phosphorus in soils by extraction with sodium bicarbonate. US Department of Agriculture; 1954.
 25. Polemio M, Rhoades JD. Determining cation exchange capacity: A new procedure for calcareous and gypsiferous soils. *Soil Science Society of America Journal.* 1977;41(3):524-8.
 26. Hazelton P, Murphy B. Interpreting soil test results: What do all the numbers mean?. CSIRO publishing; 2016.
 27. CIMMYT Economics Program. From agronomic data to farmer recommendations: an economics training manual. CIMMYT; 1988.
 28. Souri, M.K., Hatamian, M. Aminochelates in plant nutrition: a review. *J. Plant Nutr.* 2019; 42 (1);67–78.
 29. Fattahi B, Arzani K, Souri MK, Barzegar M. Effects of cadmium and lead on seed germination, morphological traits, and essential oil composition of sweet basil (*Ocimum basilicum* L.). *Industrial Crops and Products.* 2019;138:111584.
 30. Shiferaw T. Optimization of Groundnut Crop Management Practices in Somali Regional State, Eastern Ethiopia. 2024.
 31. Lawlor DW, Lemaire G, Gastal F. Nitrogen, plant growth and crop yield. *Plant nitrogen.* 2001:343-367.
 32. Jeppsson N. The effects of fertilizer rate on vegetative growth, yield and fruit quality, with special respect to pigments, in black chokeberry (*Aronia melanocarpa*) cv. Viking'. *Scientia Horticulturae.* 2000;83(2):127-37.
 33. Aminifard MH, Aroiee H, Nemati H, Azizi M, Khayyat M. Effect of nitrogen fertilizer on vegetative and reproductive growth of pepper plants under field conditions. *Journal of plant nutrition.* 2012;35(2):235-242.
 34. Getachew Gebrehana Z, AbebleDagnaw L. Response of soybean to Rhizobial inoculation and starter N fertilizer on Nitisols of Assosa and Begi areas, Western Ethiopia. *Environmental Systems Research.* 2020;9:1-11.
 35. Abdul Rahman N, Larbi A, Kotu B, Marthy Tetteh F, Hoeschle-Zeledon I. Does nitrogen

matter for legumes? Starter nitrogen effects on biological and economic benefits of cowpea (*Vigna unguiculata* L.) in Guinea and Sudan Savanna of West Africa. *Agronomy*. 2018;8(7):120.

36. Desmae H, Sones H. Groundnut cropping guide. 2017.
37. Wakjira H, Garedew W, Nebiyu A, Shifaraw G. Effect of Earthing Up Frequency on Growth and Yield of Anchote (*Coccinia abyssinica*) Varieties at Gimbi District, Western Ethiopia. *International Journal of Agronomy*. 2024.
38. Rima SS, Mostarin T, Khatun K, Haq ME, Samad MA, Akter M, Shamim AS, Fatima K. Effect of potassium and earthing up on growth, yield and economic benefit of carrot. *Asian Journal of Research in Botany*. 2021;5(3):34-46.
39. Kugedera AT, Masvingise R, Kokerai L, Sakadzo N. Assessing the potential of earthing up and integrated nutrient management on Irish potato (*Solanum tuberosum* L.) productivity in smallholder farming systems. *International Journal of Agriculture Environment and Food Sciences*. 2022;6(2):303-310.
40. Dobocho M, Gedebo A, Haile A, Beshir HM. Improving potato productivity through optimum agronomic management to ensure food security of smallholder farmers. *Cogent Food & Agriculture*. 2022;8(1):2131995.
41. Desmae H, Sako D, Konate D. Impact of earthing up on groundnut (*Arachis hypogaea* L.) pod yield and its components. *AIMS Agriculture & Food*. 2022;7(3).
42. Da Silva PM, Tsai SM, Bonetti R. Response to inoculation and N fertilization for increased yield and biological nitrogen fixation of common bean (*Phaseolus vulgaris* L.). *Enhancement of Biological Nitrogen Fixation of Common Bean in Latin America: Results from an FAO/IAEA Co-ordinated Research Programme, 1986–1991*. 1993:123-130.
43. Amante G, Wedajo M, Temteme S. Soybean (*Glycine max* (L.) Merr.) response to application of mineral nitrogen and bradyrhizobia on Nitisols of Teppi, Southwest Ethiopia. *Heliyon*. 2024;10(9).
44. Dabessa A, Arega A, Takele F. Response of soybean [*Glycine max* L.(Merrill)] to starter nitrogen in Western Ethiopia. *Archives of Agronomy and Soil Science*. 2023;69(15):3690-3704.
45. Scott Tubbs R, Harris GH, Beasley JP, Smith AR, Smith NB. Effect of inoculant and nitrogen application at planting on peanut production in Georgia. *Crop Management*. 2012;11(1):1-11.