

**Mitigation of Daratumumab Interference in Pretransfusion Tests
by Addition of soluble CD38 to Patients' Plasma**

UNDER PEER REVIEW

Abstract

Background: CD38 is a protein highly expressed on myeloma (MM) cells that has been shown to be an effective target antigen for monoclonal antibody therapies, so treatment with anti-CD38 monoclonal antibodies is a first line therapy for patients with MM. Although CD38 is weaker expressed on erythrocytes anti-CD38 binds to CD38 on reagent RBCs and cause panreactivity *in vitro* and subsequent false positive reactions in indirect antiglobulin tests (IAT), antibody detection (screening) tests, antibody identification panels, and anti-human globulin (AHG) crossmatches. The aim of this study is to report our experience in evaluation of a new soluble CD38 based commercial method for mitigation of anti-CD38 interference in pretransfusion compatibility testing.

Methods: We evaluated the Grifols sCD38 methods, this test uses soluble CD38 to mitigate the anti-CD38 antibody interference in serological assays by adsorbing the drug in patient's plasma samples. Its performance was evaluated in 20 patients: 11 with negative IAT, of these 6 were treated with anti-CD38, 9 with positive IAT, of these 4 were treated with anti-CD38. The sCD38 based method has been compared to the current standard method using dithiothreitol (DTT 0.4M), which denatures the CD38 antigens on erythrocytes surface. All treated patients were in therapy with Daratumumab. After treatment plasma samples were further processed on the Grifols Erytra System as we internally validated the procedure.

Results: In our experience mean time to perform antibody screening and cross match was about 150 minutes using DTT method and about 50 minutes using sCD38 method. The sCD38 effectively mitigated the interference caused by anti-CD38 antibodies in 10 (100% efficacy) of patient samples tested while DTT was successful in only 9/10 (90% efficacy); no interference was observed in patients presenting anti erythrocyte antibodies. Moreover, there was no negative influence on DTT sensitive blood group systems such as KEL upon sCD38 treatment.

Conclusions: In our Laboratory, in the year 2023, we processed 129 patients treated with anti-CD38. Therefore the reduction from 150 to 50 minutes in the time needed to perform tests for mitigation of anti-CD38 interference appears to be relevant with a recovery of approximately 210 technical-hours per year. Another highly appreciated operational aspect was the possibility of treating the patient's plasma and perform tests using automatic instruments (Erytra Grifols) available in our laboratory. In the evaluation of this new method, we did not observe failures in the mitigation of anti-CD38 interference. Furthermore, the results obtained in the samples that presented allo or auto antibodies

were not affected by the treatment with sCD38. In our experience Grifols sCD38 assay is straightforward and quick to perform and it is superior to DTT treatment in the mitigation of anti-CD38 antibody interference in MM patients treated with Daratumumab.

Key Words

CD38, Daratumumab, Interference, Mitigation, Pretransfusion tests, Soluble CD38.

UNDER PEER REVIEW

Introduction

The CD38 antigen (cyclic ADP ribose hydrolase) is a transmembrane glycoprotein (PM approximately 31 kDa) that appears to be involved in three different cellular functions: adhesion protein, signal transduction, regulation of the calcium signalling pathway, and it is encoded by the *CD38* gene located on chromosome 4 [1,2]. The CD38 antigen is present on the surface of numerous cell lines such as B, T and NK lymphocytes, plasma cells but also on platelets and erythrocytes. CD38 is particularly expressed in neoplastic plasma cells of multiple myeloma (MM) [3,4]. Anti-CD38 monoclonal antibodies (MoAb) have a good therapeutic effect in patients with MM by inhibiting the growth and proliferation of neoplastic plasma cells both through direct mechanisms such as enzymatic inhibition and induction of apoptosis, and through immune-mediated mechanisms such as antibody-dependent or complement-dependent cellular toxicity [5,6].

The fact that CD38 is expressed on the surface of red blood cells implies a binding of anti-CD38 MoAb to erythrocytes, which will then be recognized and removed from the bloodstream at the level of the splenic reticuloendothelial system, mediated by binding to the Fc receptor. This extra-vascular haemolysis can lead to anaemia that in 25-30% of subjects will be of such an extent as to require transfusion support [7,8]. On the other hand, the presence of anti-CD38 MoAb on the surface of mature erythrocytes is the basis of the interferences observed in the Immunohematology Laboratory [7,8].

Currently, different anti-CD38 MoAbs with different characteristics are available or in an advanced stage of experimentation: Daratumumab (human IgG1-k), Isatuximab (chimeric IgG1-k), Felzartamab (IgG1-L), Mezagitamab (IgG1-L), which are directed towards different epitopes of the CD38 antigen, and which present different percentages of interference with pre-transfusion tests: from 100% observed for Daratumumab to 65% reported for Isatuximab. This interference, in the case of Daratumumab (Dara), can be detected up to 3-6 months after the last administration of the drug [9,10].

In Immunohematology Laboratory, DARA interference with serological tests occurs in all methods that use Coombs serum (antiglobulin serum) while usually the auto control test (patients' plasma with patients' RBCs in antihuman immunoglobulin) is negative because CD38 expression is downregulated during treatment therefore no haemolysis is observed *in vivo* [11,12]. Table I lists the main immunohematology tests with the possibility of interference from Dara.

To mitigate the interference of DARA in alloantibody screening and pretransfusion testing various methods have been proposed. For example, methods for the detection of anti-

erythrocyte antibodies have been proposed that do not involve the use of antiglobulin serum, such as tests in solid phase tube tests using potentiating agents such as polybrene; however, these methods have often demonstrated a less than fully satisfactory sensitivity in the detection of clinically significant antibodies [13-15]. Alternatively, methods have been proposed that involve the use of proteolytic enzyme-treated enzyme panels, however, enzyme treatment destroys some erythrocyte blood group antigens (i.e. M,N Lea and Leb) [16]. Treatment of test cells with dithiothreitol (DTT) is widely used in serological laboratories. DTT denatures CD38 on the cell surface by reducing disulfide bonds. However, DTT also destroys or modifies some other blood group antigens, e.g., KEL, DO, JMH, LU, IN, and YT, which results in impaired sensitivity to detect alloantibodies against these blood group antigens [17-20]. Recently has been suggested a method to overcome DARA interference by blocking the CD38 epitopes on RBCs by DARA-Fab fragments which were generated by papain proteolysis of DARA antibodies [21]. Blocking of the antigen binding site of DARA by incubation of patients' plasma with anti-idiotypic antibodies was also suggested and may also be intriguing; however, these antibodies are not commercially available. In addition, this assay requires umbilical cord RBCs as screening cells that are not typically available in a routine transfusion laboratory, and, furthermore, cord cells may have altered expression of some antigens [22]. An additional alternative to mitigate Daratumumab interference on pre-transfusion testing is to treat the patient's serum with soluble CD38 peptides with the intent of blocking the DARA binding site. [23,24]. See Table II for a quick overview of the methods that can be used to mitigate interference from anti-CD38 on pre-transfusion tests.

Aim of the study: In this paper we report a comparative evaluation of two methods for mitigation of anti-CD38 interference on pre-transfusion tests. As matter of facts we compared the in-house method, based on the pretreatment of red blood cells with Dithiothreitol (DTT) 0.2M, with a commercial method based on the pretreatment of patient plasma with soluble CD38 antigen (sCD38).

Materials and Methods

The Immunohematology Laboratory of Transfusion Medicine in Mestre:Dell'Angelo is a large General Hospital that acts as a provincial hub for a network of ten Spoke Hospitals in an area of about one million people. Our Transfusion Medicine has implemented a second-level immunohematology laboratory that has routine access to three different platforms for performing serological tests: test tubes, using a semi-automated in-house method; test cards, using a commercial automated method (Grifols); solid-phase test, using a

commercial automated method (Werfen). For genotyping we adopted a commercial automated method (Bead-Chip, Werfen). Our extended phenotyping protocol includes the study of the following erythrocyte blood group antigens: ABO, C, E, c, e, K, k, Kpa,b, Fya,b, Jka,b, M, N, S, s, Lua,b, P1, Xga, Lea,b. Our extended genotyping protocol includes the study of the genes coding for the following erythrocyte blood group antigens: C, E, c, e, K, k, Kpa,b, Jsa,b, Fya,b, Jka,b, M, N, S, s, Lua,b, Dia,b, Doa,b, Hy, Joa, Coa,b, Sc1,2, LWa,b. The Laboratory also can search for anti-platelet antibodies and anti-HLA antibodies using a Luminex analyser. In our Provincial Department of Transfusion Medicine (DIMIT) second level immunohematology assays, including the management of patients treated with anti-CD38, have been centralized in the Mestre Laboratory, from August 2023 to July 2024 our Laboratory processed 129 transfusion requests for patients treated with anti-CD38, all relating to patients with multiple myeloma treated with Daratumumab.

Mitigation of interference from anti-CD38 with DTT method: To mitigate interference from Daratumumab in our Laboratory we use a gel card method with pretreatment of red blood cells with 0.2M dithiothreitol (DTT) which can denature the CD38 antigen present on the surface of the red blood cells, minimizing the interference of the drug on immunohematology tests. To prepare the 0.2M DTT solution, dissolve 1 gram of DTT in 32 mL of PBS buffer or saline. Once prepared, the solutions can be aliquoted and stored for up to 5 days at +4°C, or frozen at -30°C; the aliquot of the DTT solution must be brought to room temperature and shaken gently before use. The red blood cells treated with DTT can be stored for up to 5 days at +4°C if resuspended in PBS or saline. Table III provides a brief description of the method. When interpreting the results, it must be kept in mind that treatment with DTT can destroy some erythrocyte blood group antigens (Kell, Penney, Sutter, Dombrock, Lutheran and others), thus making some antibody specificities undetectable [17-20].

Mitigation of anti-CD38 interference with sCD38 method: The test uses a protein reagent composed of a recombinant human CD38 protein (extracellular domain) in soluble form that, once mixed with the serum of the patient treated with Anti-CD38, binds to the monoclonal antibody neutralizing it, thus eliminating/reducing its interference on immunohematology tests such as the indirect Coombs test. The use of the sCD38 reagent does not remove the positivity of the indirect Coombs test due to the presence of alloantibodies, even if the reactivity of the anti-Fya and anti-Fyb antibodies may be decreased. All the necessary reagents are ready for use [23,24]. Table III shows a

schematic comparison between the 0.2M DTT method routinely used in our laboratory and the sCD38 method.

Samples: As reported in table IV for comparison of DTT and sCD38 methods efficiency in mitigation of DARA interference on pretransfusion tests we selected twenty samples: ten obtained from patients in treatment with DARA and ten obtained from patient not in therapy with DARA. In the second group five patients had a positive IAT (anti-E, anti-D, anti-K, anti-c, anti-Fya); in the first group four patients had a positive IAT (anti-c, anti Jka, anti-Fya, anti-M).

To study stability of sCD38 mitigation we performed a IAT in three samples obtained from patients in therapy with DARA for a MM, before sCD38 treatment, immediately after sCD38 treatment and aftermath every 24 hours for a week (storage performed at +4°C).

Pretransfusion tests: To perform the evaluations reported in this study, the methods currently in routine use at our Laboratory were used. For the search for anti-erythrocyte antibodies, we used a Liss-Coombs microcolumn method, testing the patient's serum against a three-cell screening. For each patient, two units of compatible concentrated red blood cells were cross-matched using a Liss-Coombs microcolumn method [25].

Results

In samples obtained from the ten subjects not in therapy with DARA no interference was detected in IAT. Five samples without anti-erythrocyte alloantibodies showed a negative IAT using native plasma tested against untreated or DTT-treated red cell panels; the sCD38-treated plasma was also non-reactive when tested against standard red cell panels. Five samples with anti-erythrocyte alloantibodies showed a positive IAT using native plasma tested against untreated or DTT-treated red cell panels; the sCD38-treated plasma was also reactive when tested against standard red cell panels. Considering ten MM patients treated with DARA, all presented a positive IAT when immunohematology assays were performed without methods to mitigate interference from anti-CD38. In six samples without anti-erythrocyte alloantibodies, DTT treatment eliminated anti-CD38 interference in five (83%) of them, while sCD38 treatment was effective in six patients (100%). The four patients with anti-erythrocyte alloantibodies remained IAT positive using both the DTT and sCD38 methods. These data were reported in table IV.

In figure 1 we reported the images of cards relating to the pre-transfusion tests: research on erythrocyte antibodies by IAT (three micro-columns on the left) and crossmatch in Liss-Coombs (two micro-columns on the right) in three patients treated with DARA without

alloantibodies. As can be observed, the three samples showed a marked positivity when tested without using methods to mitigate DARA interference. Using the 0.2M DTT method for the treatment of the red blood cells before performing the IAT and the crossmatch, we obtained a marked mitigation of the interference from DARA. However, the microcolumns were never frankly negative, thus requiring manual interpretation by the operator. Furthermore, in the third sample, one of the two crossmatches was positive. Using the sCD38 method to treat the patient's plasma before performing the IAT and crossmatch, we observed a complete elimination of DARA interference. Red blood cells had precipitated forming a compact layer on the bottom of the micro columns while the overlying phase appeared transparent making it possible to read the cards in complete automation. To evaluate the duration of the mitigation of interference from Daratumumab on pre-transfusion tests, we performed on selected patient on DARA therapy, we performed an IAT before the treatment with sCD38 and then for a week every 24 hours. As reported in figure 2, the IAT performed on untreated plasma showed a marked positivity from interference that was eliminated after treatment with sCD38 with a mitigation that remained constant for seven days.

The average execution time of the DTT method in our Laboratory was 150 minutes versus 50 minutes for the sCD38 method as reported on table III.

Discussion

In patients with Multiple Myeloma, the frequency of occurrence of anti-erythrocyte alloantibodies is rather low (around 3%). This reduced incidence of alloimmunizations can be attributed to immune depression due both to the disease itself and to its treatment [26-28]. However, despite these low levels of alloimmunization, considering the difficulties in identifying and characterizing any anti-erythrocyte alloantibodies, the SIMTI (Italian Society of Transfusion Medicine and Immunohematology) guidelines recommend that patients with Multiple Myeloma, before starting therapy with DARA, undergo at least typing for ABO, Rh, Kell, TCD and TCI. Extended serotyping and/or genotyping are also recommended [29]. We have therefore agreed with the Oncology and Hematology Departments of the Province of Venice that in all patients who are candidates for anti-CD38 therapy, before starting the treatment, a whole blood sample in EDTA will be taken which will be sent to our Immunohematology Laboratory for the execution of extended serotyping and genotyping [30]. If the patient comes to our attention after having already started therapy with anti-CD38, our protocol does not include the execution of extended phenotyping but only genotyping [29,30].

At our Transfusion Medicine Department (DIMIT), we perform extensive phenotyping and genotyping of regular donors of group O and A, selected based on the presence of particular Rh and Kk phenotypes, using high-throughput automated systems. So we have a database, directly connected with containing several thousand extensively typed regular donors that allows to select units of packed red blood cells with level III compatibility with the patient candidate for transfusion therapy [29,30].

Our protocol for assigning red blood cell concentrates in patients with MM treated with anti-CD38 is extremely prudent and includes both the search for irregular antibodies by IAT and the execution of the crossmatch (gel-card method with Liss-Coombs). All transfusion requests of patients treated with anti-CD38 and the related samples, are transferred to Mestre. In our immunohematology laboratory, the routine method for mitigation of DARA interference is pre-treatment with DTT (0.2M) both for the polyantigenic group zero test red blood cells used for antibody screening and for the red blood cells of the units to be transfused. This is an internationally validated method that has been validated in-house at our Laboratory [17-20]. This method is quite complex and requires trained and expert personnel for its execution. Furthermore, in our operational experience, it requires on average 150 minutes for the result and therefore is not suitable for managing urgent requests. Even from the point of view of interpretation of the results, trained and expert personnel are necessary since, as reported in figure 1, treatment with DTT, although mitigating the interference, will never give a complete negativization of the pre-transfusion tests. If the card shown in the figure is read automatically, the expert system for interpreting the results will give a weak positivity (+---) for TCI and an incompatibility for crossmatch. It will therefore be necessary, to proceed with a manual validation of the results. We therefore welcomed the possibility of adopting a method such as the one based on the mitigation with sCD38 of the interference from anti-CD38 on pre-transfusion tests which, although more expensive, promised to be quicker and easier to perform [31,32]. Also in this case, it is a method validated at an international level [33-36] and has been validated in house at our Laboratory [37]. Results obtained in this study have allowed us to highlight how the treatment of the patient's plasma with sCD38 is able to eliminate the interference from DARA on the IAT and on the crossmatches in an effective and complete manner without interfering with the detection of anti-erythrocyte alloantibodies possibly present in the samples. Moreover, the sCD38 based method is relatively simple that involves pre-treatment patients' plasma with a ready-to-use reagent. Plasma treated with sCD38 can be used to perform the necessary pre-transfusion tests using highly

automated analysers. The effective mitigation of interference allows the management of results through the middleware in use in our Transfusion Medicine, normally using the pre-set automated validation rules, although manual intervention by operators is always possible, of course. Plasma treated with sCD38, if stored at +4°C, shows persistent mitigation of DARA interference and pre-transfusion tests can be performed up to seven days after treatment. In our operational experience, the total time from plasma treatment with SCD38 to IAT and crossmatches results is about fifty minutes, therefore, in our opinion, this method is suitable for managing urgent requests [38]. Retrospectively considering the 129 samples of MM patients treated with DARA that we processed in 12 months, the treatment with sCD38 was completely effective in mitigation of the interference on pre-transfusion tests in 128 cases (99.2%). In this patient we repeated the plasma treatment by doubling the dose of sCD38, opting for the complete mitigation of the interference with complete resolution of the interference itself, presumably due to a high concentration of the drug not neutralized by the standard dosage of ligand.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Authors hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT etc.) and text-to-image generators have been used during writing or editing of the present manuscript

CONSENT

As per international standards, patient(s) written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standards, written ethical approval has been collected and preserved by the author(s).

References

1. Leleu X, Martin T, Weisel K, et al. Anti-CD38 antibody therapy for patients with relapsed/refractory multiple myeloma: differential mechanisms of action and recent clinical trial outcomes. *Ann Hematol.* 2022; 101: 2123-2137. doi: 10.1007/s00277-022-04917-5.
2. De Novellis D, Fontana R, Giudice V, et al. Innovative Anti-CD38 and Anti-BCMA Targeted Therapies in Multiple Myeloma: Mechanisms of Action and Resistance. *Int J Mol Sci.* 2022; 24:645. doi: 10.3390/ijms24010645.
3. Gozzetti A, Ciofini S, Simoncelli M, et al. Anti-CD38 monoclonal antibodies for multiple myeloma treatment. *Hum Vaccin Immunother.* 2022;18:2052658. doi: 10.1080/21645515.2022.2052658.
4. Shen F, Shen W. Isatuximab in the Treatment of Multiple Myeloma: A Review and Comparison With Daratumumab. *Technol Cancer Res Treat.* 2022; 21: 15330338221106563. doi: 10.1177/15330338221106563.
5. Offidani M, Corvatta L, Morè S, et al. Daratumumab for the Management of Newly Diagnosed and Relapsed/Refractory Multiple Myeloma: Current and Emerging Treatments. *Front Oncol.* 2021; 10 :624661. doi: 10.3389/fonc.2020.624661.
6. Yanq E, Mushem I, Samarkandi H, at al. Role of Anti-CD38 Monoclonal Antibodies in the Treatment of Adult Immune Hematological Disease. *Hematol Oncol Stem Cell Ther.* 2023; 17:4-12. doi: 10.56875/2589-0646.1108.
7. Mei Z, Wool GD. Impact of Novel Monoclonal Antibody Therapeutics on Blood Bank Pretransfusion Testing. *Hematol Oncol Clin North Am.* 2019; 33: 797-811. doi: 10.1016/j.hoc.2019.05.007.
8. Nedumcheril MT, DeSimone RA, Racine-Brzostek SE, et al. Overcoming Drug Interference in Transfusion Testing: A Spotlight on Daratumumab. *J Blood Med.* 2021; 12: 327-336. doi: 10.2147/JBM.S213510.
9. Chami B, Okuda M, Moayeri M, et al. Anti-CD38 monoclonal antibody interference with blood compatibility testing: Differentiating isatuximab and daratumumab via functional epitope mapping. *Transfusion.* 2022; 62: 2334-2348. doi: 10.1111/trf.17137.
10. Agrawal S, Chowdhry M, Karna P, et al. Daratumumab: The perplexity in immunohematology with emerging horizons in myeloma therapy. *Asian J Transfus Sci.* 2020; 14: 200-202. doi: 10.4103/ajts.AJTS_19_19.
11. Sullivan HC, Gerner-Smidt C, Nooka AK, Arthur CM, Thompson L, Mener A, et al. Daratumumab (anti-CD38) induces loss of CD38 on red blood cells. *Blood.* 2017;129: 3033–7. DOI: 10.1182/blood-2016-11-749432
12. Chapuy CI, Nicholson RT, Aguad MD, Chapuy B, Laubach JP, Richardson PG, et al. Resolving the daratumumab interference with blood compatibility testing. *Transfusion.* 2015 Jun;55(6 Pt 2):1545–54. DOI: 10.1111/trf.13069
13. Perram J, Blayney B, Ackerman L, et al. Solid phase antibody screening in the presence of anti-CD38 monoclonal antibodies: a potential alternative to avoid interference. *Pathology.* 2020; 52: 492-494. doi:10.1016/j.pathol.2020.01.683.
14. Zhou Y, Chen L, Jiang T, et al. 2-Mercaptoethanol (2-ME)-based IATs or Polybrene method mitigates the interference of daratumumab on blood compatibility tests. *Hematology.* 2021; 26: 365-370. doi: 10.1080/16078454.2021.1918916.
15. Feng CC, Chang CW, Lien ZY, et al. Better resolving of anti-CD38 antibody interference with blood compatibility testing by using manual polybrene method compared with dithiothreitol-pretreatment indirect antiglobulin test. *J Clin Lab Anal.* 2023; 37:e24891. doi: 10.1002/jcla.24891.

16. Carreño-Tarragona G, Cedena T, Montejano L, et al. Papain-treated panels are a simple method for the identification of alloantibodies in multiple myeloma patients treated with anti-CD38-based therapies. *Transfus Med.* 2019; 29: 193-196. doi: 10.1111/tme.12508. Epub 2018 Jan 25. PMID: 29369430.
17. Chapuy CI, Aguad MD, Nicholson RT, et al. DARA-DTT Study Group* for the BEST Collaborative. International validation of a dithiothreitol (DTT)-based method to resolve the daratumumab interference with blood compatibility testing. *Transfusion.* 2016; 56: 2964-2972. doi: 10.1111/trf.13789.
18. Furumaki H, Takeshita A, Ohto H, et al. A newly devised flow cytometric antibody binding assay helps evaluation of dithiothreitol treatment for the inactivation of CD38 on red blood cells. *Vox Sang.* 2021; 116:725-734. doi: 10.1111/vox.13052.
19. Gessoni G, Arreghini N, Pivetta M, et al. Adattamento in micro colonna delle metodiche per ridurre le interferenze da Daratumumab sui test pre trasfusionali. *Blood Transfusion* 2020; 18 suppl 2: s72.
20. Hosokawa M, Kashiwagi H, Nakayama K, et al. Distinct effects of daratumumab on indirect and direct antiglobulin tests: a new method employing 0.01 mol/L dithiothreitol for negating the daratumumab interference with preserving K antigenicity (Osaka method). *Transfusion.* 2018; 58: 3003-3013. doi: 10.1111/trf.14900.
21. Habicht CP, Ridders M, Grueger D, et al. Mitigation of therapeutic anti-CD38 antibody interference with fab fragments: How well does it perform? *Transfusion.* 2023; 63: 808-816. doi:10.1111/trf.17253.
22. Aung F, Spencer J, Potter D, et al. Efficient neutralization of daratumumab in pretransfusion samples using a novel recombinant monoclonal anti-idiotypic antibody. *Transfusion.* 2022; 62:1511-1518. doi: 10.1111/trf.17006.
23. Binda M, Favalaro V, Jody D, Berry JD, Schwind P. Novel Soluble CD38 for Efficient Neutralization of High Titer Anti-CD38 Antibodies. *Transfusion.* 2018; 58(S2): 171A.
24. National Health Service. International Blood Group reference Laboratory. Protein Development and Production Unit Bristol. Soluble recombinant CD38 (srCD38). <https://ibgri.blood.co.uk/ibgri-research-products/recombinant-blood-group-proteins/>. Citato 04 Aprile 2024.
25. Gessoni G. Immunohematology. In clinical and laboratory medicine textbook, Ciaccio M editor. First edition, Springer London; 2024.
26. Oostendorp M, Lammerts van Bueren JJ, et al. When blood transfusion medicine becomes complicated due to interference by monoclonal antibody therapy. *Transfusion.* 2015; 55:1555-62. doi: 10.1111/trf.13150.
27. Bullock T, Foser A, Clinkard B. Alloimmunisation rate of patients on Daratumumab: A retrospective cohort study of patients in England. *Transf Med* 2021; 31: 474-480. Doi 10.1111/tme.12808.
28. Zhan Ye, Laurie A. Wolf, Daniel Mettman, Fred V. Plapp. Risk of RBC alloimmunization in multiple myeloma patients treated by Daratumumab. *Vox Sanguinis* 2020; 115: 207-212. Doi 10.1111/vox.12864.
29. Matteocci A, Coluzzi S, De Martino S et al. Raccomandazioni per la gestione trasfusionale dei pazienti in trattamento con anticorpi monoclonali anti-CD38 e anti-CD47. Edizioni SIMTI 2023.
30. Coluzzi S, Londero D, Manfroi S, et al. Raccomandazioni per l'impiego delle metodiche in biologia molecolare in Immunoematologia. Edizioni SIMTI 2018.
31. Del Bello A, Kamar N, Cointault O, et al. Anti-CD38 Monoclonal Antibodies Interfere With Isoagglutinin Detection. *Transplantation* 2023; 107: e74-e75. doi: 10.1097/TP.0000000000004453.

32. Anani WQ, Marchan MG, Bensing KM, et al. Practical approaches and costs for provisioning safe transfusions during anti-CD38 therapy. *Transfusion*. 2017; 57: 1470-1479. doi: 10.1111/trf.14021.
33. Lejon Crottet S, Locher N, Caesar A et al. sCD38: a soluble protein for mitigation of therapeutic anti-CD38 in the indirect antiglobulin test. *SwisstransfusionRossschach* 2023.
34. Bise T, Moskwa C, Selleng J et al. Neutralization of anti-CD38 in patient samples by a soluble CD38 protein to allow alloantibody detection procedures. 56° annual meeting German Society for Transfusion Medicine and Immunohematology.(DGTI). Berlin 2023.
35. Engstrom C, Scherrer E, Falconer J, et al. A multicenter evaluation of soluble CD38 plasma pre-treatment to neutralize anti-CD38 pan-reactivity with RBCs, 34° regional ISBT Congress Cape Town 2023.
36. Legoff I, Dubosc-Marchenay N, Le Bouar M et al. Neutralisation par la molecule sCD38 des anti-CD38 plasmatiques en immuno-hematologie; une etude multicentrique. XXX congres de la Societe Francophone de Transfusion Sanguine – SFTS – Toulouse 2023.
37. Pivetta M, Vicentini P, Gessoni G. Evaluation of a soluble CD38 based method for mitigation of anti-Cd38 interference in pretransfusion compatibility testing. 35° regional ISBT Congress, Barcellona 2024.
38. Phou S, Costello C, Kopko PM, Allen ES. Optimizing transfusion management of multiple myeloma patients receiving daratumumab-based regimens. *Transfusion*. 2021; 61: 2054-2063. doi: 10.1111/trf.16425.

UNDER PEER REVIEW

Table I: Daratumumab interference in Immuno Haematology Assays.

Assays description	Interference
ABO typing	No interference
RhD typing	No interference
Test for D weak	Yes presence of Interference
RhCE typing	No interference
Kell antigen	No interference
k antigen	Yes presence of Interference
Autocontrol	No interference
Indirect Antiglobulin Test	Yes presence of Interference
Direct Antiglobulin Test	Yes presence of Interference
Crossmatch	Variable depending on the method used
Typing for other erythrocyte blood group antigens	Variable depending on the method used

This table schematically reports the routine immune haematological tests and the possibility of interference by treatment with Daratumumab [11,12]

Table II: Laboratory methods to mitigate Daratumumab interference in pretransfusion tests

Mitigation Strategy	Mechanism of action	Advantage	Disadvantages
Solid phase tests without antiglobulin serum [13]	DARA interference is reduced by not using antiglobulin serum.	Fast, simple and inexpensive,	sub-optimal sensitivity
Tube tests without antiglobulin serum [14,15]	DARA interference is reduced by not using antiglobulin serum.	Fast, simple and inexpensive,	sub-optimal sensitivity
Red blood cells treated with proteolytic enzymes [16]	Proteolytic enzymes destroy CD38 molecules on the surface of erythrocytes	Fast, simple and inexpensive,	Highly variable efficacy. Treatment with proteolytic enzymes destroys some erythrocyte blood group antigens
Treatment of red blood cells with DTT at various concentrations (0.04M, 0.1M, 0.2M). [17-20]	DTT denatures CD38 present on the surface of red blood cells	inexpensive	Long to perform, requires expert personnel. Destroys some erythrocyte blood group antigens
treatment of the patient's plasma with FAB fragment [21]	Fab fragments bind to CD38 and prevent it from binding to DARA	Commercial test. Simple and fast execution.	Expensive
treatment of the patient's plasma with anti-idiotypic antibodies. [22]	Anti-idiotypic antibodies neutralize DARA in the patient's plasma	Commercial test. Simple and fast execution.	Expensive
treatment of the patient's plasma with soluble CD38 [23,24]	il CD38 solubile neutralizza il DARA nel plasma del paziente	Commercial test. Simple and fast execution.	Expensive

This table schematically reports the various strategies that can be used to mitigate the interference of daratumumab on pre-transfusion tests. The first column schematically describes the mitigation strategy with the relative bibliographic references. The second column indicates the mechanism of action, the third column reports the advantages, and the fourth column reports the disadvantages of the proposed method.

Table III: Comparison of methods adopted in our Laboratory to mitigate Daratumumab interference on pre transfusion tests.


Routine Method: DTT 0.2M	New methos: Soluble CD38
Before performing the procedure, it is recommended to perform an antibody test (indirect Coombs test) using the plasma of the untreated patient.	Before performing the procedure, it is recommended to perform an antibody test (indirect Coombs test) using the plasma of the untreated patient.
Preparation of 0.2M DTT solution dissolve 1 g of DTT in 32 ml of PBS (pH 7.3). Divide into aliquots to be stored at -30°C.	_____
Defrost an aliquot of DTT 0.2M and wait until it reaches room temperature and mix. Allow reagents and samples to reach room temperature before performing the test.	Allow reagents and samples to reach room temperature before performing the test.
Select the test red blood cells to be used as controls: the positive control must be E+ (EE or Ee), the negative control must be K+ (Kk or Kk).	_____
Appropriately identify the tubes: screening panel cells, donor cells to be crossed, autocontrol, positive control, negative control.	Appropriately identify the tubes: screening panel cells, donor cells to be crossed, autocontrol, positive control, negative control.
Prepare the 3% red blood cell suspensions Red blood cells Text: Commercial panels are made up of a 0.8% red blood cell suspension, therefore the reagents must be prepared manually by centrifuging 500 microliters of solution, eliminating the supernatant and resuspending the bottom in 150 microliters of buffer solution. To prepare the positive and negative control, the self-control and the samples relating to the units to be compatibilized, resuspend 10 microliters of red blood cells in 300 microliters of buffer solution.	The red blood cell test panels are made up of a 0.8% red blood cell suspension and can therefore be used as such. To prepare a 1% suspension of the units to be compatibilized, resuspend 10 microliters of red blood cells in 1000 microliters of buffer solution
Wash the red cells of each tube 4 times with PBS buffer pH 7.3 (500 uL PBS for each wash). Add 400 uL (8 drops) of 0.2 M DTT to each tube. Mix well and incubate at 37°C for 30 minutes. During the incubation mix gently by inverting the tubes 3-4 times. Wash the red cells again as described above. Resuspend the washed red cells at 0.8%. The red cells treated with DTT can be stored at + 4°C for up to 5 days.	_____
Label Coombs card wells with cell number or type Inoculate 50 uL of red cell suspension Add 25 uL of patient plasma Incubate for 10 minutes at 37°C Centrifuge in a dedicated card centrifuge Examine each well for hemolysis and agglutination and record the results Interpret the results.	Pretreat plasma with 2 ul sCD38 Grifols for every 25 ul patient plasma to eliminate drug interference. Incubate for 10 minutes at 37°C. Dispense 25 ul pretreated plasma and 50 ul red cell suspension into the card well. Incubate for 10 minutes at 37°C Centrifuge in a dedicated card centrifuge Examine each well for hemolysis and agglutination and record the results Interpret the results.
Average time required to process a sample 150 minutes	Average time required to process a sample 50 minutes

Table IV: Samples selected for the comparison study

DARA therapy	Sample Description	IAT before Any treatment	IAT after DTT treatment	IAT after sCD38 treatment
NO	Negative	Negative	Negative	Negative
NO	Negative	Negative	Negative	Negative
NO	Negative	Negative	Negative	Negative
NO	Negative	Negative	Negative	Negative
NO	Negative	Negative	Negative	Negative
NO	Positive (anti-E)	Positive	Positive	Positive
NO	Positive (anti-D)	Positive	Positive	Positive
NO	Positive (anti-K)	Positive	Positive	Positive
NO	Positive (anti-c)	Positive	Positive	Positive
NO	Positive (anf.Fya)	Positive	Positive	Positive
YES	Negative	Positive	Negative	Negative
YES	Negative	Positive	Positive	Negative
YES	Negative	Positive	Negative	Negative
YES	Negative	Positive	Negative	Negative
YES	Negative	Positive	Negative	Negative
YES	Negative	Positive	Negative	Negative
YES	Positivo (anti-c)	Positive	Positive	Positive
YES	Positivo (anti-Jka)	Positive	Positive	Positive
YES	Positivo (anti-Fya)	Positive	Positive	Positive
YES	Positivo (anti-M)	Positive	Positive	Positive


IAT: Indirect antiglobulin test

Figure 1: Comparison of DTT and cSD38 methods for mitigation of Daratumumab interference in pre transfusion tests.



For each samples we performed indirect antiglobulin test (three micro-columns on the left) an a cross-match in liss.Coombs (two micro-columns on the right).

Figure 2: Mitigation of CD38 interference on pre transfusion test, evaluation of stability of treated plasma until 168 hours.



Using a sCD38 soluble based assay for mitigation of Daratumumab interference on pretransfusion test is maintained unchanged for up to seven days.