

Influence of major abiotic factors on the population trend of *Tetranychusurticae* Koch on okra at Samastipur, Bihar

Abstract:

Field experiments were conducted to study ~~of the~~ population fluctuation of ~~the~~ phytophagous mite (*Tetranychusurticae* Koch) in okra and its relation with different abiotic factors ~~during 2023 crop seasons~~ under ~~unprotected field~~ conditions ~~during Summer 2023~~ at Rajendra Prasad Central Agricultural University, Pusa, Samastipur, Bihar. The ~~findings on population of fluctuation~~ *Turticae* ~~of in okra revealed that mite were~~ commenced ~~from the~~ first fortnight of March and reached its peak of 34.60 mites/ 2.5cm² leaf area on ~~the~~ first fortnight of June. ~~A simple correlation (r) was established between population of mite and weather parameters in okra. It was observed that~~ There was a positive and significant correlation ~~of the population of~~ *Turticae* with maximum temperature ($r= 0.779^{**}$) and minimum temperature ($r= 0.699^{*}$), ~~and then it was observed that there is a negative and non significant relation with relative humidity ($r= -0.375$ morning 700 hrs) and ($r= -0.068$ evening 1400 hrs), positive non-significance with rainfall $r=0.306^{NS}$.~~

Keyword: Seasonal incidence, okra mites, Correlation, Tetranychidae

Introduction:

Okra, *Abelmoschus esculentus* (L.) Moench (family: Malvaceae), is one of the world's oldest, traditional cultivated vegetables. Its primary centre of origin in Africa probably Ethiopia and is grown in many Asian countries and other parts of the world. Within India, its cultivation occurs during both the summer and rainy seasons. It is commonly farmed as a summer crop in Northern India and is also prevalent as a winter crop in states like Gujarat, Andhra Pradesh, Karnataka, and Tamil Nadu. (Lal and Sinha, 2005). ~~In India, Out of that o~~ okra ~~have crop is cultivated in an~~ area of 0.531 million hectares and ~~its~~ production is 6.47 million tons ~~in India~~ (Anonymous, 2020-21).

The low yield ~~of okra~~ is attributed to the attack of different pests from sowing to harvesting. Recently, there ~~is has been~~ a change in agricultural scenario and mites are becoming serious ~~pest pests~~ in most ~~of the~~ crops and okra is no exception. The spider mite, *Tetranychusurticae* Koch (*Acari: Tetranychidae*), poses ~~a~~ serious threat to

Formatted: Font: Not Bold

Comment [mj1]: Add the information on the lowest population in which month and how much?

Comment [mj2]: Arrange alphabetically

Formatted: Font: Not Bold

okra crop particularly during spring, summer, and post-rainy seasons. This crop is infested mainly by six different mite pest species, viz., *Tetranychus urticae*, *T. macfarlanei*, *T. ludeni*, *Brevipalpus phoenicis*, *Polyphagotarsonemus latus*, and *Aceria lycopersici* (Gupta, 1985; Prasad and Singh, 2011). Out of these mite species, *T. urticae* is responsible for causing the loss of foliage of the crop plant resulting in a reduction of the economic yield of fruits ranging from 20-45 % depending upon cropping season and agro-climatic conditions. *T. urticae* is well adapted to various environmental conditions, causing loss of quality and yield or death of plants by sucking out the contents of leaf cells (Mondel and Ara, 2006, Kumaran *et al.*, 2007).

Comment [mj3]: Add author's name in all these species

Comment [mj4]: Remove space between number and % throughout the manuscript

T. urticae causes direct damage in terms of loss of chlorophyll, stunting of growth, stippling, webbing, leaf yellowing, defoliation, leaf burning, reduction in size and quality of fruits, the appearance of various types of plant deformities, followed by death, etc. which severely affect the yield and in extreme outbreaks, plant death. Indirect effects of mite feeding may include decreased photosynthesis and transpiration. Due to high reproductive potential and extremely short life cycle, combined with frequent acaricide applications this mite has developed resistance to almost all conventional pesticides in vogue (Chiasson *et al.*, 2004; Van Leeuwen *et al.*, 2005). The mites become serious pests because they have several generations per season. Phytophagous nature, high reproductive potential, and short life cycle contributed to rapid resistance development to many acaricides even after few applications (Devine *et al.*, 2001; Stumpf and Nauen, 2001). Since the degree of incidence of red spider mite changes with the season, it is desirable to have a thorough understanding of the seasonal incidence of the mite, which will lead to the development of suitable management programmes/programs. Hence, an attempt was made to correlate the effect of weather factors on the incidence and population dynamics of the spider mite in okra.

Materials and Methods:

Field trials were conducted from the first fortnight of March 2023 to the second fortnight of July 2023 during the Summer, 2022-23 at the Vegetable Research plot of Dr Rajendra Prasad Central Agricultural University, Pusa, Bihar. Okra cultivar of the variety "Kashi Kranti" was sown on 3rd February 2023. The plot selected

Formatted: Font: Italic

had uniform topography with ~~well-drained well-drained~~ sandy loam soil. ~~Latitude~~The ~~latitude~~, longitude, and altitude of the place are 85.67° E 25.98° N, and 52.92 m, respectively. ~~Climate~~The climate in this region is hot dry summers with mild ~~winter winters~~. Other climatic factors such as rainfall, temperature, and humidity in this region are highly favourable for the growth of vegetables and other crops in all seasons with the proliferation of ~~mite-mites~~ infesting a variety of crops.

A bulk-plot was raised to conduct research on the population fluctuation of major mites associated with okra. ~~The crop was cultivated by following appropriate agricultural practices and managed without the application of any insecticide.~~ The population of mites ~~were was~~ recorded at fortnight intervals. ~~Five places were selected for each host and three leaves were plucked randomly from host plants for recording population. All the leaves were mixed, labeled, and brought to the laboratory of the Department of Entomology, Dr. Rajendra Prasad Central Agricultural University, Pusa, Bihar for counting the mites. Care was taken to avoid disturbance to the natural population of mites on the leaf. All the mites were counted under a stereo binocular microscope. The whole population was recorded on 2.5cm^2 areas at four spots per leaf and the mean population was recorded.~~

The meteorological data were taken from ~~the meteorological section, Dr Rajendra Prasad Central Agricultural University, Pusa, Bihar.~~ ~~university observatory.~~ After that, these abiotic parameters were correlated with the corresponding pest population. ~~For testing the significance of the simple correlation coefficient, the 't' test was applied. The multiple regression coefficients and multiple correlation coefficient (R) of the mite population in relation to abiotic factors were also computed. The significance of all the multiple correlation coefficients (R) was tested for their significance by applying the 'F' test. The value of the coefficient of determination (R^2) was calculated to highlight the joint contribution of independent variables on mite population.~~

~~Okra cultivar was selected as a test variety "Kashi Kranti" and sown on 3rd February, 2023. The crop was cultivated by following appropriate agricultural practices and managed without the application of any insecticide.~~

Sampling and Observations

Formatted: Indent: First line: 0.5"

Population Fluctuation of two spotted mites, *Tetranychus urticae* Koch was found more injurious to the vegetables crop in the Samastipur district, Bihar. The investigations on population dynamics were carried out during the period from March 2023 to July 2023, on okra at vegetable research plot of Dr.RPCAU. The population was recorded from first fortnightly of March to second fortnightly of July. Five places were selected for each host and three leaves plucked randomly from host plants for recording population. All the leaves were mixed, labeled and were brought to the laboratory Department of Entomology, Dr. Rajendra Prasad Central Agricultural University, Pusa, Bihar for counting the mites. Care was taken to avoid disturbance to the natural population of mites on the leaf. All the mites were counted under a stereo binocular microscope. Whole population was recorded on 2.5cm² areas at four spot per leaves and mean population was recorded. The weather records were collected from the meteorological section, Dr Rajendra Prasad Central Agricultural University, Pusa, Bihar.

Correlation study

To investigate the effect of abiotic factors on the population of mites in this region. The observation for abiotic factors were recorded at fortnightly interval, used to compute the simple correlation coefficients (r) values of dependent (mite population) and various independent variables (abiotic factors i.e., temperature, relative humidity, and rainfall). For testing the significance of simple correlation coefficient 't' test was applied which is expressed as-

$$t = \frac{r\sqrt{n-2}}{\sqrt{1-r^2}}$$

Where, 'r' is the simple correlation coefficient and 'n' is total number of observation. The multiple regression coefficients and multiple correlation coefficient (R) of mite population in relation to abiotic factors were also computed.

The 't' value was compared with tabulated 't' on (n-2) degrees of freedom. The significance of all the multiple correlation coefficients (R) was tested for their significance by applying 'F' test as-

$$F = \frac{R^2}{1-R^2} \times \frac{n-k-1}{k}$$

Formatted: Normal, Justified, Space Before: 6 pt, Line spacing: 1.5 lines

Formatted: Justified, Indent: First line: 0", Space Before: 3 pt, After: 0 pt

Formatted: Indent: First line: 0", Space Before: 3 pt

Formatted: Justified, Indent: First line: 0.5", Space After: 0 pt

Where, 'n' is the total number of observation and 'k' is the total number of independent variables. The calculated 'F' was compared with the tabulated 'F' on 'k' and (n - k - 1) degrees of freedom. The value of coefficient of determination (R^2) was calculated highlight the joint contribution of independent variables on mite population. The correlation and regression analysis was done with the help of scientific calculator and personal computer (PC).

A linear regression of mite population on each variable (abiotic factors) was separately fitted. Regression equation is useful for the purpose of predicting (Y) when independent variables were assigned values. The purpose of trying out different regression was, to have an idea of which regression had a better fit. This is as follow-

$$\text{Linear regression } Y = a + bx$$

Results and Discussion:

The present investigation was carried out during okra crop seasons during 2023. Results of the present study revealed that population of phytophagous mites *Tetranychus T. urticae* Koch was recorded on okra in at different months during from the 1st fortnight of March to the 2nd fortnight of July in 2023 is presented in table Table1 and fig-Fig1. The *T. urticae* mite population started increasing from the March 2nd fortnight of March and reached its peak during the summer months. Maximum-The maximum population of *T. urticae* mite was recorded in the month of June 1st fortnight (34.57) during 2023 where the maximum temperature, minimum temperature ($^{\circ}\text{C}$), maximum relative humidity (%), minimum relative humidity, Maximum and Minimum, and total rainfall (mm) were 40.26 $^{\circ}\text{C}$, 23.88 $^{\circ}\text{C}$, 71.66%, 34.20%, 0.00mm, respectively and May 2nd fortnight (33.63) in 2023. The populations of *T. urticae* mite were gradually reduced from the June 2nd fortnight in 2023 and to the July 1st fortnight in 2023 with a decrease in temperature. Minimum-The minimum incidence of *T. urticae* mites was recorded in the month of March 1st fortnight (0.37), in years 2023

The present findings are somewhat consistent with that of Mandal *et al.* (2006) who observed that red spider mite populations on okra fluctuated significantly between 2000 and 2001, peaking during the 15th standard meteorological week and decreasing until the 24th SMW. The population was increased gradually from March to July. Similarly,

Formatted: Superscript

Formatted: Superscript

Kumaret al. (2015) observed ~~that the variation in the~~ population of red spider mite ~~mites~~ on okra ~~during the crop seasons of 2010 and 2011 in the Varanasi region. The findings indicated that mite infestation~~ was higher in May ~~while and~~ the temperature ~~of that~~ was also high ~~during this period.~~

Formatted: Normal, Space After: 6 pt, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers, Tab stops: -0.25", Left

Veerendra et al (2015) ~~examined-revealed that~~ the population ~~dynamics~~ of the red spider mite-*T. urticae* in grape vineyards, ~~focusing on its relationship with abiotic factors and a predator. The infestation~~ began in November 2012 and increased until April 2013.

Formatted: Normal, Indent: First line: 0", Space After: 6 pt, Don't adjust space between Latin and Asian text, Don't adjust space between Asian text and numbers, Tab stops: -0.25", Left

~~Correlation of *T. urticae* in okra with weather parameters during summer season 2023.~~

~~A simple correlation (r) was established between~~ The population of mite *T. urticae* ~~and weather parameters in okra (Table 2). It was observed that there was had~~ a positive and significant correlation with maximum temperature ($r = 0.779^{**}$) and minimum temperature ($r = 0.699^{*}$) (Table 2), ~~and then it was observed that t~~ There ~~was~~ a negative and ~~non-significant nonsignificant~~ relation of the *T. urticae* population with ~~maximum~~ relative humidity ($r = -0.375$ ~~morning 700 hrs~~) and ~~minimum relative humidity~~ ($r = -0.068$ ~~evening 1400 hrs~~), ~~The rainfall had a positive and non-significant~~ ~~teecorrelation with rainfall (r=0.306) with the *T. urticae* population. The significant correlation of both the temperature showed that there was a less difference in maximum and minimum temperature.~~ The regression equation for the data, with a population (Y) as the dependent variable and weather factors as the independent variables, was as follows ~~a determined coefficient of determination and (R^2) and regression model (Table 2):~~

Formatted: Font: Italic

$$Y = -184.644 + X_1(1.129) + X_2(4.626) - X_3(1.333) - X_4(0.810) - X_5(0.107)$$

X_1 = Maximum temperature, X_2 = Minimum temperature, X_3 = Maximum relative humidity, X_4 = Minimum relative humidity, X_5 = Rainfall.

Based on the calculation, the coefficient value (R^2) ~~was~~ derived as 0.7456 which suggests that 74% of the fluctuations in the red spider mite population were attributed to the impact of abiotic factors. In other words, ~~the~~ R^2 value indicates that per cent of ~~the~~ variation in the red spider mite population ~~were was~~ influenced by abiotic factors.

This finding ~~are is~~ somehow inconsistent with ~~my the~~ results of Mohanasundaram and Sharma (2011) who ~~recorded correlation between mite~~

~~infestations with important weather parameters~~ revealed that ~~the~~ population of *T. urticae* had a non-significant positive correlation with temperature (maximum, minimum, and average), maximum relative humidity and weekly total rainfall ~~where as while the population had a~~ significant positive correlation with minimum and average relative humidity.

The ~~present findings results obtained from my study~~ differ slightly from ~~these~~ the findings of Singh *et al.* (2018), ~~the results of the finding who~~ revealed that ~~the~~ mite population fluctuation showed a highly significant negative correlation with minimum temperature ~~moreover, and~~ a significant negative correlation ~~was observed~~ with maximum temperature and rainfall.

These findings obtained are inconsistent with ~~my the~~ results ~~where of~~ Ghosh (2019) ~~findings of correlation analysis between mite infestations and key weather parameters who~~ revealed that the mite population exhibited a ~~statistically~~ significant positive correlation with various temperature measures (maximum, minimum, and average). Additionally, there was a significant positive correlation with both minimum and average relative humidity.

CONCLUSION

~~These findings indicated that~~ population of ~~two spotted mites, Tetranychus~~ *T. urticae* Koch was observed in the experimental field during 1st fortnight of March 2023 and reached its peak ~~was the highest~~ during the 1st fortnight of June 2023 ~~i.e~~ 34.57/ 2.5 cm² leaf area. ~~Positive~~ There was a positive significant correlation ~~was observed in of the T. urticae population with~~ maximum temperature and minimum temperature ~~and also observed negative non significant correlation in relative~~ humidity.

REFERENCES

Formatted: Indent: First line: 0"

Benchasri S. Okra (*Abelmoschus esculentus* (L.) Moench) as a valuable vegetable of the world. *Field & Vegetable Crops Research/Ratarstvoipovrtarstvo*. 2012 Jan 1;49(1).

Singh P, Chauhan V, Tiwari BK, Chauhan SS, Simon S, Bilal S, Abidi AB. An overview on okra (*Abelmoschus esculentus*) and its importance as a nutritive vegetable in the world. *International journal of Pharmacy and Biological sciences*. 2014;4(2):227-33.

Gupta RC. Enhanced sensitivity of 32P-postlabeling analysis of aromatic carcinogen: DNA adducts. *Cancer research*. 1985 Nov 1;45(11_Part_2):5656-62.

Prasad A, Singh V. Behavior of confined square and rectangular footings under eccentric-inclined load. *International Journal of Geotechnical Engineering*. 2011 Apr 1;5(2):211-21.

Kumaran A, Karunakaran RJ. In vitro antioxidant activities of methanol extracts of five *Phyllanthus* species from India. *LWT-Food Science and Technology*. 2007 Mar 1;40(2):344-52.

Chiasson MW, Davidson E. Taking industry seriously in information systems research. *MIS quarterly*. 2005 Dec 1:591-605.

Van Leeuwen R, Boekhoorn S, Vingerling JR, Witteman JC, Klaver CC, Hofman A, de Jong PT. Dietary intake of antioxidants and risk of age-related macular degeneration. *Jama*. 2005 Dec 28;294(24):3101-7.

Devine DJ, Clayton LD, Dunford BB, Seying R, Pryce J. Jury decision making: 45 years of empirical research on deliberating groups. *Psychology, public policy, and law*. 2001 Sep;7(3):622.

Stumpf N, Zebitz CP, Kraus W, Moores GD, Nauen R. Resistance to organophosphates and biochemical genotyping of acetylcholinesterases in *Tetranychusurticae* (Acari: Tetranychidae). *Pesticide Biochemistry and Physiology*. 2001 Feb 1;69(2):131-42.

Veerendra AC, Udikeri SS, Karabhantanal SS. Dynamics of two spotted red spider mite *Tetranychusurticae* Koch (Acari: Tetranychidae) in grape vineyards and its co-relation with abiotic factors and a predator. *Journal of Entomology and Zoology studies*. 2015;3(6):373-6.

Kumar S, Ahlawat W, Kumar R, Dilbaghi N. Graphene, carbon nanotubes, zinc oxide and gold as elite nanomaterials for fabrication of biosensors for healthcare. Biosensors and Bioelectronics. 2015 Aug 15;70:498-503.

Comment [mj5]: Check the guidelines and make accordingly

UNDER PEER REVIEW

Table 1: Population fluctuation of *Tetranychusurticae* on okra during the crop period Summer 2023

Interval	*Mean population of <i>T. urticae</i> mites/2.5cm ² leaf area	Temperature (°C)		Relative humidity (%)		Total Rainfall (mm)
		Maximum	Minimum	Maximum	Minimum	
1 st Fortnight March	0.37	31.11	14.67	93.33	48.67	0.00
2 nd Fortnight March	2.53	30.18	16.73	89.06	50.56	18.00
1 st Fortnight April	8.23	35.67	16.20	76.80	27.46	0.00
2 nd Fortnight April	13.43	35.80	20.73	71.06	34.27	0.00
1 st Fortnight May	30.10	35.71	19.93	75.07	37.20	1.40
2 nd Fortnight May	33.63	35.20	22.52	82.56	52.18	16.80
1 st Fortnight June	34.57	40.26	23.88	71.66	34.20	0.00
2 nd Fortnight June	29.67	36.27	25.36	86.20	55.20	92.60
1 st Fortnight July	25.30	33.22	19.70	89.20	50.40	59.40
2 nd Fortnight July	6.20	32.45	20.20	85.30	51.20	24.00

* Mean of 5 plants per plot, 3 leaves of each plant and one leaf 2.5 cm² areas of four spot spots.




Fig 1-Influence of weather parameters with-on the population of *Tetranychus urticae* on okra crop during Summer 2023

Table 2: Correlation coefficients and regression equation of phytophagous mite with weather parameters on okra crop during crop period 2023

Correlation Factor	Temperature (°C)		Relative humidity (%)		Rainfall
	Maximum	Minimum	Maximum 700hr	Minimum 1400 hr	
Red spider mite	0.779**	0.699*	-0.375 ^{NS}	-0.068 ^{NS}	0.306 ^{NS}
Regression factor	Regression equation			Multiple R ² Value	R ² Value
Red Spider Mite	Y= -184.644 +X ₁ (1.129) + X ₂ (4.626) -X ₃ (1.333) -X ₄ (0.810) -X ₅ (0.107)			0.8751	0.7456

**Correlation is significant at the 0.01 level *Correlation is significant at the 0.05 level.

X₁- Maximum temperature, X₂- Minimum Temperature, X₃-Relative humidity 700hrs, X₄- Relative humidity 1400hrs, X₅- Rainfall,
R₂- Coefficient of Determination