

Contribution of PGPRs to plant growth

Abstract

PGPRs play an important role in maintaining soil equilibrium. This environment is a living medium made up of essentially heterotrophic microorganisms, and is a major reservoir of rhizobacteria. It contains a complex and varied microflora that plays essential roles for the soil ecosystem and the soil's higher organisms. In the rhizosphere, in terms of biomass and taxon diversity, PGPRs are by far the most abundant in the soil. The aim of this study is to characterise the rhizobacteria and arbuscular mycorrhizal fungi (AMF) that occur naturally in the rhizosphere of millet as a function of different soil fertility management practices.

1. Introduction

Rhizobacteria (PGPR, acronym for Plant Growth Promoting Rhizobacteria) are bacteria in the rhizosphere that are beneficial to plant growth and health. There are two main groups of PGPR: phyto-stimulators and phyto-protectants [1]. The term PGPR was first introduced at the end of the 1970s, when it was demonstrated by Kloepper and Schroth that strains of *Pseudomonas fluorescens* improved potato crop yields by up to 500% through the production of siderophores, iron chelators that deprive indigenous pathogenic bacteria of iron [1]. PGPRs are of major agronomic interest, as their use could make it possible to reduce the use of chemical fertilisers and pesticides [2]. These are bacteria that actively colonise plant roots while increasing their growth and yield [3]. PGPRs can be used in two different ways: phyto-stimulation (sometimes called bio-fertilisation), when the PGPR directly stimulates plant growth, and phyto-protection (also called bio-control), when it inhibits the development of phyto-pathogenic organisms [4]. Plant growth-promoting rhizobacteria (PGPR) are considered an alternative to the use of chemicals in agriculture [5]. These PGPR are often used as model rhizobacteria [6]. They make up a significant proportion (up to 10%) of the cultivable rhizosphere microflora [7]. The aim of this study is to analyse the results of research showing the significant influence of PGPRs on plant growth, thereby contributing to a better understanding of the nutrition of cultivated plants.

2. Plant Growth Promoting Rhizobacteria (PGPR)

Rhizobacteria are microorganisms that directly stimulate plant growth plant growth directly by increasing the uptake of nutrients from the soil nutrients from the soil, inducing and producing plant growth regulators and plant growth regulators and activating induced resistance mechanisms in plants. They indirectly stimulate plant growth through their antagonistic effect on harmful microflora, by transforming toxic metabolites. toxic metabolites. The establishment of the PGPR-plant association is essential for the expression of beneficial effects.

Expression of beneficial effects. Rhizobacteria are bacteria with the ability to the ability to colonise roots intensively. Non symbiotic bacteria that meet this definition belong to different genera and species species, the most extensively studied being : *Agrobacterium*

Comment [11]: italic letters

Comment [12]: why the same reference repeated?

radiobacter, *Azospirillum* spp, *Bacillus* spp, *Pseudomonas* spp [8]. According to [8], the PGPR/pant relationship improves nutrient uptake, the same authors also concluded that the application of bacterial inoculations considerably improves N, P, and K uptake. Certain PGPR strains of the genera *Pseudomonas*, *Bacillus*, *Paenibacillus*, *Rhodobacter* and *Azospirillum* have recently been described for their direct positive effect on plant growth and increased crop yield [9]. PGPRs can promote host plant growth through various mechanisms such as nitrogen nitrogen (N₂) fixation and solubilisation of trace elements such as phosphate (P) [10]. Basically, RMPs are defined by three intrinsic characteristics [11].

- (1) they must be able to colonise the root ;
- (2) they must survive and multiply in the microhabitats associated with the root surface, in competition with other microbiota;
- (3) they must promote plant growth [12].

2.1. Diversity of RMPs in the rhizosphere

In the rhizosphere, bacteria are by far the most abundant microorganisms in terms of both biomass and taxonomic diversity[13]. [14]estimated that one gram of natural forest soil contains nearly 1.5. 10¹⁰ bacteria. Generally speaking, it is now considered that one gram of soil contains several thousand species and that the abundance of these species can vary from 10⁸ to 10¹¹ cells per gram of soil [15]. PGPRs can be classified into two types according to their degree of association with root cells. They are divided into extracellular (ePGPR) and intracellular (iPGRP) [16, 17,18]. PGPRs reside in the rhizosphere (mainly the rhizoplane) or in the intracellular space of the root cortex. iPGPRs, on the other hand, mainly reside inside nodules. The bacterial genera belonging to extracellular PGPRs are *Azotobacter*, *Serratia*, *Azospirillum*, *Bacillus*, *Caulobacter*, *Chromobacterium*, *Agrobacterium*, *Erwinia*, *Flavobacterium*, *Arthrobacter*, *Micrococcus*, *Pseudomonas* and *Burkholderia*. In contrast, endophytic bacteria belonging to intracellular PGPRs include *Allorhizobium*, *Bradyrhizobium*, *Mesorhizobium*, and *Rhizobium*, as well as *Frankia* species, which can fix atmospheric nitrogen in actinorhizal plants [19].

2.2. Taxonomic diversity of rhizobacteria

PGPR colonise the rhizosphere using root exudates as nutrient substrates, but unlike other rhizospheric bacteria they in turn have a beneficial effect on the plant via a multitude of mechanisms (Figure 1). The enormous taxonomic and genetic diversity of rhizobacteria means that they are heavily involved in numerous environmental functions in the soil. Certain PGPRs are involved in plant health and growth, the most extensively studied being rhizobial and mycorrhizal symbioses. The soil microbial component also plays an active role in the biogeochemical cycles of sulphur, phosphorus, iron and nitrogen. As far as the latter is concerned, their involvement in atmospheric nitrogen fixation, ammonification, nitrification and denitrification processes is well established [20]. PGPRs are among the main players controlling the decomposition of organic matter. Due to their metabolic plasticity, these soil microorganisms are also involved in the degradation and immobilisation of pollutants (pesticides) brought in from agricultural or industrial sources. At soil level, the distribution of microorganisms is heterogeneous and is conditioned by the organisation of the soil (texture, structure, composition, etc.) according to [21,22]. The distribution of microorganisms in the soil is also highly dependent on nutrient resources and their spatiotemporal distribution.

Comment [I3]: italic letters why the same reference repeated again

Comment [I4]: italic letters

Comment [I5]: italic letters

Comment [I6]: repeated word

Comment [I7]: repeated word

Comment [I8]: what does this appropriation mean?

Comment [I9]: italic letters

Comment [I10]: italic letters

Comment [I11]: ?? what is the meaning of this speech?

Comment [I12]: please rewrite this paragraph

Consequently, the presence of a particular trophic niche will strongly structure the distribution of microorganisms in the soil [21,22]. [23]. Have classified RMPs into four sub-groups according to their modes of action:

- Bio-fertilisers (increase in the availability of nutrients to the plant);
- Phyto-stimulators (increase in plant growth, ability to produce phytohormones);
- Rhizoremediators (degradation of organic pollutants);
- bio-pesticides (disease control, production of fungicidal and antibiotic metabolites).

Cultivable micro-organisms, with a diversity of genera and species, belong mainly to the following three phyla: Proteobacteria, Firmicutes and Actinobacteria [20]. At present, many bacterial genera include PGPRs, revealing very diverse taxa [5].




Figure 1: Effects of rhizobacteria on root system architecture and root function [24].

3. Mechanisms of action of PGPR

The beneficial effects of PGPRs on plant growth result from different mechanisms exerted by rhizobacteria whose modes of action are direct or indirect, although the difference between the two is not always obvious.

Indirect mechanisms are generally those that occur outside the plant, whereas direct mechanisms are those that occur inside the plant and directly affect its metabolism. These mechanisms (Figure 2) may be active simultaneously or sequentially at different stages of plant growth:

1. Solubilisation of phosphates, nitrogen fixation and mineral nutrients, making these foods available to the plant;
2. The production of phytohormones such as 3-indoleacetic acid (IAA);
3. The repression of pathogenic soil microorganisms (through the production of hydrogen cyanide, siderophores, antibiotics, and/or competition for nutrients [25]. In addition, PGPRs can contribute to improving plant resistance to biotic and abiotic stresses (salinity, dryness and heavy metal toxicity) on the basis of their activities. [23] have classified PGPRs as biofertilisers (increasing the availability of nutrients to plants), phytostimulators (improving

Comment [I13]: repeated speech

plant growth, usually through the production of phytohormones), rhizoremediators (degrading organic pollutants) and biopesticides (controlling disease, mainly through the production of antibiotic and antifungal metabolites).




Figure 2: Overview of the two direct and indirect modes of RMPs [26].

Conclusion

This review has revealed the role played by PGPRs in the plant rhizosphere. The rhizosphere is home to various microorganisms that interfere with the plant. PGPRs promote the growth of host plants through mechanisms such as nitrogen fixation and the solubilisation of trace elements such as phosphate. The beneficial effects of PGPRs on plant growth result from different mechanisms exerted by rhizobacteria whose modes of action are direct or indirect, although the difference between the two is not always obvious. Certain PGPRs are involved in plant health and growth, the most widely studied of which are rhizobial and mycorrhizal symbioses. The soil microbial component also plays an active role in the biogeochemical cycles of sulphur, phosphorus, iron and nitrogen. As far as nitrogen is concerned, the strong involvement in the fixation of atmospheric nitrogen, ammonification, nitrification and denitrification processes no longer needs to be demonstrated.

Comment [114]:

UNDER PEER REVIEW

Références

1. García-Arenal F, McDonald B (2003) An analysis of the durability of resistance of plant to viruses. *Phytopathology* 93:941-952
1. Garcia J-L., Roger PA (2001), Introduction à la microbiologie du sol. Polycopié de cours (Université de Provence, Université de la Méditerranée, Ecole Supérieure d'Ingénieurs de Luminy. 191pp
1. García Lucas, J.A., Schloter, M., Durkaya, T., Hartmann, A. Gutierrez-Maero, F.J (2003). Colonisation des racines de poivrons par une souche de *Pseudomonas fluorescens* favorisant la croissance des plantes. *BiolFertil Soils* 37(6) : 381-385.
2. Morrissey, O. et D. Nelson (2002). The Role of the World Bank in the Transfer of Policy Knowledge on Trade Liberalisation ».politiques sur la libéralisation du commerce ». Document présenté à la conférence sur Political Economy of Policy Reform, Tulane University, 9-10 novembre. Disponible à l'adresse : www.tulane.edu/~dnelson/PEReform.htm
3. Wu, H., Hayes, M.J., Wilhite, D.A., et al. (2005) The Effect of the Length of Record on the Standardized Precipitation Index Calculation. *International Journal of Climatology*, 25, 505-520. <https://doi.org/10.1002/joc.1142>
4. Lucy, M., Reed, E., Glick, B.R., 2004. Application of free living plant-promoting rhizobacteria. *Antonie van Leeuwenhoek* 86, 1-25.
5. Kloepper, J. W. (1993). Plant-growth-promoting rhizobacteria as biological control agents, In : *Soil Microbial Ecology*, (Ed.) F. B. Jr, Metting. Marcel Dekker inc., N. Y. p.255-273.
6. Sorensen, P. W. (2001). Laboratory assessment of the role of a larval pheromone and natural stream odor in spawning stream localization by migratory sea lamprey (*Petromyzon marinus*). *Journal canadien des sciences halieutiques et aquatiques*, 58(12), 2374-2385.
7. Kragelund, B. B., Knudsen, J., Aplin, R. T., Poulsen, F. M., & Dobson, C. M. (1996). Probing the nature of noncovalent interactions by mass spectrometry. A study of protein- CoA ligand binding and assembly. *Journal of the American Chemical Society*, 118(36), 8646-8653.
8. Hallmann, J., Quadt-Hallmann, A., Rodriguez-Kibana, R., et Kloepper, J.W. 19976. Interactions between *Meloidogyne incognita* and endophytic bacteria in cotton and cucumber. *SoilBiol. Biochem.* In press.
10. ORHAN, M., SCOTT, D. (2001), « Why women enter into entrepreneurship : an explanatory model », *Women in Management Review*, vol. 16, n° 5, pp. 232-247.
11. Barea JM, Werner D, Azcón-Aguilar C, Azcón R. 2005. Interactions of arbuscular mycorrhiza and nitrogen fixing symbiosis in sustainable agriculture. In : Werner D, Newton WE, eds. *Agriculture, forestry, ecology and the environment*. Pays-Bas : Kluwer Academic Publishers.
12. Niranjana, S. R. et Hariprasad. P. 2014. Understanding the Mechanism Involved in PGPR-Mediated Growth Promotion and Suppression of Biotic and Abiotic Stress in Plants. In : *Future Challenges in Crop Protection Against Fungal Pathogens*, Fungal Biology, Springer 2014, pp 59-108.
13. Schmidt, F. L. et Hunter, J. E. (2002). Bias in standardized educational and employment tests as justification for racial preferences in affirmative action programs. In K. T. Leicht (Ed.), *The future of affirmative action* (Vol. 17, pp. 285-302). Stanford, CT : JAI (PDF) Le

rôle de l'aptitude cognitive générale et de la performance professionnelle : Pourquoi il ne peut y avoir de débat.

14. Torsvik V, Sorheim R, Goksoyr J.. Total bacterial diversity in soil and sediment communities-a review. *J Ind Microbiol* 17 : 170-178

15. Roesch, L., Fulthorpe, R., Riva, A. et al. Pyrosequencing enumerates and contrasts soil microbial diversity. *ISME J* 1, 283-290 (2007). <https://doi.org/10.1038/ismej.2007.53>

16, 17, 18 Gray, N. S., Brown, A. S., MacCulloch, M. J., Smith, J. et Snowden, R. J. (2005). An Implicit Test of the Associations Between Children and Sex in Pedophiles. *Journal of Abnormal Psychology*, 114(2), 304-308. <https://doi.org/10.1037/0021-843X.114.2.304>

19. Gouda, A. I., Toko, I. I., Salami, S. D., Richert, M., Scippo, M. L., Kestemont, P., & Schiffers, B. (2018). Pratiques phytosanitaires et niveau d'exposition aux pesticides des producteurs de coton du nord du Bénin. *Cahiers Agricultures*, 27(6), 65002.

20. Hugenholtz, P., & Huber, T. (2002). Chimeric 16S rDNA sequences of diverse origin are accumulating in the public databases. *International journal of systematic and evolutionary microbiology*, 53(1), 289-293.

21, 22 Ranjard, L. et Richaume, A. (2001). Quantitative and qualitative microscale distribution of bacteria in soil. *Research in microbiology*, 152(8), 707-716.

23. Somers, D. E. (2004). La formation d'un complexe SCFZTL est nécessaire pour une bonne régulation du rythme circadien. *The Plant Journal*, 40(2), 291-301.

24. Belotti, F., Daidone, S., Ilardi, G., & Atella, V. (2013). Stochastic frontier analysis using Stata. *The Stata Journal*, 13(4), 719-758.

25. Understanding the Mechanism Involved in PGPR-Mediated Growth Promotion and Suppression of Biotic and Abiotic Stress in Plants. In : *Future Challenges in Crop Protection Against Fungal Pathogens*, Fungal Biology, Springer 2014, pp 59-108.

26. Basu, R. et Ferreira, J. (2021). Mobilité durable dans la région métropolitaine de Boston dominée par l'automobile : Challenges and opportunities post-COVID-19. *Transport Policy*, 103, 197-210.

UNDER PEER REVIEW

