Exploring Trait Interactions for Yield Improvement in Linseed(*Linumusitatissimum*L.): A Correlation and Path Analysis Approach

Abstract

Linseed (*Linumusitatissimum*L.)is a self-pollinated (2n = 20)annual Rabi season crop mainly cultivated for flax fiber and oil for its commercial use. The present investigation was carried out to study association analysis among the yield and yield attributes using 75 linseed genotypes including 10 checks in randomized blocked design with three replications at Oilseeds Research Farm, Kalyanpur, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, Uttar Pradesh, (India) during Rabi season 2019-20. The correlation analysis revealed that the traits number of secondary branches per plant followed the number of primary branches per plant and number of capsules per plant at both the genotypic and phenotypic levels.The results of the path coefficient analysis revealed that traits such as the number of secondary branches per plant, number of capsules per plant, number of seeds per capsule, and plant height exhibited a strong and positive direct effect on seed yield per plant at both the genotypic and phenotypic levels.The seed yield can be improved by selecting the traits viz., number of primary branches per plant, number of secondary branches, number of capsules per plant, and plant height.

Keywords: Linseed, Seed yield, Correlation, and Path analysis.

Introduction:

Linseed (*Linumusitatissimum*L.) is an annual, self-pollinated oilseed crop (Yadava*et al.*, 2012) primarily cultivated during the rabi season. Belonging to the genus "Linum"in the familyLinaceae, ~~linseed~~ originated in Southwest Asia, particularly in India (Richharia, 1952). It is grown both as an oilseed and a fibre crop, with flax fibre derived from a taller plant variant with greater straw length (Pavelek *et al.*, 2020). The oil content of linseed ~~is~~ varied from 36 - 48 % (Patel *et al.*, 2023). ~~Due to its drying properties, linseed oil is highly valued for producing premium-grade paints, soap, linoleum putty, varnishes, and medicinal products (Goyal *et al.*, 2014).~~In India, it is mainly cultivated in the states of Chattisgarh, Jharkhand,

Comment [H1]: not clear from the manuscript

Comment [H2]: Use single quotation mark

Comment [H3]: Comma should be replaced with full stop as your sentence is complete here

Comment [H4]: Belonging to the genus 'Linum'in the family Linaceae, originated in Southwest Asia, particularly in India.

Comment [H5]: What is the relevance of writing this sentence?
If not better remove it

UNDER PEER REVIEW

Madhya Pradesh, Maharashtra, and Odisha. The area, production, and productivity of linseed in India is2.39 lakh hectares, 1.67 lakh tones and 698 kg/hectares (Indiastat 2023) respectively. The main reason for the low productivity of linseed is ~~due to it is grown in~~ input starved conditions, marginal and sub-marginal lands which ~~are~~ risk-prone and poor fertility status. More than 63% of the area is under rain-fed, an adaptation of primitive sowing method relay (utera) cropping (Srivastava, 2009) and lack of awareness/dissemination of better agro techniques among the farmers and perpetual scarcity of basic input viz, improved varieties, irrigation, fertilizers, weed control,etc (Chandrawati*et al*., 2016).The present status of linseed production could be increased 2-3fold through the adoption of improved varieties coupled with recommended production and protection technologies. The idea of the factors which are responsible for higher yield is a complicated problem. Because yield is a very complex character and an interactive effect/ multiplication of different traits and is highly influenced by environmental factors. Therefore, to achieve a high yield level, the breeder requires simplifying this complex situation by handling yield components that have a negative association with each other. In order to achieve this rational approach, it is vital to obtain information on the nature and extent of the correlation between different yield components and to resolve quality/quantity and how they contribute to yield. The current research was carried out to study the association analysis between yield and yield attributes.

Comment [H6]: Crosscheck the data

Comment [H7]: Have

Comment [H8]: Which area? Cultivated or else

Comment [H9]: Rephrase the sentence in meaningful way

Materials and methods

Comment [H10]: Capitalized the word

The experiment was conducted to estimate the association of traits for seed yield per plant using 75 linseed (*Linumusitatissimum*L.) genotypes including 10 checks in Randomized Block Design with three replications atOilseeds Research Farm, Kalyanpur, Chandra Shekhar Azad University of Agriculture and Technology, Kanpur, Uttar Pradesh, (India) during Rabi season 2019-20. Sowing was done in a 2-row plot of 5-meter length. The spacing is 30 cm × 7 cm distance was maintained by proper thinning. A healthy and good crop is raised by ~~the proper~~ recommended package of agronomic practices. Equally competitive five plants were tagged for recording the observations on traits viz., days to 50% flowering, number of primary branches per plant, number of secondary branches per plant, plant height, number of capsules per plant, number of seeds per capsule, days to maturity, oil content, thousand seed weight and seed yield per plant. The oil content was estimated by weight of 50 g seeds taken after threshing and drying on a Nuclear Magnetic Resonance (NMR) technique. Both the genotypic correlation coefficient and phenotypic

Comment [H11]: adopting

UNDER PEER REVIEW

correlation coefficient were estimated from the formula given by Aljibouri*et al.* (1958).The path analysis was performed by a process as described by Dewey and Lu, 1959.

Comment [H12]: Correct the name, mentioned in reference

Comment [H13]: (Aljibouri*et al.*, 1958)

Comment [H14]: it should be like this (Dewey and Lu, 1959)

Comment [H15]: Follow the authors guideline for the citation

Results and Discussion

Correlation analysis

The observed genotypic values of the correlation coefficient were higher than the phenotypic correlation coefficient value of most of the characters, which significantly impacts seed yield (Table 1).Indirect selection is effective under the traits, which is low heritability (Singh *et al.*, 2024). This correlation studies guides the breeder to select the traits which is responsible for yield and quality improvement in linseed. Seed yield per plant was high and significantly correlated with the number of secondary branches per plant (G = 0.567), number of primary branches per plant (G = 0.498)and number of capsules per plant (G = 0.438) at the genotypic level. At the phenotypic level, seed yield per plant showed positive and significantly associated number of secondary branches per plant (P = 0.544), number of capsules per plant (P = 0.4356) and number of primary branches per plant (P = 0.426).Similar type results were reported by Tariq *et al.* (2014), Patel *et al.* (2023), for capsules per plant and primary branches per plant;Singh*et al.* (2024) andJain *et al.* (2011)for primary branches per plant, secondary branches per plant and capsules per plant.The results revealed that the seed yield per plant was positively correlated with thousand seed weight (G = 0.0934, P = 0.0897), number of seeds per capsule (G = 0.0584, P = 0.0494), days to 50 per cent flowering (G = 0.0318, P = 0.0315) and oil content (G = 0.0282, P = 0.0288). These results agree with those of Muhammad Akbar *et al.* (2001) and Ibrar*et al.* (2016).The interrelationship were positive and significant among yield component traits like days to 50 per cent flowering with days to maturity (G = 0.638, P = 0.616), plant height (G = 0.252, P = 0.247), number of capsules per plant (G = 0.158, P = 0.155) and number of primary branches per plant (G = 0.156, P = 0.141), number of primary branches with number of secondary branches per plant (G = 0.798, P = 0.709), number of capsules per plant (G = 0.457, P = 0.383) and days to maturity (G = 0.182, P = 0.1567), number of secondary branches per plant with number of capsules per plant (G = 0.550, P = 0.531), plant height with days to maturity (G = 0.264, P = 0.257), oil content (G = 0.178, P = 0.176) and thousand seed weight (G = 0.131, P = 0.129), oil content with test weight (G = 0.511, P = 0.5023)at both the genotypic and phenotypic levels, except for traits viz., number of primary branches with days to 50 per cent flowering and plant height with thousand seed weight which is at genotypic

Comment [H16]: Significant associations with the

Comment [H17]: (Tariq *et al.* (2014); Patel *et al.* 2023)
Follow the journal rules and rectify such errors everywhere

Comment [H18]: follow the guideline for the authors and rectify such errors

Comment [H19]: Follow author citation rule

Comment [H20]: attributing

UNDER PEER REVIEW

level only.These type results are in agreement with Paul *et al.* (2020), Kumar and Paul (2016), Kasana*et al.* (2018), Chaudhary *et al.* (2016), Sharma *et al.* (2016) and Ankit *et al.* (2019). The interrelationships were negative and significant among the yield-contributing traits like days to 50 per cent flowering with oil content (G = -0.343, P = -0.336)and thousand seed weight (G = -0.202, P = -0.198), plant height with number of capsules per plant (G = -0.296, P = -0.293), number of capsules per plant with oil content (G = -0.278, P = -0.275) at both the genotypic and phenotypic levels (Patial *et al.*, 2018). The number of seeds per capsule was negative and significant at genotypic and negative at phenotypic level correlated with days to maturity (G = -0.261, P = -0.192), number of primary branches (G = -0.233, P = -0.129), oil content (G = -0.199, P = -0.1622), number of secondary branches per plant (G = -0.177, P = -0.133) and thousand seed weight (G = -0.1578, P = -0.128). Such type of results were matched with Patel *et al.*, 2023.

Path analysis

The direct effect of phenotypic path coefficient value is higher for most of the characters compared to genotypic path coefficient value, it indicates that there is a significant genotype x environment interaction for traits expressed.The findings of path analysis (presented in Table 2 and figures 1 &2)revealed that the number of secondary branches per plant (G = 0.3344, P = 3556), number of capsules per plant (G = 0.2014, P = 0.2535), number of primary branches (G = 0.1928, P = 0.1004) andnumber of seeds per capsule (G = 0.1772, P = 0.1075)and plant height (G = 0.0998, P = 0.1045) exhibited positive and direct effect on seed yield per plant at both genotypic and phenotypic levels.Plant height (P = 0.1045) showed positive direct effect on seed yield per plant at phenotypic level. These results are matched with Ronika *et al.* (2020), Ibrar*et al.*(2016), Naik and Setapathy (2002) andRahimi*et al.* (2011), Gaurah and Rao (2011) and Yadav (2001)for the number of primary branches per plant, capsules per plant and seeds per capsule.The number of secondary branches per plant showed a positive indirect effect on seed yield per plant through the number of primary branches per plant (G = 0.2669, P = 0.2552) and number of capsules per plant (G = 0.1840, P = 0.1887) at both genotypic and phenotypic levels. Similar results were reported by Gudmewad*et al.* (2016). On the other hand, the number of primary branches per plant (G = 0.1539, P = 0.0712) and number of capsules per plant (G = 0.1108, P = 0.1345) exhibited indirect effect on seed yield per plant via number of secondary branches per plant at both the genotypic and phenotypic levels (Gudmewad*et al.*,2016; Ghiday*et al.* (2023).The remaining

Comment [H21]: Use multiplication symbol instead of alphabet 'X'

UNDER PEER REVIEW

characters exerted negligible direct and indirect contributions towards the seed yield per plant.

Conclusion

The current research concluded that the number of secondary branches, number of primary branches per plant, and number of capsules per plant demonstrated positive and significant associations with seed yield per plant at both genotypic and phenotypic levels. The results of the path coefficient analysis revealed that traits such as the number of secondary branches per plant, number of capsules per plant, number of seeds per capsule, and plant height exhibited a strong and positive direct effect on seed yield per plant at both the genotypic and phenotypic levels.The secondary branches per plant showed positive and indirect effect on seed yield per plant via number of primary branches per plant and number of capsules per plant. Accordingly, the seed yield can be improved by selecting the traits viz., number of primary branches per plant, number of secondary branches, number of capsules per plant, and plant height.

UNDER PEER REVIEW

Table 1: Estimates of genotypic (G) and phenotypic (P) correlation among different traits in Linseed

Sr No.	Characters		D50F	NPBPP	NSBPP	PH (cm)	NCPP	NSPC	DM	OC (%)	SW (g)	SYPP (g)
1.	D50F	r_g	**1.000**	0.156*	0.034	0.252**	0.158*	0.091	0.638**	-0.343**	-0.202**	0.0318
		r_p	**1.000**	0.141	0.038	0.247**	0.155*	0.071	0.616**	-0.336**	-0.198**	0.0315
2.	NPBPP	r_g		**1.000**	0.798**	-0.031	0.457**	-0.233**	0.182**	0.022	-0.034	0.498**
		r_p		**1.000**	0.709**	-0.031	0.383**	-0.129	0.1567*	0.0178	-0.0281	0.426**
3.	NSBPP	r_g			**1.000**	-0.045	0.550**	-0.177**	0.053	-0.001	0.079	0.567**
		r_p			**1.000**	-0.045	0.531**	-0.133	0.047	-0.001	0.070	0.544**
4.	PH (cm)	r_g				**1.000**	-0.296**	-0.085	0.264**	0.178**	0.131*	0.008
		r_p				**1.000**	-0.293**	-0.062	0.257**	0.176**	0.129	0.011
5.	NCPP	r_g					**1.000**	0.1167	0.0268	-0.278**	-0.068	0.438**
		r_p					**1.000**	0.096	0.0267	-0.275**	-0.066	0.4356**
6.	NSPC	r_g						**1.000**	-0.261**	-0.199**	-0.1578*	0.0584
		r_p						**1.000**	-0.192	-0.1622	-0.128	0.0494
7.	DM	r_g							**1.000**	-0.1156	-0.028	0.0068
		r_p							**1.000**	-0.1106	-0.0252	0.0057
8.	OC (%)	r_g								**1.000**	0.511**	0.0282
		r_p								**1.000**	0.5023**	0.0288
9.	SW (g)	r_g									**1.000**	0.0934
		r_p									**1.000**	0.0897
10.	SYP (g)	r_g										**1.0000**
		r_p										**1.0000**

***Indicates significant at 5% level, ** indicates significance at 1% level**

D50F: Days to 50 % flowering; NPBPP: Number of primary branches per plant; NSBPP: Number of secondary branches per plant; PH: Plant height; NCPP: Number of capsules per plant; NSPC: Number of seeds per capsule; DM: Days to maturity; OC: Oil content; SW: Thousand seed weight; SYP: Seed yield per plant.

Comment [H22]: Indicate the legend at the table

Comment [H23]: What is this please mention below the table

UNDER PEER REVIEW

Table 2: Estimates of direct and indirect effects for various traits studied towards seed yield per plant at genotypic (G) and phenotypic (P) levels in linseed

Sr No.	Characters		D50F	NPBPP	NSBPP	PH (cm)	NCPP	NSPC	DM	OC (%)	SW (g)	SYPP (g)
1.	D50F	G	**-0.0665**	-0.0106	-0.0022	-0.0168	-0.0105	-0.0060	-0.0425	0.0228	0.0134	0.0318
		P	**-0.0319**	-0.0045	-0.0012	-0.0079	-0.0049	-0.0023	-0.0197	0.0107	0.0063	0.0315
2.	NPBPP	G	0.0308	**0.1928**	0.1539	-0.0060	0.0882	-0.0449	0.0351	0.0042	-0.0066	0.4987**
		P	0.0141	**0.1004**	0.0712	-0.0031	0.0385	-0.0130	0.0157	0.0018	-0.0028	0.4259**
3.	NSBPP	G	0.0113	0.2669	**0.3344**	-0.0150	0.1840	-0.0591	0.0175	-0.0003	0.0266	0.5673**
		P	0.0137	0.2522	**0.3556**	-0.0161	0.1887	-0.0473	0.0167	-0.0001	0.0249	0.5439**
4.	PH (cm)	G	0.0252	-0.0031	-0.0045	**0.0998**	-0.0295	-0.0084	0.0263	0.0178	0.0131	0.0086
		P	0.0259	-0.0032	-0.0047	**0.1045**	-0.0306	-0.0065	0.0269	0.0185	0.0136	0.0102
5.	NCPP	G	0.0318	0.0921	0.1108	-0.0596	**0.2014**	0.0235	0.0054	-0.0560	-0.0138	0.4382**
		P	0.0392	0.0972	0.1345	-0.0742	**0.2535**	0.0244	0.0068	-0.0697	-0.0168	0.4356**
6.	NSPC	G	0.0160	-0.0412	-0.0313	-0.0150	0.02207	**0.1772**	-0.0462	-0.0354	-0.0280	0.0584
		P	0.0077	-0.0139	-0.0143	-0.0067	0.0103	**0.1075**	-0.0206	-0.0174	-0.0138	0.0494
7.	DM	G	0.0115	0.0033	0.0009	0.0047	0.0005	-0.0047	**0.0179**	-0.0021	-0.0005	0.0068
		P	-0.0076	-0.0019	-0.0006	-0.0032	-0.0003	0.0024	**-0.0124**	0.0014	0.0003	0.0057
8.	OC (%)	G	-0.0146	0.0009	0.0000	0.0076	-0.0119	-0.0085	-0.0049	**0.0427**	0.018	0.0282
		P	-0.0200	0.0011	0.0000	0.0105	-0.0164	-0.0097	-0.0066	**0.0595**	0.0299	0.0288
9.	SW (g)	G	-0.0136	-0.0023	0.0054	0.0088	-0.0046	-0.0106	-0.0019	0.0344	**0.0674**	0.0934
		P	-0.0095	-0.0014	0.0034	0.0062	-0.0032	-0.0062	-0.0012	0.0299	**0.0482**	0.0897

Residual (G): 0.7805, Residual (P): 0.8013 **Dark figure denotes direct effect**

Comment [H24]: What is indicate

Comment [H25]: Give the detail at the end of table or on the body of manuscript

UNDER PEER REVIEW



Fig. 1. Direct and indirect effects of yield components on seed yield per plant at the genotypic level.



Fig. 2. Direct and indirect effects of yield components on seed yield per plant at the phenotypic level.

References

1. Al-Jibouri HA, Miller PA, Robinson HF. Genotypic and environmental variance and covariance in an upland cotton cross of inter-specific origin. Agron J. 1958; 50:633-637.
2. Ankit KS, Singh SP, Singh VK, Singh A, Tawari A. Estimates indirect selection parameters through correlation and path analysis in linseed (*Linumusitatissimum*L.). *International Journal of Chemical Studies*. 2019; *7*(1):2461-2465.
3. Chandrawati, Singh N, Kumar R, Kumar S, Yadav HK. Genetic variability and interrelationship among morphological and yield traits in Linseed (*Linumusitatissimum*L.). Genetika. 2016; 48 (3):881–892.
4. Chaudhary M, Rahul VP, Singh V, Chauhan MP. Correlation and path coefficient analysis for yield and yield-related traits in linseed (*Linumusitatissimum*L.). International Quarterly Journal of Life Sciences. 2016; 11(2): 939-942.
5. Dewey DR, Lu. K. H. A correlation and path coefficient analysis of components of of rested grass. Agronomy Journal. 1959; 51:515-518.
6. Gaurah, Rao SS. Analysis of yield Characters in linseed. Research Journal of Agricultural Sciences. 2011; 2 (2):258-260.
7. Ghiday T, Mohamed W, Tsehaye Y, Wakjira A, Daba C, Dissasa T. 2023. Association and Path Analyses among Morpho-agronomic Traits, Fatty Acids Contents and Oil Yield in Linseed (*Linumussitatisimum* L.) Genotypes in Ethiopia.
8. Goyal A, Sharma V, Upadhyay N, Gill S, Sihag M. Flax and flaxseed oil: an ancient medicine & modern functional food. J. Food Sci. Technol. 2014; 51:1633–1653.
9. Gudmewad R, Misal A, Dhutmal R, Pole S. Correlation and path analysis studies in linseed. *Annals of Plant and Soil Research*. 2016; 18(2):123-126.
10. Ibrar D, Ahmad R, Mirza MY, Mahmood T, Khan MA, Iqbal M.S. Correlation and Path analysis for yield and yield components in Linseed (*Linumusitatissimum*L.). Journal of Agricultural Research. 2016; 54(2):153-159.
11. Indiastat 2023. Selected-State-wise-Area-Production and Productivity of Linseed in India (2022-23).https://www.indiastat.com/table/agriculture/selected-state-wise-area-production-productivity-l/1456968#. Accessed on 12th September 2024.
12. Jain RJ. Correlation study of flowering performance and flowering pattern with the yield in (*Linumusitatissimum*L.). African J Plant Sci. 2011;5(3):146-151.

Comment [H26]: it should be at the end of tile of the paper in reference

UNDER PEER REVIEW

13. Kasana RK, Singh PK, Tomar A, Mohan S, Kumar S. Selection parameters (heritability, genetic advance, correlation and path coefficient) analysis in linseed (*Linumusitatissimum*L.). The Pharma Innovation Journal. 2018; 7(6):16-19.
14. Kumar N, Paul S. Selection criteria of linseed genotypes for seed yield traits through correlation, path coefficient and principal component analysis. The Journal of Animal and Plant Sciences. 2016; 26:1688-1695.
15. Muhammad Akbar, Norul Islam khan and Khalid Mahmood Shabir. Correlation and Path coefficient studies in linseed. *Journal of Biological Sciences*. 2001; 1(6):446- 447.
16. Naik BS, Satapathy PC. Selection strategy for improvement of seed yield in late sown linseed. Research on Crops. 2002; 3(3):599-605.
17. Patel MD, Patel PC, Donga AR, Soni NV, Vaghela GK, Shah S. K. Genetic Architecture and Association Analysis of Yield and Related Attributes in *Linumusitatissimum*(L.) *Frontiers in Crop Improvement.*2023; 11: 430-438.
18. Patial R, Paul S, Sharma D. Correlation and path coefficient analysis for improvement of seed yield in linseed (*Linumusitatissimum*L.). *International Journal of Current Microbiology and Applied Sciences*. 2018; *7*(3):1853-1860.
19. Paul S, Satasiya P, Kumar A. Genetic variability, correlation and path coefficient analysis of introduced genotypes of linseed (*Linumusitatissimum*L.) in mid-hills of North-West Himalayas. Journal of Pharmacognosy and Phytochemistry. 2020; 9(1):1189-1192.
20. Pavelek M, Vrobova-Prokopova M, Ondrackova E, Ludvikova M, Griga M. 2020. Developments in fibrous flax and linseed breeding and cultivation. In: Handbook of Natural Fibres, second ed. 19. Woodhead Publishing, Sawston, UK, pp. 1–95.
21. Rahimi MM, Zarei MA, Arminian A. Selection criteria of flax (*Linumusitatissimum*L.) for seed yield, yield components and biochemical compositions under various planting dates and nitrogen. African Journal of Agricultural Research. 2011; 6(13): 3167-3175. 29.
22. Richharia, R.H. 1952. Linseed. The Indian Central Oilseeds Committee, Hyderabad, India. 155.
23. Sharma D, Paul S, Patial R. Correlation and path coefficient analysis of seed yield and yield-related traits of linseed (*Linumusitatissimum*L.) in mid-hills of north-west Himalayas. International Quarterly Journal of Life Sciences. 2016; 11(4): 3049-3053.
24. Singh SP, Aware SA, Tiwari A, Sindha SB, Chellem SR, Singh DV. Estimates Indirect Selection Parameters Through Correlation and Path Analysis in Linseed

Comment [H27]: rectify it and find such errors

UNDER PEER REVIEW

(*Linumusitatissimum* L.). Plant Cell Biotechnology and Molecular Biology. 2024;25(3-4):54-61.

25. Srivastava RL. 2009. Research and development strategies for Linseed in India. In: National Symposium on Vegetable Oils Scenario: Approaches to Meet the Growing Demands, Jan. 29-31, 2009, Indian Society of Oilseeds Research, Directorate of Oilseeds Research, Rajendra Nagar, Hyderabad.
26. Tariq MA, Hussain T, Ahmad I, Saghir M, Batool M, Safdar M, Tariq M. Association analysis in linseed (*Linumusitatissimum*L.). *Journal of Biology, Agriculture and Healthcare*. 2014; *4*(6):60-62.
27. Yadav RK. Association studies over locations in linseed, Progressive Agriculture. 2001; 1(1):11-15.
28. Yadava DK, Vasudev S, Singh N, Mohapatra T, Prabhu KV Breeding major oil crop: present status and future research needs. Technol. Innov. Major World Oil Crops Breed. 2012;1:17–51.

Comment [H28]: rectify

UNDER PEER REVIEW