

**Dynamics of legume cropping and agro-ecological transition in the cotton savannah zones of West Africa: the case of Koumbia district of Burkina Faso**

**Abstract:**

**Aims:** This paper shows the contribution of participatory research to upgrade the integration of legumes into cropping systems.

**Place and Duration of Study:** The investigation is based on survey data collected in 2008 and 2011 (two reference years) and 2021 and 2022 (current situation) in the village of Koumbia in western Burkina Faso.

**Methodology:** Data was collected through surveys and field measurements. The surveys were conducted with a sample of fifty (50) volunteer farmers. In 2008, 2011 and 2022, the same questionnaire was administered to all 50 farmers. Field measurements with GPS (Global Positioning System) were conducted on crop area.

**Results:** The results show that the proportion of area under legumes crops increased by 77,38% after 15 years. Other legumes crops (*Glycine max* and *Mucuna sp.*) were introduced in addition to those previously grown (groundnuts and cowpeas) by farmers. Surface cropped in sorghum (35.36%) and millet (82.79%) declined while cotton and maize remain the main crops in the zone. The results on farmers' perception show that they prefer single cropping of legume crops as their insertion modality in the existing farming systems compared to their association with cereals.

**Conclusion:** The study shows that the role of legumes is increasing in the cropping system, and represents an opportunity to succeed in the agro-ecological transition, even though the intercropping of cereal/legume, which is put forward by the theoretical models of the agro-ecological transition, is still little practiced. The structuration of the legume sector can be a condition for achieving it.

**Keys words:** perception, single cropping, service plant, cropping system, Burkina Faso.

## 1. Introduction

Agro-ecology (AE) is optimizing interactions between plants, animals, humans and the environment while addressing social aspects of food system such as sustainability and equity (FAO, 2018). With regard to their triple role of soil fertility improvement, food and forage crops, legumes crops represent an opportunity for agro-ecologic transition in West Africa. Ranged as service plants likely to provide various ecosystem services, legumes prevent soils from physical degradation while improving their nutrient content through fixing atmospheric nitrogen. Indeed, *Mucuna sp.* was shown to reduced runoff by 30-25% and erosion by 25% in five years (Azontondé, 1993). Zougmore et al. (2000) reported that the association of cowpea with sorghum, led to the reduction of runoff in the range of 20 to 30% compared with single cropping of sorghum, and 5 to 10% compared to cowpea in single cultivation. As consequences, they found reduction in soil erosion in range of 45% to 80% according to the association method of leguminous plants to cereals compared to the cereals copped alone. Gebhard et al. (2013) reported that the nitrogen accumulated by legumes was mainly from symbiotic fixation while ranging it from few kg to 150 kg N/ha in three months of vegetation, and varies significantly according to the legumes species. Toé et al. (2022) obtained fodder yields of 1302 kg/ha for K VX-745-11P cowpea and 3231 kg/ha for IS-5874 cowpea in single cropping, compared to 3261 kg/ha for sorghum. The same authors ranged total nitrogen contained in cowpea fodder from 19.2 and 20.8% respectively for K VX 745 11P and IS-5874; and 7% for sorghum. Legumes play high socio-economic roles in the investigation areas; as reported by Koné et al. (2019); legumes play an important socio-economic role. They are consumed alone and also use in the preparation of daily dishes. Therefore, legumes crops production do not only generate significant income, but also offers households a varied range of food products.

In terms of human consumption, Koné et al. (2019) report that groundnuts are consumed in several forms (roasted, boiled, processed into oil), and that oilcake is processed into patties. For Koné et al. (2019), peanut production not only generates significant income, but also offers households a varied range of food products, most of which are used to accompany meals.

Despite these agronomic advantages and their economic and dietary importance, many challenges are been met for integrating legumes plants into the cropping systems. These challenges are both technical (phosphorus requirements and storage difficulties) and socio-economic, such as the low nutritional value of certain legumes like mucuna and the lack of structured outlets for other legumes like soya and cowpea (Carsky et al., 2003).

In western Burkina Faso, legumes plants such as groundnuts and cowpeas, occupied only a marginal share (5%) of crop rotations in 2008, are considered secondary crops compared to the main crops such as cotton, maize and sorghum. Since 2008, research has been carried out in this area in a number of areas, including the insertion of legumes into the cropping system (Coulibaly et al., 2012a, Barro et al., 2016;


Ouattara et al., 2016, Perinelle et al., 2021). Coulibaly et al. (2012a) worked on the co-design of technical itineraries for single cropping of cowpea and mucuna. Barro et al. (2016) worked on maize/ cowpea association options that would reduce competition between two crops in association for better agronomic and economic performance. Ouattara et al. (2016) analyzed the large-scale effects of maize-cowpea association for intensification and optimization of agricultural production in agropastoral systems in the commune of Koumbia. Perinelle et al. (2021) explored new ways of reintroducing legumes by tracking innovations in legume cropping practices in the commune of Boni (Tuy province, Burkina Faso). For addressing climate change, volatile prices for agricultural products and inputs, and the emergence of new markets, farmers are tending crops diversification on their own initiative.

With the current situation, the question raised herein is to know if legume integration in cropping systems have been driving by research activities and farming contexts. This investigation was initiated in 2022 in order to respond to the question. The objective of this paper is show and explain that legume crops insertion trend un the cropping systems of the district of Koumbia in Burkina Faso after 15 years of farmers 'fields research activities

## **2. Material et methods**

### **2.1. Study area**

The study was carried out in the village of Koumbia (11° 14' 11" North; 3° 41' 47" West), located 67 km northeast of Bobo-Dioulasso in the rural district of Koumbia in western Burkina Faso (Figure 1). The village of Koumbia is characterized by a Sudanian climate. At Koumbia, the rainfall is ranged from 775 mm to 1,202 mm, with an average of 966.24| over the last 10 years. In the village of Koumbia, cotton-based farming systems are the most common ones with Agri livestock integration (Vall et al., 2006). The cropping system is dominated by cotton, grown in rotation with maize.



**Figure 1 : Location of Koumbia**

## 2.2. Data collection

Data was collected through surveys and field measurements. The surveys were conducted with a sample of fifty (50) farmers that voluntarily participated in the activities of the Fertipartenaires project (funded by the European Union) from 2008 to 2012. In 2008, 2011 and 2022, the same questionnaire was administered to all 50 farmers. According to the simplified typology built by Vall et al. (2006), the farms surveyed were farmers (n=27), agro-pastoralists (n=17) and cattle breeders (n=6). The main areas covered by the questionnaire were related to the characteristics of these AEs for 2008, 2011 and 2022, and their cropping system (type of crop, area and crop production) for year n-1 of each survey year (2007, 2010, 2021). Data for 2008 and 2011 are considered as reference data, and data for 2021 and 2022 as current ones. In 2022, in addition to crop type and area, the questionnaire took into account farmers' perceptions of how legumes could be integrated into the existing cropping systems. To capture this perception, the best/worst scale method was used to collect data from respondents (Amadou, 2021). Thus, 15 modalities of legume insertion were identified for analysis. These modalities are the single cropping of mucuna, cowpea, groundnut, soybean and bambara groundnut, the association maize and mucuna, maize and cowpea, maize and groundnut, maize and soybean and maize-Bambara groundnut and the association sorghum and mucuna, sorghum and cowpea, sorghum- and groundnut, sorghum and soybean and

sorghum and Bambara groundnut. From these 15 modalities, each respondent was asked to choose 05 modalities that were the most important, and 05 modalities that were the least important in terms of agricultural production. Each respondent also gave the advantages and weaknesses of single cropping of legumes and cereal/legume associations.

Field measurements were conducted on crop area. Using a GPS (Global Positioning System) for crosschecking of the areas of crops declared by the respondents.

### 2.3. Data processing

Data collected was compiled using Excel, Office 2016; as well as calculations and drafting tables. Data on farm characteristics: areas and crop yields were subjected to analysis of variance (ANOVA) using Statistix 9.0 software. Means were separated at the 5% threshold using Tukey's test.

For farmers' perceptions of the different options of integrating legumes into existing cropping systems, the weight of each modality was determined according to the formula :

$$P_i = \frac{(B_i - C_i)}{aN} ; \text{Amadou (2021),}$$

with  $P_i$  = weight of modality  $i$ ,  $B_i$  = number of times modality  $i$  is cited as more important,  $C_i$  = number of times modality  $i$  is cited as less important,  $a$  = constant and  $N$  = size of sample.

For this study, the constant  $a=15/3 = 5$  with 15 the number of modalities and 3 the possibility that each modality is selected by the respondent as more important or less important or neither (indicating respondent neutrality). Survey population  $N = 50$ .

## 3. Results

### 3.1. Characteristics of the farms

The results show that from 2008 to 2022, the respondent's age increased from 41 to around 47 (Table 1). The number of household members and workers in the fields, the total area of fields and the number of cattle have not changed significantly after 15 years. With the exception of respondent age, the difference in farm characteristics surveyed between 2008 and 2022 is not significant ( $p>0.05$ ).

**Table 1: Variation of farm characteristics from 2008 to 2022**

|  | 2008                     | 2011                         | 2022                     | Pr(>F) | Significativity |
|--|--------------------------|------------------------------|--------------------------|--------|-----------------|
| Respondants' age                                     | 41.02 <sup>b</sup> ±1.68 | 43.08 <sup>ab</sup><br>±1.65 | 46.98 <sup>a</sup> ±1.76 | 0.045  | Yes             |
| Number of person                                     | 16.38±2.67               | 16.18±1.87                   | 15.22±1.05               | 0.907  | No              |
| Size of the human ressource in the housholds (olader | 7.64±0.82                | 8.68±1.17                    | 8.18±0.69                | 0.725  | No              |

|                        |            |            |            |       |    |  |
|------------------------|------------|------------|------------|-------|----|--|
| than +12yeras)         |            |            |            |       |    |  |
| Number of plots        | 2.00±0.13  | 1.88±0.14  | 2.10±0.24  | 0.688 | No |  |
| area of the plots (ha) | 7.77±1.15  | 10.17±1.21 | 8.11±1.03  | 0.271 | No |  |
| Number of oxen         | 14.50±2.87 | 14.28±2.48 | 14.30±2.74 | 0.998 | No |  |

### 3.2. Farming plots areas

For each investigated farmers the plots areas were determined using farmers 'declaration during the surveys and the GPS records as well. And, the GPS field measurements did not showed significant difference between the declared ones (Table 2). When ranking field size by crop, cotton and maize occupied first and second place respectively, with mean areas in the range of 4.75±0.75 ha (declaration) to 4.64±0.53 ha (GPS records) and from 2.80±0.38 ha (declaration) to 2.57±0.24 ha (GPS records). The difference between declared and measured area was higher for sesame (65.71%) than for the other crops.

**Table 2: Variation between cropped areas estimated by the surveys and field measurements**

| crops             | declaration |    | GPS records |    | Pr(>F) | Significativity |
|-------------------|-------------|----|-------------|----|--------|-----------------|
|                   | areas (ha)  | Nb | areas (ha)  | Nb |        |                 |
| Cotton            | 4.75±0.75   | 29 | 4.64±0.53   | 23 | 0.906  | No              |
| Maize             | 2.80±0.38   | 48 | 2.57±0.24   | 42 | 0.619  | No              |
| Sorghum           | 0.64±0.12   | 17 | 0.56±0.16   | 11 | 0.678  | No              |
| Millet            | 0.21±0.03   | 6  | 0.30±0.16   | 4  | 0.478  | No              |
| peanut            | 0.47±0.07   | 23 | 0.50±0.07   | 16 | 0.798  | No              |
| Cowpea            | 0.47±0.06   | 42 | 0.41±0.05   | 28 | 0.482  | No              |
| Rice              | 0.52±0.07   | 15 | 0.55±0.14   | 8  | 0.844  | No              |
| Bambara groundnut | 0.22±0.09   | 4  | 0.10±0.06   | 2  | 0.46   | No              |
| Sesame            | 1.05±0.65   | 8  | 0.36±0.24   | 4  | 0.49   | No              |
| soybean           | 1.78±0.22   | 40 | 1.91±0.26   | 31 | 0.699  | No              |
| Mucuna            | 0.25±0.05   | 7  | 0.31±0.11   | 4  | 0.548  | No              |

*Nb = number*

### 3.3. Cropping area dynamic from 2008 to 2022

From 2008 to 2022, the number of cotton, sorghum and millet farmers decreased (Table 3). On the other hand, the number of cowpea, soybean and rice farmers has increased. For maize, the number of farmers remained almost stable after 15 years.

Over the same period, cotton and maize occupy first and second rank respectively in terms of area (Table 3). Areas allocated to Sorghum and millet declined by 35.36% and 82.79% respectively after 15 years. In contrast, the area under sesame, soybean and rice increased by 31.25%, 57.52% and 23.81% respectively. Mucuna, which was absent in 2008 and 2010, was introduced with average areas of 0.33±0.08 ha and

0.25±0.05 ha in 2021 and 2022 respectively. Analyses of variance show that there is no significant difference ( $P>0.05$ ) between crop areas from 2008 to 2022.

Averaging cereals and pulses, we noticed that cotton takes first place in 2008 with an mean area of 4.51±0.86 ha, compared with 1.88±0.19 ha, 0.50±0.06 ha and 0.80±0.30 ha for cereals, pulses and sesame respectively (Table 3). The ranking remains the same in 2022. Area under pulses has increased significantly ( $P<0.001$ ) in 2021 (1.05±0.17 ha) and 2022 (0.88±0.09 ha), compared with 2008 (0.50±0.06 ha) and 2010 (0.42±0.04 ha). Also, cotton, cereal and sesame areas did not vary significantly ( $P>0.05$ ) after 15 years.

UNDER PEER REVIEW

**Table 3: Variation of cropped surface and number of crop users from 2008 to 2022**

| Crops                        | 2008                          |           | 2010                         |           | 2021                         |           | 2022                          |           | Pr(>F)           | Significativity |
|------------------------------|-------------------------------|-----------|------------------------------|-----------|------------------------------|-----------|-------------------------------|-----------|------------------|-----------------|
|                              | Area (ha)                     | Nb        | Area (ha)                    | Nb        | Area (ha)                    | Nb        | Area (ha)                     | Nb        |                  |                 |
| <b>Cotton</b>                | <b>4.51±0.86</b>              | <b>37</b> | <b>4.36±1.06</b>             | <b>27</b> | <b>4.48±0.66</b>             | <b>29</b> | <b>4.75±0.75</b>              | <b>29</b> | <b>0.978</b>     | No              |
| Maize                        | 2.88±0.42                     | 49        | 2.41±0.51                    | 50        | 2.95±0.42                    | 48        | 2.80±0.38                     | 48        | 0.818            | No              |
| Sorghum                      | 0.99±0.11                     | 37        | 1.28±0.17                    | 41        | 0.86±0.19                    | 15        | 0.64±0.12                     | 17        | 0.057            | No              |
| Millet                       | 1.22±0.33                     | 15        | 1.84±0.54                    | 17        | 0.25±0.06                    | 6         | 0.21±0.03                     | 6         | 0.081            | No              |
| Rice                         | 0.42±0.12                     | 6         | 0.73±0.17                    | 25        | 0.42±0.08                    | 16        | 0.52±0.07                     | 15        | 0.650            | No              |
| <b>Mean for cereal crops</b> | <b>1.88±0.19</b>              | <b>50</b> | <b>1.64±0.25</b>             | <b>50</b> | <b>2.11±0.31</b>             | <b>49</b> | <b>1.99±0.24</b>              | <b>49</b> | <b>0.584</b>     | No              |
| peanut                       | 0.49±0.06                     | 27        | 0.41±0.04                    | 19        | 0.49±0.06                    | 21        | 0.47±0.07                     | 23        | 0.788            | No              |
| cowpea                       | 0.50±0.06                     | 30        | 0.44±0.07                    | 17        | 0.49±0.10                    | 32        | 0.47±0.06                     | 42        | 0.967            | No              |
| Bambara groundnut            | 0.33±0.08                     | 5         | 0                            | 0         | 0.25                         | 1         | 0.22±0.09                     | 4         | 0.676            | No              |
| Soybean                      | 1.13±0.88                     | 2         | 0                            | 0         | 2.13±0.35                    | 31        | 1.78±0.22                     | 40        | 0.546            | No              |
| Mucuna                       | 0                             | 0         | 0                            | 0         | 0.33±0.08                    | 5         | 0.25±0.05                     | 7         | 0.248            | No              |
| <b>Mean for legum crops</b>  | <b>0.50<sup>bc</sup>±0.06</b> | <b>38</b> | <b>0.42<sup>c</sup>±0.04</b> | <b>29</b> | <b>1.05<sup>a</sup>±0.17</b> | <b>45</b> | <b>0.88<sup>ab</sup>±0.09</b> | <b>48</b> | <b>&lt;0.001</b> | <b>Yes</b>      |
| Sesame                       | 0.80±0.30                     | 10        | 0.85±0.13                    | 12        | 1.15±0.51                    | 5         | 1.05±0.65                     | 8         | 0.925            | No              |
| <b>Total</b>                 | <b>13.25</b>                  |           | <b>12.32</b>                 |           | <b>13.47</b>                 |           | <b>12.90</b>                  |           |                  |                 |

*Nb*

=

*number*

### 3.4. Global yields trends from 2007 to 2021

Overall, the surveys show that crop yields increased after 15 years (Table 4). Cotton yield increased from 845.11 kg/ha (2007) to 1256.28 kg/ha (2021). As for maize it was from 1146.02 kg/ha in 2007 to 2190.29 kg/ha in 2021. Sorghum yield, in contrast, fell by 23.32% after 15 years. Statistical analysis shows that the difference observed for cotton, maize and sesame is significant at the 5% threshold. This is not the case for rice, sorghum and millet.

For legumes, difference in yields was not significant at the 5%. However, there was an upward trend in yields of 22.48%, 27.62% and 442.18% respectively for groundnuts, cowpeas and soybeans after 15 years.

**Table 4: Variation of crop yields from 2007 to 2021**

| Crops               | 2007                         |    | 2010                         |    | 2021                         |    | Pr(>F)  | Significativity |
|---------------------|------------------------------|----|------------------------------|----|------------------------------|----|---------|-----------------|
|                     | Yields (kg/ha)               | Nb | Yields (kg/ha)               | Nb | Yields (kg/ha)               | Nb |         |                 |
| Cotton              | 845.11 <sup>b</sup> ±79.83   | 37 | 902.31 <sup>b</sup> ±69.19   | 27 | 1256.28 <sup>a</sup> ±73.91  | 29 | <0.001  | Oui             |
| Sesame              | 78.50 <sup>b</sup> ±14.45    | 8  | 215.83 <sup>ab</sup> ±60.52  | 12 | 433.33 <sup>a</sup> ±88.19   | 4  | 0.008   | Oui             |
| <i>Céréales</i>     |                              |    |                              |    |                              |    |         |                 |
| Maize               | 1146.02 <sup>c</sup> ±116.84 | 49 | 1739.39 <sup>b</sup> ±102.05 | 50 | 2190.29 <sup>a</sup> ±113.97 | 48 | <0.0001 | Oui             |
| Sorghum             | 851.85±214.12                | 36 | 638.78±83.43                 | 41 | 619.05±98.13                 | 14 | 0.527   | Non             |
| Millet              | 429.73±95.18                 | 15 | 353.80±56.04                 | 17 | 619.13±76.20                 | 5  | 0.212   | Non             |
| Rice                | 933.33±217.05                | 6  | 1960.53±639.02               | 25 | 2758.33±453.71               | 15 | 0.334   | Non             |
| <i>Légumineuses</i> |                              |    |                              |    |                              |    |         |                 |
| Peanut              | 1155.93±234.99               | 27 | 1288.89±219.46               | 18 | 1415.79±271.26               | 19 | 0.745   | Non             |
| Cowpea              | 408.89±65.80                 | 30 | 502.35±70.62                 | 17 | 521.81±45.80                 | 27 | 0.343   | Non             |
| Soybean             | 160.00±40.00                 | 2  | -                            | -  | 867.48±89.70                 | 31 | 0.0574  | Non             |

*Nb = Number*

### 3.5. Farmers' perceptions on the integration of legumes to the existing cropping systems

The overall results on farmers' perceptions show that single cropping of legume crops, cowpea (weight = 0.18), soybean (weight = 0.16), groundnut (weight = 0.15) and mucuna (weight = 0.10) are the legume insertion methods perceived by farmers as the most important in terms of agricultural production (Table 5). In contrast, associations of legumes plants with maize are perceived as the least important by these farmers, with weights of -0.17 for the maize and cowpea association, -0.15 for the maize and groundnut association, -0.14 for the maize and soybean association and -0.07 for the maize and Bambara groundnut association.

**Table 5: Proportion of farmers' responses on the most common and least common legume insertion methods according to agricultural production (n=50)**

| <b>Modalities for integrating legumes into cropping systems</b> | <b>Common practice</b> | <b>Lesscommon practice</b> | <b>Weigh</b> |
|---|------------------------|----------------------------|--------------|
| Single cropping of Mucuna                                       | 36                     | 11                         | 0.10         |
| Association Maize and Mucuna                                    | 16                     | 27                         | -0.04        |
| Association Sorghum and Mucuna                                  | 5                      | 13                         | -0.03        |
| Single cropping of Peanut                                       | 44                     | 6                          | 0.15         |
| Association Maize-Peanut  | 5                      | 43                         | -0.15        |
| Association Sorghum and Peanut                                  | 11                     | 19                         | -0.03        |
| Single cropping of Cowpea                                       | 48                     | 2                          | 0.18         |
| Association Maize and Niébé                                     | 1                      | 44                         | -0.17        |
| Association Sorghum and Niébé                                   | 9                      | 10                         | 0.00         |
| Single cropping of Soybean                                      | 41                     | 2                          | 0.16         |
| Association Maize and -Soybean                                  | 3                      | 37                         | -0.14        |
| Association Sorghum and Soybean                                 | 0                      | 13                         | -0.05        |
| Single cropping of Bambara groundnut                            | 22                     | 5                          | 0.07         |
| Association Maize and Bambara groundnut                         | 0                      | 17                         | -0.07        |
| Association Sorghum and Bambara groundnut                       | 0                      | 0                          | 0.00         |

Specifically for farmers, the single cropping of cowpea (weight = 0.20) is perceived as the first option for legume insertion (Table 6); while for agro-pastoralists, it's the single cropping of groundnut with a weight of 0.18. In the herder population, single croppings of mucuna, groundnut and cowpea were found to have the same weight (0.20). For the three (3) groups of farmers, the maize and cowpea combination was perceived as the least important modality. The maize and groundnut association and the one of maize and cowpea had the same with a weight of -0.18.

**Table 6: Proportion of farmers' responses according to farm typology on the most important and least important legume insertion methods according to agricultural production**

| Modalités d'insertion des légumineuses    | Farmers (n=27) |            |       | Agropastoralists (n=17) |            |       | Cattlebreeders (n=6) |            |       |
|---|----------------|------------|-------|-------------------------|------------|-------|----------------------|------------|-------|
|   | Common         | Lesscommon | Weigh | Common                  | Lesscommon | Weigh | Common               | Lesscommon | Weigh |
| Single cropping of Mucuna                 | 19             | 5          | 0.10  | 11                      | 6          | 0.06  | 6                    | 0          | 0.20  |
| Association of Maize and Mucuna           | 9              | 12         | -0.02 | 4                       | 12         | -0.09 | 3                    | 3          | 0.00  |
| Association of Sorghum and Mucuna         | 4              | 6          | -0.01 | 1                       | 4          | -0.04 | 0                    | 3          | -0.10 |
| Single cropping of Peanut                 | 22             | 5          | 0.13  | 16                      | 1          | 0.18  | 6                    | 0          | 0.20  |
| Association of Maize and Peanut           | 1              | 25         | -0.18 | 3                       | 13         | -0.12 | 1                    | 5          | -0.13 |
| Association Sorghum and Peanut            | 5              | 11         | -0.04 | 6                       | 5          | 0.01  | 0                    | 3          | -0.10 |
| Single cropping of Cowpea                 | 27             | 0          | 0.20  | 15                      | 2          | 0.15  | 6                    | 0          | 0.20  |
| Association Maize and Cowpea              | 0              | 24         | -0.18 | 1                       | 15         | -0.16 | 0                    | 5          | -0.17 |
| Association Sorghum and Cowpea            | 5              | 6          | -0.01 | 4                       | 2          | 0.02  | 0                    | 2          | -0.07 |
| Single cropping of Soybean                | 24             | 0          | 0.18  | 13                      | 2          | 0.13  | 4                    | 0          | 0.13  |
| Association Maize and Soybean             | 1              | 20         | -0.14 | 2                       | 13         | -0.13 | 0                    | 4          | -0.13 |
| Association Sorghum and Soybean           | 0              | 9          | -0.07 | 0                       | 2          | -0.02 | 0                    | 2          | -0.07 |
| Single cropping of Bambara groundnut      | 14             | 1          | 0.10  | 5                       | 3          | 0.02  | 3                    | 1          | 0.07  |
| Association Maize and Bambara groundnut   | 0              | 11         | -0.08 | 0                       | 4          | -0.05 | 0                    | 2          | -0.07 |
| Association Sorghum and Bambara groundnut | 0              | 0          | 0.00  | 0                       | 0          | 0.00  | 0                    | 0          | 0.00  |

### 3.6. Farmers' perception of the advantages and weaknesses of pure legume crops and cereal/legume associations

Farmers attributed 09 advantages to single cropping of legume crops, and 07 to cereal/legume associations (Table 7). Its main advantages are improving soil fertility (24.58% of respondents), better yielding (23.73%), low labour requirements (23.73%), and their role as food for humans and animals (10.17%). For cereal/legume associations, the main advantages cited are crop diversification (32.65%), improvement soil fertility (22.45%), food for humans and animals (16.33%), better yielding (14.29%), and efficient land use (10.20%).

The results show 09 weaknesses attributed to single cropping of legumes, and 06 weaknesses attributed to cereal/legume associations (Table 7). Difficulty in conducting some farm operations (27.54% of respondents), soil impoverishment (21.74%) and pest attacks (17.39%) were the main weaknesses attributed by farmers to single cropping legume. Difficulty conducting some farm operations (58.62% of respondents), low production (19.54%) and competition between crops (18.39%) were the main weaknesses attributed to cereal/legume combinations.

**Table 7: Advantages and weaknesses of legume insertion in the cropping system according to farmers**

| N°           | Criteria   | Advantages of single cropping of legumes (%) | Advantages of legumes/cereals association (%) |
|--------------|--|--|---|
| 1            | food (household and cattle.)                               | 10.17  | 16.33   |
| 2            | Convenient management of soil                              | -  | 10.20   |
| 3            | Precocity of the production cycle                          | 1.69   | -   |
| 4            | Crops diversification                                      | -  | 32.65   |
| 5            | Economic advantages  | 5.08   | 2.04  |
| 6            | High market demands  | 5.93   | -   |
| 7            | Weeding  | -  | 2.04  |
| 8            | Easy to cropping   | 3.39   | -   |
| 9            | Soil fertility management                                  | 24.58  | 22.45   |
| 10           | High yielding  | 23.73  | 14.29   |
| 11           | Low labour demand  | 23.73  | -   |
| 12           | Easy processing  | 1.69   | -   |
| <b>Total</b> |  | <b>100</b>                                   | <b>100</b>                                    |
| N°           | Criteria   | Weaknesses of single cropping of legumes (%) | Weaknesses of legumes/cereals association (%) |
| 1            | Soil fertility depletion                                   | 21.74  | 2.30  |
| 2            | Pest attack  | 17.39  | -   |
| 3            | Pesticide expenditures                                     | 2.90   | -   |
| 4            | Shelters for reptiles                                      | -  | 1.15  |
| 5            | Competition between cultures                               | -  | 18.39   |
| 6            | Difficulty for conducting some farming operations cultural | 27.54  | 58.62   |

|              |   |            |            |
|--------------|---|------------|------------|
| 7            | Damage to crops and harvests                | 11.59      | -          |
| 8            | Low production (low yield)                  | 1.45       | 19.54      |
| 9            | Market prices Instability                   | 2.90       | -          |
| 10           | Lack of equipment (cultivation and harvest) | 7.25       | -          |
| 11           | Requires more labor                         | 7.25       | -          |
| <b>Total</b> |   | <b>100</b> | <b>100</b> |

#### 4. Discussion

##### 4.1. Over 150% increase in legume area after 15 years

The results show that the characteristics of the farms surveyed have not changed significantly after 15 years. However, the area allocated to legumes in the cropping system increased by 156.47% from 2008 to 2022. In addition to the traditional legumes (cowpea, Bambara groundnut and groundnut), soybean and mucuna have also been introduced. This increase in the area planted with legumes in Koumbia can be explained by the intervention of research and development projects and non-governmental organizations (NGOs). Indeed, the farmers' community of Koumbia have been working with researchers since 2005 on innovative farming systems building (CPASI) using co-design approach (Vall et al., 2006; Coulibaly et al., 2012b; Vall et al., 2016). Also, Coulibaly et al. (2012b) investigated the integration of two legumes (cowpea and mucuna) both for single cropping and in association. In Koumbia, the introduction of mucuna (a forage legume) as an innovation in the cropping systems was driving by co-design work conducted by farmers in partnership with researchers (Vall et al., 2016). As for the soybean, it was also facilitated by the national policy and development projects in charge of promoting soybean sector in Burkina Faso (Guilloux et al., 2018). Indeed, a national policy for promoting soybean production was established in Burkina Faso targeted to 100,000 tones by 2022. The increase of farm areas under legumes crops can also be due to land pressure. With the expansion of areas under crops, there is a restriction of grazing land, and consequently the need for fodder production to address forage demand during the lean periods. Crop diversification to strengthen households' resilience to climate change and income source diversification is also one of reasons why legumes crop production is in increase; compared to most of the other crops, legume have high economic value.

Although the area planted with legumes has increased, yields have not varied significantly after 15 years. The noncompliance of farmer in conducting farm operations can explain this yield trends. Legume yields may also be underestimated, because for some crops such as cowpeas, part of the harvested product is directly used for family consumption specially during the lean season. Consequently, this part is not accounted during yield estimation. Soybean yields have risen from 160 kg/ha in 2007 to 867 kg/ha in 2021, according to growers. This yield, achieved in 2021, is in the same range as those obtained by Sermé

et al. (2018) with the farmers' practice (600 to 900 kg/ha). This may reflect the growing interest of producers in this legume crop, which is used for semi-industrial and artisanal processing into flour, oil, soumbala, poultry feed (roasted soybeans) and kebabs (Bila et al., 2009).

Despite the increase in the area under legumes, cotton and maize remain at the backbone of the cropping system in the village of Koumbia, with a slight increase in their area. This is justified by the fact that cotton, which is grown in rotation with maize, is a well-organized sector by the cotton companies. Yields of both crops even increased significantly after 15 years.

#### **4.2. Single cropping as the preferred method for integrating legumes into the cropping systems**

Growers with their own constraints and habits prefer integrating legumes crops in the existing cropping systems using single cropping approaches. This agroecological configuration of farming plots is in discordance with crop association generally perceived by agronomists as agroecology as a means of optimizing the Land Equivalent Ratio (LER) and the symbiotic relations between legumes and cereal crops. The choice for single cropping of legumes can also be explained by the advantages that farmers attribute to pure legume crops, namely high forage and grain production, low labour requirements and improved soil fertility. Research has also shown that yields in association are low compared with pure crops (Coulibaly et al., 2017; Zongo et al., 2021). The high production of fodder in single cropping of legumes is a way of building up fodder for cattle. For Toé et al. (2022), fodder crops are effective means of addressing fodder deficit in the farming systems. In addition to their role as fodder, groundnuts and cowpeas are high protein sources when consumed by household members. Due to their early maturation compared to the other crops, cowpea is even used as food during the cotton and maize harvest. Indeed, in this area, the harvest period often corresponds to the lean periods for households. The role played by legumes in soil fertilization is well demonstrated (Muhammad et al., 2010; Hamidou et al., 2018; Guinet et al., 2019). It is why legume insertion in the farming system is recommended either in rotation with other crops or in association. Guinet et al. (2019) found improvement in wheat yield when cropped in rotation after a large majority of legumes compared to the ones grown after cereal crops. Yeo et al. (2022), who showed a positive background-effect of groundnut and cowpea on eggplant, suggest that these two (2) legumes should be used as an organic alternative to mineral fertilizers. Current field management practices seriously compromise the possibilities of cereal-legume association (use of herbicides, mechanization of farm operations such as weeding, etc.).

However, with the reduction in arable land coupled with land pressure, the option of single cropping of legume crops may not last. It is therefore appropriate to work on the association options of legumes in association with other crops. These include crop diversification on the same plot, and optimization of farm lands used with land equivalent ratios higher than 1 (Toé et al., 2022).

As conclusion, the driving force behind agro-ecological transition that include legumes in Burkina Faso's western cotton zone, the limitations raised by farmers need to be address. These constrains include (i) the difficulty of conduction some farm operations (mechanical weeding, phytosanitary treatment, ridging, harvesting, threshing, post-harvest processes, etc.), (ii) pest preasure in the field and in store. weak value chain for fodder legumes and/or seed legumes can also be a hindrance to the large-scale integration of legumes into the cropping system.

## **5. Conclusion**

The inclusion of multi-interest legumes in a context of promoting agro-ecology appears to be both opportunities and challenges. The results of the study reported herein showed an increase in the area under legumes; over 150% after 15 years. The number of producers also increased for legumes, notably cowpeas and soybeans. *Mucuna*, a forage legume introduced by research, is also being adopted. Moreover, analysis of perceptions shows that single cropping of legumes remain the favorite insertion modality of farmers. One of the most important explanatory factors is the partnership-based co-design of innovative agricultural systems (CPSAI) implemented in western Burkina Faso. In order to maintain this dynamic and using legumes as a lever for agro-ecological transition in western Burkina Faso, addressing the obstacles to their introduction (mechanization of cultivation and harvesting operations, pest control) and on structuring their value chain (exchange platform between legume industry players) would seem appropriate.

## References

1. Amadou, Z., (2021). "Econometric analysis of the determinants of food consumption in the commune of Tahoua (Niger)". Rev. Mar. Sci. Agron. Vét. 9 (2): 293-300. French.
2. Azontondé, A., (1993). Degradation and restoration of barre lands (weakly desaturated ferralitic clay-sandy soils) in Benin. Cah. Orst. Sér. Pédol., 28: 217-226. French.
3. Barro, A., Sangaré, M., Coulibaly, K., Koutou, M. & Diallo, MA., (2016). Study of the modalities of corn/cowpea association in the villages of Koumbia and Gombêlédougou in the cotton-growing area of western Burkina Faso. Science and Technology Review of the CNRST, Natural and Applied Sciences Series, Special Issue No. 2: 151-163. French.
4. Bila, NK., Bonkougou, I. & Ouédraogo, M., (2009). Study on the soybean sector in Burkina Faso. Final report, Agency for the promotion of small and medium-sized enterprises in agriculture and crafts (APME 2A), Burkina Faso, 62 p, <http://www.ungana.org> > IMG > pdf > ungana\_11 (accessed 26/12/2022). French.
5. Carsky, RJ., Douthwaite, B., Manyong, VM., Sanginga, N., Schulz, S., Vanlauwe, B., Diels, J., & Keatinge, J., (2003). Improving soil management by introducing legumes into cereal systems in African savannahs. Cahiers Agricultures, 12, 227-233, <https://revues.cirad.fr/index.php/cahiers-agricultures/article/view/30394> .French.
6. Coulibaly, K., Vall, E., Autfray, P., Bacyé, B., Somda, I., Nacro, HB. & Sédogo, MP., (2012a). Co-design of technical itineraries for the pure cultivation of cowpea (*Vigna unguiculata* L. Walp.) and mucuna (*Mucuna deeringiana* [Bort], Merrill) in the western cotton zone of Burkina Faso: interests and limits. Journal of Agriculture and Environment for International Development – JAEID, 106 (2): 139-155, DOI: 10.12895/JAEID.20122.78. French.
7. Coulibaly, K., Vall, E., Autfray, P., Nacro, HB., & Sédogo, PM. (2012b). First results of ecological intensification and participatory approach in cotton-growing area of western Burkina Faso. African Agronomy, 24(2):129-141. French, <https://www.ajol.info/index.php/aga/article/view/82318>
8. Coulibaly, K., Gomgnimbou, APK., Traoré, M., Nacro, HB., & Sédogo, PM. (2017). Effects of maize-legume associations on maize (*Zeamays* L.) yield and fertility of a tropical ferruginous soil in western Burkina Faso. Afrique science, 13(6): 226 – 235. French. 19.pdf (afriquescience.net)
9. FAO. 2018. The 10 elements of agroecology: Guiding the transition to sustainable food and agricultural systems. 15p, <https://www.fao.org/3/i9037fr/i9037fr.pdf> (accessed 27/12/2022). French.

10. Gebhard, CA., Büchi, L., Liebisch, F., Sinaj, S., Ramseier, H., & Charles R. (2013). Screening of legumecrops for plant cover: nitrogen and weeds. *Swiss Agricultural Research* 4 (9): 384–393, French. <https://agriculture-de-conservation.com/Screening-de-legumineuses-pour.html>
11. Guinet, M., Nicolardot, B., Durey, V., Revellin, C., Lombard, F., Pimet, E., Bizouard, F., & Anne-Sophie Voisin, A-S. (2019). Symbiotic nitrogen fixation and the preceding effect: are all grain legumes equal? *Agronomic Innovations*, 74: 55–68. French. 10.15454/jj5qv. hal-02158172
12. Le Guilloux, G., David, A. & Pouzet, A. (2018). Differentiation through the organization of sectors: the experience of AGROPOL. *OCL* 25(2): D210. French. <https://doi.org/10.1051/ocl/2018023>
13. Hamidou, F., Harou, A., Achirou, BF., Halilou, O. & Bakasso, Y. (2018). Nitrogen fixation in peanut and cowpea under drought conditions for improving productivity in the Sahel. *Tropicultura*, 36(1): 63-79, French. DOI: 10.25518/2295-8010.984
14. Koné, M., Ouattara, Z.S. & Yeo L. 2019. Peanut production: what contribution to food security in the Dianra department (Côte d'Ivoire). *African Agronomy Special Issue (8) / AGRIEDAYS*: 93-102, French. <https://www.ajol.info/index.php/aga/article/view/192418>
15. Muhammad, A., Dikko, AU., Audu, M. & Singhet, A. (2010). Comparative Effects of Cowpea and Soybean Genotypes on N<sub>2</sub> - Fixation and N-Balance in Sokoto Dry Sub-Humid Agro-Ecological Zone of Nigeria. *Nigerian Journal of Basic and Applied Science*, 18(2): 297-303. <http://ajol.info/index.php/njbas/index>
16. Ouattara, B., Sangaré, M., Coulibaly, K. (2016). Options for Sustainable Intensification of Agricultural and Fodder Production in the Agropastoral Production System of the Cotton-Growing Areas of Burkina Faso. *CNRST Science and Technology Review, Natural and Applied Sciences Series, Special Issue No. 2*: 133-149. French.
17. Perinelle, A., Meynard, J.-M., & Scopel, E. (2021). Combining on-farm innovation tracking and participatory prototyping trials to develop vegetable-based cropping systems in West Africa. *Agricultural Systems*, 187(102978), 16 p. <https://doi.org/10.1016/j.agsy.2020.102978>
18. Sermé, I., Pouya, MB., Nignan, I. & Ouattara, K. (2018). Effect of applying soil fertilizer (NPK and Urea) in microdose on soybean and corn in Burkina Faso. *CNRST Science and Technology Review, Natural and Applied Sciences Series*, 34 to 37: 67-80. French.

19. Toé, A., Sanon, HO., Obulbiga, F. & Bougouma, VMC. (2022). Improving forage availability by different dual-purposesorghum and cowpea cultivation methods. *Journal of Animal & Plant Sciences*, 54 (1): 9808-9821, French. <https://doi.org/10.35759/JAnmPISci.v54-1.2>
20. Vall, E., Dugué, P. & Blanchard, M. (2006). Weaving agriculture-livestockrelationshipsthroughcotton. *Cahiers Agricultures*, 15 (1): 72–79, French. <https://revues.cirad.fr/index.php/cahiers-agricultures/article/view/30562>
21. Vall, E., Koutou, M., Blanchard, M., Coulibaly, K., Diallo, MA. & Andrieu, N. (2011). Agriculture-livestockintegration and ecological intensification in agro-sylvo-pastoral systems in western Burkina Faso, Tuy province. Proceedings of the ASAP seminar, November 2011, Bobo-Dioulasso, Burkina Faso. CD-ROM. French.
22. Vall, E., Chia, E., Blanchard, M., Koutou, M., Coulibaly, K. & Andrieu, N. (2016). Co-design in partnership of innovative agricultural systems. *Cahiers Agricultures*, 25, 15001. French. DOI: 10.1051/cagri/2016001
23. Yéo, KT., Fondio, L., Kouakou, KL., N’Gbesso, MFP. & Coulibaly, ND. (2022). Characterization and diversity of marketgardening production systems in the center (Bouaké) of Côte d’Ivoire with a view to an agroecological transition. *J. Anim. Plant Sci.*, 52(3): 9538-9551. French. <https://doi.org/10.35759/JAnmPISci.v52-3.3>
24. Zongo, KF., Hien, E, Maré, BT. & Guébré, D. (2021). Performance of the mixed sorghum-cowpea association on sorghum and soilproductivities in the Sudano-Sahelian zone of Burkina Faso. *Int. J. Biol. Chem. Sci.* 15(3): 987-1005. French. DOI: <https://dx.doi.org/10.4314/ijbcs.v15i3.12> .
25. Zougmore, R., Guillobez, S., Kambou, NF. & Ouattara, K. (2000). Sorghum-cowpeaintercropping: an effective technique againstrunoff and soilerosion in the Sahel (Saria, Burkina Faso). *AridSoilResearch&rehabilitation* 14: 329-342; <https://doi.org/10.1080/08903060050136441>