

Effect of Optimization use of Nutrient on Yield and Productivity of Mung bean (*Vigna radiata* L.)

Abstract

A field experiment was conducted at experimental farm, Department of Agronomy, Faculty of Agriculture and Veterinary Sciences, Mewar University Gangrar, Chittorgarh (Rajasthan) during Rabi season of 2023-24 to effect of optimization use of nutrient on yield and productivity of mung bean, variety "SML-832" was used in this study. The required quantities of fertilizers as per treatments were applied. The experiment was laid out in randomized block design with three replications consisting of ten treatments. The data recorded maximum yield attributes such as number of pods per plant (13.35), number of seed per pod (8.52), grain yield (1170.55 kg/ha), stover yield (3260.47 kg/ha) and maximum net return (67329.45 Rs/ha) and B:C ratio (2.28) was recorded with T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium. The minimum yield and profitability obtained with control treatment. Therefore, conclude be application of 100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium are indigenous sources of nutrient to enhance yield and productivity of mung bean.

Keywords: -Productivity; FYM; Vermicompost; Yield; Mung bean

1. Introduction

Green gram [*Vigna radiata* (L.) Wilczek] is one of the most ancient and extensively grown leguminous crops of India. It is a native of India and Central Asia and commonly known as mung bean. Green gram protein is deficient in methionine and cysteine but rich in lysine making it an excellent complement to rice. It is a good source of mineral, pro-vitamin A, B complex and ascorbic acid. India is one of the important mung bean growing countries in Asia with an area 8.7 million hectares and production of 8.83 million tonnes with a productivity of 1014 kg ha⁻¹ (Anonymous., 2020).

The productivity of this crop is very low because of its cultivation on marginal and sub marginal lands of low soil fertility where little attention is paying to adequate fertilization. In summer green gram, a high reduction in yield has been reported to occur due to non-use of fertilizers. Although, chemical fertilizer is playing a crucial role to meet the nutrients need of the crop, the imbalance and continuous use of chemical fertilizers has adverse effect on soil

physical, chemical and biological properties thus affecting the sustainability of crop production, besides causing environmental pollution (Singh and Sekhon 2008).

Consumption of chemical fertilizers will also be quite a limiting factor of agricultural production in future. Because of escalating energy cost, chemical fertilizers are not available at affordable price to the farmers. Therefore, there is an urgent need to reduce the usage of chemical fertilizers and in turn increase in the usage of organics is needed to check the yield and quality levels. On the other hand, use of organics alone does not result in spectacular increase in crop yields, due to their low nutrient status. Therefore, the aforesaid consequences have paved way to grow green gram by integration of organic and inorganic fertilizers along with biofertilizers (Ali *et al.* 2010).

Vermicompost has a higher nitrogen, phosphorus, and potassium content than typical heap manure, making it an excellent alternative to commercial fertilizers. Vermicompost typically contains 0.40–0.75%, 0.13–0.22% P, and 0.6–1.2% N. It also contains significant amounts of nutrients, a sizable population of beneficial microorganisms, and biologically active metabolites, in particular gibberellins, cytokinins, auxins, and group B vitamins (Jack *et al.* 2011).

Integration of organic manures and inorganic fertilizer materials has been found to be promising not only in maintaining higher productivity of crops and for providing stability in crop production, besides improving soil physical conditions (Verma *et al.* 2012). Farmyard manure and vermicompost have been advocated as good organic manure for use in integrated nutrient management programme in field crops. They are low cost and eco-friendly inputs, which have tremendous potential of fixing atmospheric nitrogen and can reduce the chemical fertilizer dose by 25–50% (Pattanayak *et al.* 2007).

2. Materials and Methods

A field experiment was conducted during Rabi season of 2023-24 at experimental farm, Department of Agronomy, Faculty of Agriculture and Veterinary Sciences, Mewar University Gangrar, Chittorgarh (Rajasthan). Soil of the experimental field was sandy loam in texture, saline in reaction with a pH value of 7.6, poor in organic carbon (0.16%), deficient in available zinc (0.48 ppm) and iron (1.2 ppm) low in available nitrogen (176 kg/ha) and phosphorus (20.2 kg/ha) but medium in available potassium (320 kg/ha). The experiment was laid out in randomized block design with three replications consisting of treatments viz. T₁-Control, T₂-RDF + Rhizobium, T₃-FYM @ 4 t ha⁻¹ + Rhizobium, T₄-100%

RDF + FYM @ 2 t ha⁻¹ + Rhizobium, T₅-75% RDF + FYM @ 2 t ha⁻¹ + Rhizobium, T₆-50% RDF + FYM @ 2 t ha⁻¹ + Rhizobium, T₇-Vermicompost @ 2 t ha⁻¹ + Rhizobium, T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium, T₉-75% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium and T₁₀-50% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium. The required quantities of fertilizers as per treatments were applied. The doses of NPK were applied in the form of urea, diammonium phosphate, and muriate of potash respectively. The half dose of nitrogen gives basal dose and remain two split doses after irrigation and full dose of phosphorus and potassium at basal dose. Vermicompost and FYM apply in field at field preparation before sowing. The seed treatment with Rhizobium culture. The yield parameters were calculated from output from the field. The profitability and productivity of mung bean was calculated from cost of field preparation to harvesting and threshing cost and out put from straw yield and grain yield as per market rate.

3. Results and Discussion

3.1 Yield attributes and yield

Data pertaining to effect of different organic and inorganic sources of nutrient on yield attributes and yield are presented in Table 1.0 and Figure 1.0. The organic and inorganic sources of nutrient were showed significant effect on yield attributes and yield of mung bean. The showed that maximum number of pods per plant with T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (13.35), it was at par with T₄-100% RDF + FYM @ 2 t ha⁻¹ + Rhizobium and T₉-75% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (13.25 and 12.76). The minimum number of pods per plant recorded with control treatment (68.58). Singh and Singh (2013), Hamza *et al.* (2016) and Yadav *et al.* (2017) reported similar findings. The maximum number of seed per pod with T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (8.52), it was at par with T₄-100% RDF + FYM @ 2 t ha⁻¹ + Rhizobium and T₉-75% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (8.30 and 8.00). The minimum number of seed per pod recorded with control treatment (6.75). The maximum grain yield with T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (1170.55 kg/ha), it was at par with T₄-100% RDF + FYM @ 2 t ha⁻¹ + Rhizobium and T₉-75% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (1145.25 and 1128.45 kg/ha). The minimum grain yield recorded with control treatment (775.25 kg/ha). Solanki *et al.* (2018), Sudipta *et al.* (2019), Tyagi and Singh (2019) and Sachan *et al.* (2020) stat that same conclusion. The maximum straw yield with T₄-100% RDF + FYM @ 2 t ha⁻¹ + Rhizobium (3260.47 kg/ha), it was at par with T₈-100% RDF +

Vermicompost @ 1.0 t ha⁻¹ + Rhizobium and T₉-75% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (3185.45 and 3110.58 kg/ha). The minimum straw yield recorded with control treatment (2536.45 kg/ha). Similar results were reported by Rajkhowa *et al.* (2002), Kinkar (2007), Vadgave (2010), Kushwaha (2013) and Somalraju *et al.* (2021).

3.2 Economics

Data pertaining to effect of different organic and inorganic sources of nutrient on economics presented in Table 2.0. The organic and inorganic sources of nutrient were showed significant effect on economic variability of mung bean. Data showed that the maximum cost of cultivation was recorded with treatment was recorded T₄ and T₈ (29500 Rs/ha). The minimum cost of cultivation was recorded with control treatment (24500 Rs/ha). The maximum gross return was recorded with treatment was recorded T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (96829.45Rs/ha). The minimum gross return was recorded with control treatment (60500.00 Rs/ha). Patil *et al.* (2011) and Verma *et al.* (2017) supported by similar findings. The maximum net return was recorded with treatment was T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (67329.45Rs/ha). The minimum net return was recorded with control treatment (36000.00 Rs/ha). The maximum B:C ratio was recorded with treatment was recorded T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium (2.28). The minimum B:C ratio was recorded with control treatment (1.47). Similar result also reported by Singh *et al.* (2018) and Marimuthu *et al.* (2023).

Conclusion

The findings of present investigation revealed that significant impact of different organic and inorganic nutrient sources on the yield and productivity of the mung bean. Among all treatment T₈-100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium registered the maximum production with higher net return. So, it was concluded that the treatment 100% RDF + Vermicompost @ 1.0 t ha⁻¹ + Rhizobium superior among all treatments.

Table.1.0Effect of different organic and inorganic sources of nutrient on yield attributes and yield of mung bean

Treatments	Number of pods per plant	Number of seeds per plant	Grain yield (kg/ha)	Straw yield (kg/ha)
T ₁ -Absolute control	10.02	6.75	775.25	2536.45
T ₂ -RDF + Rhizobium	11.32	8.15	1080.45	3085.45
T ₃ -FYM @ 4 t ha ⁻¹ + Rhizobium	10.95	7.82	1018.65	2862.45
T ₄ -100% RDF + FYM @ 2 t ha ⁻¹ + Rhizobium	13.25	8.30	1145.25	3260.47
T ₅ -75% RDF + FYM @ 2 t ha ⁻¹ + Rhizobium	12.70	7.89	1100.36	3012.45
T ₆ -50% RDF + FYM @ 2 t ha ⁻¹ + Rhizobium	11.28	7.75	1052.25	2905.78
T ₇ -Vermicompost @ 2 t ha ⁻¹ + Rhizobium	11.00	7.89	1022.47	2935.45
T ₈ -100% RDF + Vermicompost @ 1.0 t ha ⁻¹ + Rhizobium	13.35	8.52	1170.55	3185.45
T ₉ -75% RDF + Vermicompost @ 1.0 t ha ⁻¹ + Rhizobium	12.76	8.00	1128.45	3110.58
T ₁₀ -50% RDF + Vermicompost @ 1.0 t ha ⁻¹ + Rhizobium	11.35	7.78	1070.65	3028.85
S. Em. (±)	0.20	0.18	15.25	50.12
C.D. at 5%	0.59	0.53	45.85	151.02

Table.2.0Effect of different organic and inorganic sources of nutrient on economics of mung bean

Treatment	Cost of cultivation (Rs/ha)	Gross return (Rs/ha)	Net return (Rs/ha)	B:C ratio
T ₁ -Absolute control	24500.00	60500.00	36000.00	1.47
T ₂ -RDF + Rhizobium	28500.00	76580.00	48080.00	1.69
T ₃ -FYM @ 4 t ha ⁻¹ + Rhizobium	27000.00	78250.00	51250.00	1.90
T ₄ -100% RDF + FYM @ 2 t ha ⁻¹ + Rhizobium	29500.00	90500.00	61000.00	2.07
T ₅ -75% RDF + FYM @ 2 t ha ⁻¹ + Rhizobium	28500.00	85050.00	56550.00	1.98
T ₆ -50% RDF + FYM @ 2 t ha ⁻¹ + Rhizobium	26500.00	78500.00	52000.00	1.96
T ₇ -Vermicompost @ 2 t ha ⁻¹ + Rhizobium	26300.00	82500.00	56200.00	2.14
T ₈ -100% RDF + Vermicompost @ 1.0 t ha ⁻¹ + Rhizobium	29500.00	96829.45	67329.45	2.28
T ₉ -75% RDF + Vermicompost @ 1.0 t ha ⁻¹ + Rhizobium	28500.00	88500.00	60000.00	2.11
T ₁₀ -50% RDF + Vermicompost @ 1.0 t ha ⁻¹ + Rhizobium	27500.00	81500.00	54000.00	1.96




Figure 1.0 Effect of vermicompost and zinc application on yield of gram

References

- Ali MA, Abbas G, Mohy-ud-Din Q, Ullah K, Abbas G and Aslam M. 2010. Response of mung bean (*Vigna radiata*) to phosphatic fertilizer under arid climate. *The Journal of Animal and Plant Sciences* **20** (2): 83-86.
- Anonymous (2020). District Wise Area, Production and Yield per Hectare of important food and non-food crop. Directorate of Agriculture, Jaipur, Krishi Bhavan (Raj.).
- Hamza, B. A., Chowdhury, M. A. K., Rob, M. M., Miah, I., Habiba, U. and Rahman, M. Z. (2016). Growth and yield response of mung bean as influenced by phosphorus and boron application. *American J. Experit. Agri.* **11**(3):1-7.
- Jack, A.L.H., Rangarajan, S.W., Culman, T., Sooksa, N. and Thies, J.E. 2011. Choice of organic amendments in tomato transplants has lasting effects on bacterial rhizosphere communities and crop performance in the field. *Applied Soil Ecology*, **48**(1):94-101.
- Kinkar R. 2007. Response of green gram [*Vigna radiata* (L.) Wilczek] to integrated nutrient management (INM) in vertisols of Malwa Plateau. M.Sc. (Ag.) Thesis, JNKVV, Jabalpur. p 51.
- Kushwaha R. 2013. Effect of Different N, P and K Doses on Growth, Yield and Economics of Sesame (*Sesamum indicum* L.) Under Rainfed Conditions. M.Sc. (Ag) Thesis, JNKVV, Jabalpur. p 78.
- Madholiya, N. S. (2015). Integrated crop management practices for maximization of yield of black gram [*Vigna mungo* (L.) hepper]. M. Sc. (Ag.) Thesis, Agronomy, Rajmata Vijayaraje Scindia Krishi Vishwa Vidyalaya, Gwalior (MP).
- Marimuthu, S., Gnanachitra, M., Prabu Kumar, G., & Surendran, U. (2023). Effect of organic and inorganic sources of phosphorus for enhancing productivity and phosphorus use efficiency in blackgram under acid soils. *Journal of Plant Nutrition*, **46**(9), 1845-1855.
- Patil, S.C., Jagtap, D. N. and Bhale, V.M. (2011). Effect of phosphorus and sulphur on growth and yield of moong bean. *Internat. J. agric.Sci.*, **7**(2):348-351.
- Pattanayak SK, Rao DLN and Mishra KN. 2007. Effect of biofertilizers on yield, nutrient uptake and nitrogen economy of rice-peanut cropping sequence. *Journal of the*

Indian Society of Soil Science 55: 184–9.

- Rajkhowa DJ, Saikia M and Rajkhowa KM. 2002. Effect of vermicompost with and without fertilizer on green gram. *Legume Research*, **25** (4): 295-296.
- Singh G, and Sekhon HS. 2008. Effect of various inputs on the growth and yield of summer green gram (*Vigna radiata*). *Indian Journal of Agricultural Sciences* **78** (1): 87-89.
- Singh, A., Kumar, B. and Pandey, A.K. (2018). Response of phosphorus, PSB and press mud on dry matter accumulation, phosphorus content and uptake of urdbean. *Int. J. of Chemical studies*, **5**(5): 2351- 2353.
- Singh, S. K. and Singh, G. C. (2013). Effect of phosphorus, sulphur and zinc on nutrient composition in black gram. *J. Rural and Agril. Res. Vol.* **13** No.2, 63-64.
- Solanki, R.L., Sharma, M., and Indoria, D. 2018. Effect of Phosphorus, Sulphur and PSB on Yield of Indian Mustard (*Brassica juncea* L.) and Available Macronutrients in Soil. *Journal of the Indian Society of Soil Science*, **66**(4), 415-419.
- Somalraju, S., Goyal, G., Singh Gurjar, L., Chaturvedi, M., & Singh, R. (2021). Effect of organic and inorganic fertilizer on the growth and yield of green gram (*Vigna radiata* L.). *The Pharma Innovation Journal*, **10**(12), 1959-1962.
- Vadgave SM. 2010. Studies on integrated nutrient management on seed yield, quality and storability in green gram [*Vigna radiata* (L.) Wilczek]. M.Sc. Thesis, University of Agriculture, Kerala (India).
- Verma S, Singh HV and Saxena R. 2012. Relative performance of soybean (*Sesamum indicum* L.) under organic, inorganic and integrated nutrient management. *Indian Journal of Agricultural Sciences* **83** (3): 143–149.
- Verma, G., Kumawat, N. and Morya, J. (2017). Nutrient management in mung bean [*Vigna radiata* (L.) Wilczek] for higher production and productivity under semi-arid tract of central India. *Int. J. Curr. Microbial. App. Sci* **6** (7): 488-493.
- Yadav, M., Yadav, S. S., Kumar, S., Yadav, T. and Yadav, H. K. (2017). Effect of phosphorus and bio-fertilizers on growth and yield of urd bean [*Vigna mungo* (L.) Hepper]. *Int. J. of Plant and Soil Sci.*, **18** (5): 1-7.