

Growth and Yield of Vertically Trained Watermelon [*Citrullus lanatus* L. (Thunb.) Matsum. & Nakai] Under Different Levels of Spacing, Training and Pruning under Rain Shelter

ABSTRACT

This study investigated the effects of spacing, training and pruning on the growth and yield of watermelon hybrid 'Saraswati' trained vertically under rain shelter. The treatments were factorial combinations of three planting densities (1.50 m × 0.60 m, 1.50 m × 0.45 m and 1.50 m × 0.30 m), two training levels (nipping to one vine, nipping to two vines) and two pruning heights (pruned at 2m and unpruned) arranged in randomized block design with two replications and control with horizontal training of vines in open condition. Spacing, training and pruning levels had varying influence on growth and yield parameters of vertically trained watermelon. Wider spacing of 1.5 m x 0.6 m recorded the highest vine length, highest fruit weight and yield per plant. Earliness in days to flowering as well as node number was observed in widely spaced plants trained to single vine. Plants trained to two vines exhibited the highest number of fruits per plant, yield per plant and yield per plot. Unpruned plants trained to two vines at closer spacing of 1.50 m × 0.30 m recorded the highest number of fruits per plant and yield per plot (90.10 kg).

Key Words: *Citrullus lanatus*, watermelon, rain shelter, spacing, training, pruning

INTRODUCTION

Watermelon [*Citrullus lanatus* (Thunb.) Matsum & Nakai] is an important *cucurbitaceous* crop grown in tropical and subtropical areas of the globe. It is primarily grown for its refreshing juice and sweet flesh, and it holds the highest global consumption among cucurbits (Goreta *et al.*, 2005). China is the leading producer of watermelon globally. In India, watermelon is cultivated in 1.24 lakh hectares with a production of 35.05 lakh tonnes (GOI, 2022).

The idea of using rain shelters for year-round vegetable cultivation is becoming more popular in Kerala, with the financial assistance from the government. Rain shelters are being encouraged to support small-scale farming and reduce the use of harmful pesticides in commercial vegetable cultivation. Currently, only few vegetables like cucumber, capsicum, yard long beans, and tomatoes are cultivated under protected conditions in Kerala. It is important to promote cultivation of more vegetables in protected structures (Narayanankutty *et al.*, 2014).

Watermelon is a highly sought after fruit in Kerala, but its cultivation is limited. Compared to the traditional horizontal method of trailing, growing watermelon plants vertically in rain shelters allows accommodation of greater number of plants in unit area. Managing plant density is crucial for achieving higher number of commercially viable fruits. Altering plant densities through

Formatted: Font: Italic

Commented [A1]: <https://www.fao.org/faostat/en/#data/QCL/visualize>

Commented [A2R1]: FAOSTAT data is more common in articles. Maybe the values can be updated with FAO values according to the given address.

varying levels of spacing is a significant factor in agricultural management. This technique can be utilized to improve crop productivity as it significantly impacts the growth and yield of each plant (Diepenbrock, 2000).

To improve productivity and fruit quality, it is crucial to effectively train watermelon plants in a vertical system. Appropriate training makes it simpler to apply pesticides, improves ventilation for plants, and guarantees that solar radiation is distributed uniformly throughout the canopy. Additionally, this method enables higher plant density, leading to a greater number of fruits per unit area. A study showed that training to two vines notably boosted the number of fruits and yield per plant when done under a rain shelter. However, it was observed that when training was done in an open environment, it had no discernible effect (Nisha and Sreelathakumary, 2020).

Pruning vines is a crucial practice in crop management that offers several agricultural benefits. Primarily, it facilitates mechanical harvesting by ensuring that vines are well-organized and manageable, allowing machines to navigate and harvest effectively. Pruning also supports the production of hybrid seeds, as it helps control the plant's flowering and pollination patterns. By reducing dense foliage, pruning improves air circulation and sunlight penetration, which limits the moist, shaded environments, thereby reducing pest and disease incidence. Additionally, pruning enables farmers to increase plant population per unit area without risking significant yield reduction, as it helps manage vine growth and nutrient distribution, which maintains optimal productivity. Pruning also promotes the production of uniform fruits by directing the plant's energy to fewer, healthier fruits, which leads to consistent size and quality. According to Oga and Umekwe (2016), adopting pruning as a regular cultural practice can significantly improve crop growth and yield, making it a recommended approach for maximizing production in vine crops.

The potential of commercial cultivation of watermelon, especially mini and icebox types, is significant due to their small size. Therefore, it is important to focus on studying plant density and training levels in watermelon under rain shelter. Consequently, this research aims to assess the impact of various spacing, training, and pruning levels on the growth and yield of watermelon under rain shelter.

MATERIALS AND METHODS

The experiment was carried out at Department of Vegetable Science, College of Agriculture, Vellayani, Kerala Agricultural University during 2021-22 with the watermelon hybrid Saraswati. The treatments were factorial combinations of three levels of spacing (1.50 m × 0.60 m, 1.50 m × 0.45 m and 1.50 m × 0.30 m), two training levels (nipping to one vine, nipping to two vines) and two pruning heights (pruned at 2 m and unpruned) arranged in randomized block design with two replications and control with horizontal training of vines in open condition.

Experimental site

The site of rain shelter was located at 8°25'53.5" N latitude and 76°59'14.9" E longitude at an altitude of 29 m above sea level. The location of open field was 8°25'53.7" N latitude and 76°59'15.8" E longitude. The soil of experiment site is red loam of Vellayani series, texturally classified as sandy clay loam.

Commented [A3]: Watermelons are generally grown under greenhouses and in the open. Therefore, the differences between rain shelter and greenhouse and open cultivation can be mentioned in the introduction.

Commented [A4]: More resources should be used in this section

Commented [A5]: There are many sources in the literature. At this stage, a given source is quite insufficient. The searches are insufficient compared to a comprehensively written paragraph.

Commented [A6]: Will mini and icebox types of watermelons be used in the study? If so, information about these watermelon types should be provided at the introduction stage.

Commented [A7]: This study should be set up as a trial plan with at least 3 replications. In trials conducted with 2 replications, the margin of error is high and the trial is not reliable. The number of replications of the study should be checked again.

Field preparation and planting

The experimental area was deeply ploughed up to 50 cm and weeds and stubbles were removed. Farm yard manure @ 25 t ha⁻¹ and Rock Phosphate (125 kg ha⁻¹) was applied before last ploughing. Raised beds of one meter width and one foot height were taken with channels of 50 cm between the beds. The beds were mulched with silver on black polyethylene sheet of 50 µ thickness after laying drip lines. Seedlings were raised in pro trays using cocopeat and vermicompost as media. Twelve days old seedlings at 2-3 true leaf stage were transplanted at required spacing.

Commented [A8]: On what date were the seedlings planted? Planting dates are important in cultivation.

Training and Pruning

The plants were trained vertically under the rain shelter. In T₁ (nipping to one vine) only the main vine was allowed to grow. In T₂ (nipping to two vines), two vigorously growing laterals were allowed after nipping off the main vine. GI wires were tied over the cropping rows at 2.5 m height and plastic strings were tied on them to trail the plants. The first three secondary branches were removed and the rest were pruned after the third leaf (Campagnol *et al.*, 2012). The plants were trailed horizontally under open.

Fertigation

Drip lines were laid with a lateral per bed and drippers with a discharge rate of 2 l/hr at required spacing (60 cm, 45 cm and 30 cm) were used. Fertigation was done at three days interval using fertigation pump. Water soluble fertilizers, 19:19:19, 13:0:45, urea and 12:61:0 was used (Nisha, 2017). Submains and laterals are flushed prior to fertigation process and continuous drip irrigation was given for five to ten minutes after each fertigation. Disc filter and screen filter were cleaned at regular intervals.

Observations on growth and yield parameters were recorded and data were analysed statistically by applying the techniques of analysis of variance (Panse and Sukhatme, 1985).

Commented [A9]: What parameters were used in the examination? This stage must be written. How many plants were used for each parameter and how?

RESULTS AND DISCUSSION

The length of the vines was significantly influenced by the different plant densities as shown in Table 1. The longest vine, measuring 3.84 m, was observed in D₁ (1.50 m × 0.60 m), while the shortest vine (3.14 m), was in D₃ (1.50 m × 0.30 m). This could be attributed to reduced competition for water, nutrients and light among crops in wider spacing. Unpruned plants exhibited longer vines (5.00 m) compared to pruned ones (2.00 m). These findings are consistent with the research conducted by Sabo *et al.* (2013) and Lemos *et al.* (2022). The different training levels did not have a significant impact on vine length, aligning with the conclusions drawn by Nisha and Sreelathakumary (2020). A notable difference was observed between the treatments and the control as indicated in Table 2, with the control displaying longer vines (5.85 m) compared to the treatments.

The number of days taken for first female flower opening is a crucial factor in predicting early yield. Planting density had a notable impact on the timing of flowering and the location of the first flower (Table 1). The earliest flowering occurred in plants with wider spacing, single vines, and those that were pruned. This could be attributed to better interception of sunlight and

Commented [A10]: 15.56 days (1.50 x0.60)

Commented [A11]: Single vine T1(17.15 days)

Commented [A12]: 16.88 number

nutrient absorption from soil, leading to the accumulation of more photosynthates and prompting earlier flowering compared to plants with closer spacing. Similar findings were documented by Oga and Umekwe (2016). There was a significant contrast between the treatments and the control (Table 2), with the control exhibiting late flowering.

The position of the first female flowering node is an important character in watermelon. First female flower at the lowest node signifies early entry of watermelon into reproductive phase and can potentially capture the market early. The study showed that the spacing and training significantly influenced the nodal position of the first male and female flowers. In D₁ and T₁ plants, the female flower appeared at lower nodes (Table 1). Pruning did not have a notable impact on the nodal position of the flowers, which is consistent with the findings of Jaffar and Wahid (2014) and Sharma *et al.* (2018). While there were no significant differences in treatment interactions, there was a notable contrast between the treatments and the control (Table 2). The control had the first flowers at higher nodes.

Table 1. Effect of spacing, training and pruning on growth parameters in vertically trained watermelon under rain shelter

Treatments	Vine length (m)	Days to first male flower	Node to first male flower	Days to first female flower	Node to first female flower
Spacing					
D ₁ (1.50 m x 0.60 m)	3.84 ^a	15.56 ^c	4.68 ^c	22.41 ^c	13.64 ^c
D ₂ (1.50 m x 0.45 m)	3.52 ^b	16.92 ^b	5.55 ^b	23.53 ^b	14.57 ^b
D ₃ (1.50 m x 0.30 m)	3.14 ^c	18.45 ^a	6.37 ^a	24.12 ^a	15.34 ^a
SE _m (±)	0.034	0.035	0.075	0.044	0.070
CD (0.05)	0.106	0.109	0.235	0.138	0.216
Training					
T ₁ (Single vine)	3.61	16.79 ^b	5.36 ^b	23.14 ^b	14.36 ^b
T ₂ (Two vines)	3.42	17.15 ^a	5.67 ^a	23.57 ^a	14.67 ^a
SE _m (±)	0.087	0.029	0.062	0.036	0.057
CD (0.05)	NS	0.089	0.192	0.113	0.177
Pruning					
P ₁ (Pruned at 2.0 m)	2.00 ^b	16.88 ^a	5.52	23.24 ^b	14.53
P ₂ (Unpruned)	5.00 ^a	17.07 ^b	5.55	23.47 ^a	14.50
SE _m (±)	0.068	0.029	0.062	0.036	0.057
CD (0.05)	0.087	0.089	NS	0.113	NS
Control (open condition)	5.85	18.9	6.38	26.38	16.62

NS – Non significant

Table 2. Interaction effects of treatments on growth characters in vertically trained watermelon under rain shelter

Commented [A13]: There are many current literature studies on watermelon flowering.

Commented [A14]: values must be written explicitly

Treatments	Vine length (m)	Days to first male flower	Node to first male flower	Days to first female flower	Node to first female flower
D × T × P					
d1t1p1	2.00	15.33	4.50	22.10	13.50
d1t1p2	5.66	15.48	4.48	22.30	13.47
d1t2p1	2.00	15.64	4.84	22.47	13.80
d1t2p2	5.70	15.78	4.89	22.75	13.76
d2t1p1	2.00	16.52	5.37	23.25	14.42
d2t1p2	5.04	16.86	5.37	23.45	14.42
d2t2p1	2.00	17.05	5.68	23.62	14.70
d2t2p2	5.04	17.23	5.75	23.80	14.73
d3t1p1	2.00	18.18	6.20	23.75	15.23
d3t1p2	4.31	18.39	6.25	24.00	15.10
d3t2p1	2.00	18.54	6.50	24.25	15.50
d3t2p2	4.27	18.66	6.52	24.50	15.52
SE _m (±)	0.068	0.07	0.151	0.089	0.139
CD (0.05)	NS	NS	NS	NS	NS
Control	4.55	18.9	6.38	26.38	16.62
Control vs Treatment	S	S	S	S	S

NS – Non significant

All treatments had a notable impact on the average weight of fruit, as shown in Table 3. The highest fruit weight of 2.84 kg was observed when planting at 1.50 m × 0.60 m. Wider spacing allowed the plants to grow longer and develop more branches to accommodate larger fruits. Additionally, the wider spacing provided more room for the developing fruits to grow bigger. These findings are consistent with the research of Dahake *et al.* (2020) and Da Silva *et al.* (2021). Single vine training resulted in higher fruit weight compared to double vine plants. Similarly,

pruned plants produced heavier fruits than unpruned plants. Pruning facilitates the distribution of assimilates generated through photosynthesis, leading to cell enlargement and improved fruit characteristics. Campagnol (2012) found that training and pruning levels did not significantly influence fruit weight. Vertically trained watermelon fruits under rain shelter were lower than those grown horizontally in open conditions, as indicated in Table 4. The cropping density of mini watermelons trained in a vertical system influenced leaf area and solar radiation exposure, consequently affecting photosynthetic efficiency and fruit mass (Watanabe *et al.*, 2003).

Number of fruits per plant and yield per plot increased with increasing plant densities (Table 3). Lower plant spacing increase competition for water, light and nutrients, producing smaller fruits but increases yield per unit area, since a greater number of plants can be accommodated. Training to two vines recorded higher number of fruits than single vine. Similarly unpruned plants recorded highest number of fruits. Longer vine length and extended crop duration in rain shelter might have led to an increase in the number of fruits (Nisha, 2017). Maximum number of fruits per plant and yield per plot (78.37 kg) was recorded in high planting density. Increasing the density of planting can optimize land use effectively without reducing yield. In fact, yield per unit area generally improves as plant density rises, reaching a peak at an optimal population level. Beyond this point, however, yields may decline due to increased competition among plants for resources such as nutrients, water and sunlight. These results are supported by the findings of Da Silva *et al.* (2021). Wider spacing of 1.50 m x 0.6 m recorded the highest yield per plant. This might be due to higher fruit setting and fruit retention with increased spacing (Deka *et al.*, 2024). Among interactions, unpruned double vined plants at closer spacing of 1.50 m x 0.30 m recorded the highest value of yield per plot. This is in accordance with the findings of Anwar *et al.* (2019), where greater crop density increased the yield and number of fruits per unit area, but decreased yield per plant. Neupane (2023) reported increased yield in watermelon at a spacing of 2m x 1m with black silver polythene mulch.

Table 3. Effect of spacing, training and pruning on yield parameters in vertically trained watermelon under rain shelter

Treatments	Fruit weight (kg)	Fruits per plant	Yield per plant (kg)	Yield per plot (kg)
Spacing				
D ₁ (1.50 m x 0.60 m)	2.84 ^a	1.96	5.50 ^a	55.04 ^c
D ₂ (1.50 m x 0.45 m)	2.46 ^b	2.05	4.82 ^b	62.61 ^b
D ₃ (1.50 m x 0.30 m)	2.15 ^c	2.09	3.92 ^c	78.37 ^a
SE _m (±)	0.030	0.047	0.134	1.874
CD (0.05)	0.093	NS	0.418	5.482
Training				
T ₁ (Single vine)	2.84 ^a	1.50 ^b	4.21 ^b	58.36 ^b
T ₂ (Two vines)	2.13 ^b	2.56 ^a	5.28 ^a	72.32 ^a

SE _m (±)	0.024	0.039	0.110	1.530
CD (0.05)	0.076	0.120	0.341	4.762
Pruning				
P ₁ (Pruned at 2.0 m)	2.56 ^a	1.81 ^b	4.46 ^b	61.21 ^b
P ₂ (Unpruned)	2.39 ^b	2.20 ^a	5.03 ^a	69.47 ^a
SE _m (±)	0.042	0.039	0.110	1.530
CD (0.05)	0.132	0.120	0.341	4.762
Control (open condition)	3.24	2.87	6.308	63.06

NS – Non significant

Table 4. Interaction effects of treatments on yield characters in vertically trained watermelon under rain shelter

Treatments	Fruit weight (kg)	Fruits per plant	Yield per plant (kg)	Yield per plot (kg)
D × T × P				
d1t1p1	3.18	1.33 ^g	4.22 ^{ef}	42.23 ^f
d1t1p2	3.06	1.67 ^f	5.29 ^{bcd}	52.94 ^{ef}
d1t2p1	2.53	2.17 ^{cd}	5.67 ^{bc}	56.79 ^{de}
d1t2p2	2.56	2.67 ^b	6.82 ^a	68.22 ^{cd}
d2t1p1	2.86	1.84 ^{ef}	5.18 ^{cd}	67.38 ^{cd}
d2t1p2	2.65	1.33 ^g	3.39 ^{fg}	44.18 ^f
d2t2p1	2.40	2.01 ^{de}	4.65 ^{de}	60.49 ^{de}
d2t2p2	1.93	3.01 ^a	6.03 ^{ab}	78.39 ^{bc}
d3t1p1	2.65	1.17 ^g	3.02 ^g	60.44 ^{de}
d3t1p2	2.61	1.67 ^f	4.15 ^{ef}	82.99 ^{ab}
d3t2p1	1.82	2.33 ^c	3.99 ^{ef}	79.92 ^{ab}
d3t2p2	1.52	3.17 ^a	4.50 ^{de}	90.10 ^a
SE _m (±)	0.060	0.095	0.269	3.747

CD	NS	0.295	0.836	11.664
Control	3.24	2.87	6.308	63.06
Control vs Treatment	S	NS	S	NS

NS – Non significant

The number of fruits per plant and the yield per plot both increased with increasing plant densities (Table 3). Closer planting increased fruit number but lowered mean fruit weight (Filho *et al.*, 2015). Decreasing the distance between plants led to increased competition for water, light, and nutrients, resulting in smaller fruits but a higher yield per unit area, due to accommodating a greater number of plants. Training the plants to two vines resulted in a higher number of fruits compared to a single vine. Similarly unpruned plants recorded highest number of fruits. The longer vine length and extended crop duration in a rain shelter likely contributed to the increase in the number of fruits, as noted by Nisha (2017). The highest number of fruits per plant and the highest yield per plot (78.37 kg) were recorded at high planting density. Increasing the planting density can effectively optimize land use without reducing the yield. In fact, the yield per unit area generally improves as the plant density rises, reaching a peak at an optimal population level. Beyond this point, however, yields may decline due to increased competition among plants for resources like nutrients, water, and sunlight. These results align with the findings of Da Silva *et al.* (2021). A wider spacing of 1.50 m x 0.6 m resulted in the highest yield per plant, likely due to increased fruit setting and retention, according to Deka *et al.* (2024). Among interactions, unpruned double-vined plants at a closer spacing of 1.50 m x 0.30 m resulted in the highest value of yield per plot. This is consistent with the findings of Anwar *et al.* (2019), where a greater crop density increased the yield and number of fruits per unit area but decreased the yield per plant. Neupane (2023) reported an increased yield in watermelon at a spacing of 2m x 1m with black silver polythene mulch.

CONCLUSION

The results of the present study revealed that growing watermelon vertically under rain shelter by training the plants to two vines and allowing them to grow unpruned, improves the yield per unit area. A spacing of 1.50 m x 0.30 m and unpruned plants trained to two vines was found to be the best with highest number of fruits per plant and yield per unit area.

REFERENCES

1. Anwar N A, Gad A A, Bardisi A and Zyada H G .2019. Effect of plant spacing and apical shoot pinching on growth and productivity of watermelon plants under sandy soil conditions. *Zagazig J. Agric. Res.* 46(2): 357-365.
2. Campagnol R, Mello S C and Barbosa J C. 2012. Vertical growth of mini watermelon according to the training height and plant density. *Hortic. Bras.* 30(4): 726-732.
3. Dahake L Z, Sonkamble A M, Meshram L T and Patil S R. 2020. Effect of spacing on growth and yield of watermelon. *J. Pharma Phytochem.* 9(5): 751-753.

4. Da Silva AV, Da Silva CM, Goncalves CN, Filho MCDO, De Sousa Pereira C, Andrade MJKL, Pessoa WRLS. 2021. Productive potential of watermelon under different plant spacings in the semi-arid region of Brazil. *Austr. J. Crop Sci.* 15(2):238-243
5. Deka, B, Jahnabi, H, Pritom K B, Utpal K, Ananta S, Prakash K, Bhabesh G, Soumitra G, Bikash H, and Karobi H. 2024. Influence of crop geometry, fruit thinning and nutrient management on yield and yield-related attributes of watermelon (*Citrullus lanatus* Thunb.)". *Inter. J. Environ. Climate Change* 14 (1):443-52.
6. Diepenbrock W. 2000. Yield analysis of winter oilseed rape (*Brassica napus* L.): a review. *Field Crops Res.* 67: 35-49.
7. Filho, A. B. C, Feltrim, A. L., Cortez, J. W. M., Goncalves, M. V., Pavani, L. C., Barbosa, J. C. 2015. Nitrogen and potassium application by fertigation at different watermelon planting densities. *J. Soil Science Plant Nutrition* 15:928-937.
8. GOI [Government of India]. 2022. Agricultural statistics at a glance. Ministry of Agriculture and Farmers Welfare, New Delhi, 92p.
9. Goreta S, Perica S, Dumicic G, Buca L and Zanic K. 2005. Growth and yield of watermelon on polyethylene mulch with different spacings and nitrogen rates. *Hort. Sci.* 40(2): 366-369.
10. Jaffar A and Wahid F. 2014. Effect of row spacing on growth, yield and yield components of cucumber varieties. *Sci. Letter* 2: 33-38.
11. Lemos, D. A, Hamilton, Cesar, de, Oliveira, Charlo., Antonio, Carlos, Barreto., Dinamar, Márcia, da, Silva, Vieira., Ernane, Miranda, Lemes., Jose, Luiz, Rodrigues, Torres. 2022. Production and fruit quality of watermelon hybrids under different plant spacing. *Revista Brasileira de Engenharia Agricola e Ambiental*, 26(6):459-465.
12. Narayanankutty C, Sreelatha U, Jyothi M L and Gopalakrishnan T R. 2014. Advances in protected cultivation of vegetables in Kerala. In: Singh B, Singh B, Sabir N and Hasan M. (eds.), *Advances in protected cultivation*. New India Publishing Agency, New Delhi, pp: 133-141.
13. Neupane R., Tripathi K.M., Panthi B., Sapkota, P. 2023. Effects of Mulching and Plant Spacing on Growth, Yield and Quality of Watermelon (*Citrullus lanatus*) in Rapti Riverbed, Dang, Nepal. *SAARC J. Agric.* 21(2):41-52.
14. Nisha S K. 2017. Standardization of agrotechniques for precision farming in watermelon [(*Citrullus lanatus* (Thunb.) Matsum & Nakai)]. Ph. D. (Hort.) Thesis, Kerala Agricultural University, Thrissur, 213p.
15. Nisha S K and Sreelathakumary I. 2020. Growth and yield of watermelon (*Citrullus lanatus* (Thunb.) with different levels of fertigation and drip irrigation. *J. Krishi Vigyan.* 8(2): 157.
16. Oga I O and Umekwe P N. 2016. Effects of pruning and plant spacing on the growth and yield of watermelon (*Citrullus lanatus* L.) in Unwana- Afikpo. *Int. J. Sci. Res.* 5(4): 110-115.
17. Panse V G and Sukhatme P V. 1985. *Statistical Methods for Agricultural Workers* (4thEd.). Indian Council of Agricultural Research, New Delhi 347p.
18. Sabo M U, Wailare M A, Aliyu M, Jari S and Shuaibu Y M. 2013. The effect of NPK fertilizer and spacing levels on growth and yield of watermelon (*Citrillus lanatus* L). *Scholarly J. Agric. Sci.* 3: 325- 330.
19. Sharma D, Sharma V K and Kumari A. 2018. Effect of spacing and training on growth and yield of polyhouse grown hybrid. cucumber (*Cucumis sativus* L.). *Int. J. Curr. Microbiol. App. Sci.* 7(05):1844-1852.

20. Watanabe S. 2014. Fruit productivity of vertically trained watermelon plants. *JARQ* 48(2): 121-131.

[Wahid, 2014](#)

Commented [A15]: This research is not available

UNDER PEER REVIEW