

Original Research Article

ASSESSMENT OF PLASMA LEVELS OF CALCIUM, CREATININE AND GLUCOSE AMONG HYPERTENSIVE PATIENTS IN EDO STATE, NIGERIA

ABSTRACT

Background: Hypertension is a medical condition in which blood arteries have a higher pressure, requiring the heart to work harder to pump blood. Hypertension is indicated by systolic blood pressure (SBP) of 130mmHg or higher and diastolic blood pressure (DBP) higher than 80mmHg.

Aims: This study was set out to assess the plasma levels of calcium, creatinine and glucose among hypertensive patients in Etsako west local government area (L.G.A), Edo State, Nigeria.

Study design: This study is an analytical cross-sectional study conducted among 150 hypertensive patients defined as test participants and 150 non-hypertensive individuals defined as control participants aged 20 years and above selected by simple random sampling.

Methods: Hypertension was determined by sphygmomanometer blood pressure reading of $\geq 140/90$ mmHg. Five millilitres of blood sample was collected from each participant using standard venipuncture technique and plasma levels of calcium, creatinine and glucose were determined using O'Cresolphthalein method, Jaffe-Slot alkaline picrate method and glucose-oxidase method respectively. Data was analyzed using Student's t-test. Statistical significance was set at $P < 0.05$.

Results: There was a significant difference in the mean age of hypertensive patients (52.2 ± 1.18 years) and non-hypertensive patients (35.1 ± 0.91 years) ($P = 0.001$). The systolic (156.2 ± 1.18 mmHg) and diastolic blood pressures (95.8 ± 0.70 mmHg) were significantly higher in hypertensive patients compared with the non-hypertensive controls (SBP: 118.6 ± 0.64 mmHg; DBP: 79.4 ± 0.60 mmHg) ($P = 0.001$). The mean plasma levels of calcium (10.6 ± 0.17 mg/dl), creatinine (1.83 ± 0.068) and glucose (134.6 ± 4.00 mg/dl) were significantly higher in hypertensive patients compared with the non-hypertensive individuals (calcium: 9.1 ± 0.10 mg/dl; creatinine: 0.82 ± 0.033 mg/dl; glucose: 100.5 ± 1.87 mg/dl) ($P = 0.001$).

Conclusion: Hypertension is associated with significant increase in the plasma levels of calcium, creatinine and glucose in hypertensive patients compared with the non-hypertensive individuals which indicates a significant role of these parameters in the pathogenesis of hypertension and the paramount importance of their measurement in the management of hypertension.

Keywords: Calcium, Creatinine, Glucose, Hypertension

1. INTRODUCTION

Hypertension, also known as high or increased blood pressure, is a disorder in which the blood arteries have an abnormally high pressure for an extended period of time requiring the heart to work harder to pump blood [1], [2]. According to [3], hypertension is defined as a systolic blood pressure of 140 mmHg or higher and/or a diastolic blood pressure of 90 mmHg

or higher. However, in 2017, [4] defined hypertension as systolic blood pressure of 130 mmHg or higher and a diastolic blood pressure of 80 mmHg or higher. The definition and categories of hypertension have been evolving over years, but there is a consensus that persistent BP readings of 140/90mmHg or more should undergo treatment with the usual therapeutic target of 130/80mmHg or less [5]. Pre-hypertension is defined as a grey area between 120–139 mmHg systolic BP and 80–89 mmHg diastolic BP. Despite the fact that pre-hypertension is not a medical condition in and of itself, pre-hypertensive people are more likely to acquire hypertension [6].

Hypertension has been steadily rising in recent decades, particularly in low- and middle-income nations, posing a severe danger to world health [1]. Hypertension is the leading risk factor for cardiovascular disease (CVD) globally, which if left uncontrolled, can lead to a heart attack, heart hypertrophy, and finally heart failure [3], [7]. Stroke, kidney failure, blindness, arterial aneurysm, blood vessel rupture, and cognitive impairment are all possible outcomes of hypertension [3], [5]. Along with its widespread incidence, hypertension is a major public health concern [5]. High blood pressure is responsible for approximately 7.5 million fatalities per year, or 12.8 percent of all deaths globally. In 2025, the number of adults with hypertension is expected to rise to 1.56 billion [8]. A moderate increase in arterial blood pressure reduces life expectancy [3].

Although dietary and lifestyle adjustments can improve blood pressure regulation and reduce the risk of associated health issues, pharmacological treatment may be required in patients for whom lifestyle changes are unsuccessful or insufficient [5]. Several hereditary and lifestyle factors, including nutrition, influence blood pressure. In this regard, sodium is an important mineral that, in addition to its functions in fluid balance, action potential generation, digestive secretions, and nutrient absorption, also plays a key role in blood pressure regulation, with lower sodium intake linked to lower systolic and diastolic blood pressure [5]. As a result, excessive dietary salt (sodium chloride) is considered an established risk factor for hypertension, regardless of body weight, sex, or age. Higher potassium intake or supplementation, when combined with salt decrease, has been found to lower blood pressure in people who are hypertensive [5], [9]. Hypertension is a silent killer since no symptoms are visible in the early stages until a serious medical emergency occurs, such as a heart attack, stroke, or chronic renal disease [10]. Although the majority of hypertensive individuals are asymptomatic, some suffer from headaches, lightheadedness, vertigo, impaired vision, or fainting episodes [10]. Hypertension is caused by a combination of variables. These variables differ from one country to the next. Because most people are unaware of high blood pressure, it can only be detected through measures [11].

People living with hypertension with elevated or low levels of calcium, creatinine and blood glucose are at risk of developing myocardial contractility, arrhythmias, chronic kidney disease, diabetes and other co-morbidities [12], [13], [14]. Hence, the need to assess the plasma levels of calcium, creatinine and blood glucose among hypertensive patients in Edo State, Nigeria in order to ascertain whether plasma levels of these parameters are associated with hypertension in this population which will provide improved insight on proper management of hypertensive conditions.

2. MATERIALS AND METHODS

2.1 Study Design and Setting

This was an analytical cross-sectional study conducted at Edo State University Teaching Hospital Auchi in Etsako West L.G.A, Edo State, Nigeria. Etsako west local government area is situated in Edo state and has its headquarters in the town of Auchi. Six clans make up Etsako west LGA and these include Uzairue, Auchi, South Ibie, Anwain, Jagbe and Aviele. The population of Etsako west LGA is estimated at 294,717 inhabitants with the area hosting members of different tribes which include the Agbede ethnic sub-division (Nigeria Media, 2018).

2.2 Study Population

A total of 300 individuals aged ≥ 20 years were randomly recruited in this study after a written informed consent was sought and obtained upon counselling on the purpose and requirements of the study. The test group comprised 150 hypertensive individuals while the control participants consisted of 150 non-hypertensive individuals who were matched for age. Hypertension was defined as a blood pressure of $\geq 140/90$ mmHg[3].

2.3 Sample Size Determination

The sample size was calculated using the Fisher's formula, with prevalence of 20.2% of hypertension in Edo State with an error margin (d) of 0.05 and a 95% confidence interval [15].

The sample size, $N = Z^2p(1-p)/d^2$

N = required minimum sample

Z^2 = standard normal deviation (1.96)

p = prevalence (20.2%); $20.2 / 100 = 0.202$

$d^2 = 0.05$ the inverse of 95% confidence limit.

$$N = \frac{Z^2p(1-p)}{d^2} = \frac{1.96^2 \times 0.202 \times (1-0.202)}{0.05^2} = 247$$

A non-response rate of 10% of 247 = 24.7 approximately 25

$$N = 247 + 25 = 272$$

The sample size for this study was 272 which was rounded up to 300.

2.4 Inclusion Criteria

Hypertensive and non-hypertensive individuals aged 20 years and above.

2.5 Exclusion Criteria

Individuals below 20 years old, those with history of diabetes, cardiovascular disease, hepatic disorder, cancers, human immunodeficiency virus (HIV) infection, renal dysfunction and individuals who withheld their consent. The exclusion was done after the filled questionnaire has been reviewed.

2.6 Informed Consent

A written informed consent was sought and obtained from the participants before being recruited in the study after the nature and scope of the study have been explained to them.

2.7 Ethical Consideration

Ethical approval for the study was obtained from the Ethics Committee of Edo State University Teaching Hospital (EDSUTH) Auchi.

2.8 Sample Collection and Analysis

Individuals with informed consent had their blood pressure taken using an electronic sphygmomanometer at the right arm in a sitting position after adequate relaxation. Two sets of blood pressure were taken and the average of the reading was taken as the systolic and diastolic blood pressure of each participant. Systolic and diastolic blood pressures greater than or equal to 140mmHg and 90mmHg respectively were regarded as high blood pressure [3]. Information on socio-demographic and medical history was obtained using well-structured questionnaire.

Five milliliters of blood sample was collected from each participant using standard venipuncture technique. The blood samples were added in to lithium heparin container for calcium and creatinine assay and in to fluoride oxalate container for blood glucose assay. The blood samples were centrifuged at 4000rpm for 5 minutes to obtain the plasma. The separated plasma samples were put into a plain container and stored at -20°C prior to analysis. O'Cresolphthalein method, Jaffe-Slot alkaline Picrate method and glucose-oxidase method with Randox kit were used for the quantitative determination of plasma levels of calcium, creatinine and glucose respectively.

2.9 Statistical Analysis

The data for this study was analyzed using statistical software SPSS version 16 (SPSS Inc. Chicago, Illinois). The results were expressed as mean plus or minus standard deviation (mean \pm SD) in tabular form. Independent student's t-test was used to compare the mean differences between the two groups. P-value < 0.05 was taken to be statistically significant.

3. RESULTS

As shown in table 1, the mean age of hypertensive patients (52.2 ± 1.18 years) was significantly higher compared with the non-hypertensive control group (35.1 ± 0.91 years) ($p = 0.001$). The systolic (156.2 ± 1.18 mmHg) and diastolic blood pressure (95.8 ± 0.70 mmHg) were significantly higher in hypertensive patients compared with the non-hypertensive patients (SBP: 118.6 ± 0.64 mmHg; DBP: 79.4 ± 0.60 mmHg) ($p = 0.001$).

Table 1. Comparison of mean age, SBP and DBP in hypertensive and non-hypertensive patients.

Variables	Hypertensives (N = 150)	Non- hypertensives (N = 150)	t-value	P-value
Age (years)	52.2±1.18	35.1±0.91	11.474	0.001*
SBP (mmHg)	156.2±1.18	118.6±0.64	27.996	0.001*
DBP (mmHg)	95.8±0.70	79.4±0.60	17.714	0.001*

Values are represented as mean±SD, SBP = systolic blood pressure, DBP = diastolic blood pressure, **P* = 0.001 significant.

As presented in table 2, the mean values of plasma calcium (10.6±0.17mg/dl), creatinine (1.83±0.068mg/dl) and glucose(134.6±4.00mg/dl) were significantly higher in hypertensive patients compared with the non-hypertensive control patients (calcium: 9.1±0.10mg/dl; creatinine: 0.82±0.033mg/dl; glucose: 100.5±1.87mg/dl; p = 0.001).

Table2. Comparison of mean values of plasma calcium, creatinine and glucose in hypertensive and non-hypertensive patients.

Parameters	Hypertensives (N = 150)	Non- hypertensives (N = 150)	t-value	P-value
Calcium (mg/dl)	10.6±0.17	9.1±0.10	7.639	0.001*
Creatinine (mg/dl)	1.83±0.068	0.82±0.033	13.356	0.001*
Glucose (mg/dl)	134.6±4.00	100.5±1.87	7.741	0.001*

Values are represented as mean±SD, **P* = 0.001 significant

4. DISCUSSION

Hypertension, often known as high or increased blood pressure, is a disorder in which the blood vessels have an abnormally high pressure for an extended period of time [2]. Persistent hypertension is a primary cause of chronic kidney failure and a risk factor for stroke, myocardial infarction, heart failure, and arterial aneurysm [3, 4]. The research work was centered on assessing the plasma levels of calcium, creatinine and glucose among hypertensive patients in Etsako West L.G.A, Edo State, Nigeria.

This present study showed significant increase in the mean age of hypertensive patients when compared with non-hypertensive patients. The mean age of 52.2 ± 1.18 years recorded in hypertensive patients recruited in this study is almost the same as the mean age of 50.0 ± 10.0 years reported by [16] in a similar study at Wolaita Sodo University Teaching and Referral Hospital, SNNPR, Ethiopia. Age is regarded as an unmodifiable risk factor for hypertension. Age has been positively associated with increased blood pressure levels [17]. Increase in blood pressure with age is mostly associated with structural changes in the arteries and especially with large artery stiffness [18].

Our study also showed significant increase in the systolic and diastolic blood pressures of hypertensive patients when compared with the non-hypertensive control patients. In hypertensive patients the pressure against the blood vessel walls in the body is consistently too high [5]. This finding agrees with previous studies by [1] and [2].

Our findings showed a significantly increased plasma calcium concentration in hypertensive patients compared with their non-hypertensive counterparts. This could be as a result of an overactive parathyroid gland leading to excessive secretion of parathyroid hormone and the attendant increase in plasma calcium level [19]. Total calcium has earlier been suggested as a risk factor for hypertension [20]. Hypercalcemia is associated with vascular resistance and vasoconstriction mediated via the direct effect of calcium on vascular smooth muscle as well as the indirect effect of calcium induced hypercatecholaminemia which contributes to the development of hypertension [19], [21]. This is consistent with the findings of [20] and [21]. However, in contrast to our finding, [16] in a similar study at Ethiopia reported a significant decrease in serum calcium levels in hypertensive patients compared with the non-hypertensive individuals. This could be attributed to difference in sample size and study design.

Our study found significantly elevated levels of plasma creatinine in hypertensive patients compared with the non-hypertensive control group. High blood pressure occurs when the force of the blood pushing on the walls of the arteries is too high. This can damage or weaken the blood vessels around the kidneys, impacting kidney function and leading to elevated creatinine levels [12]. Elevated plasma creatinine level is an indicator of chronic renal disease and also associated with inadequate treatment of high blood pressure [22]. Elevated plasma creatinine has been associated with increased mortality in hypertensive persons, the elderly, and patients of myocardial infarction or stroke where cardiovascular disease is the major cause of death [12]. This finding correlates with previous studies by [23] and [22] in a similar study setting.

Our study also showed a significant increase in blood glucose concentration in hypertensive patients compared with the non-hypertensive control individuals. People with high blood pressure usually have insulin resistance and have an increased risk of diabetes compared to those with normal blood pressure due to bodily processes that link both conditions such as inflammation, oxidative stress, activation of the immune system, thickening of the blood vessels and obesity [24]. This may therefore account for the significant increase in levels of

blood glucose in the hypertensive group. Elevated blood sugar decreases the elasticity of blood vessels and causes them to narrow, impeding blood flow, This can lead to a reduced supply of blood and oxygen, increasing the risk of high blood pressure and damage to large and small blood vessels [25].Our finding agrees with previous studies by [24] and [26].

5. CONCLUSION AND RECOMMENDATIONS

The present study showed a significant increase in the plasma levels of calcium, creatinine and blood glucose in hypertensive patients compared with the non-hypertensive individuals which indicates a significant role of these parameters in the pathogenesis of hypertension and the A

Regular screening for the plasma level of calcium, creatinine and blood glucose should be considered as routine tests for hypertensive patients. Adequate awareness of the risk of ignoring these parameters in hypertensive patients should be ensured.

CONSENT

A written informed consent was sought and obtained from the participants before being recruited in the study after the nature and scope of the study have been explained to them.

ETHICAL APPROVAL

Ethical approval for the study was obtained from the Ethics Committee of Edo State University Teaching Hospital (EDSUTH) Auchi.

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

REFERENCES

[1] Mills KT, Bundy JD, Kelly TN, Reed JE, Kearney PM, Reynolds K, Chen J, He J. Global disparities of hypertension prevalence and control: a systemic analysis of population-based studies from 90 countries. *Circulation*. 2016; 134: 441-450.

[2] Muntner P, Carey RM, Gidding S, Jones DW, Taler SJ, Wright Jr, JT, Whelton, PK. Potential US population impact of the 2017 ACC/AHA high blood pressure guideline. *Circulation*. 137: 109-118.

[3] World Health Organization (2013). A global brief on hypertension: silent killer, global public health crisis: World Health Day 2013. World Health Organization.

- [4] Whelton PK, Carey RM, Aronow WS, Casey DE, Collins KJ, Himmelfarb DC, DePalma SM, Gidding S, Jamerson KA, Jones DW, MacLaughlin EJ. ACC/AHA/AAPA/ABC/ACPM/AGS/APhA/ASH/ASPC/NMA/PCNA Guideline for the Prevention, Detection, Evaluation, and Management of High Blood Pressure in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Journal of the American College of Cardiology*. 2017; 71(19):127-248.
- [5] Iqbal S, Klammer N, Ekmekcioglu C. The effect of electrolytes on blood pressure. A brief summary of meta-analyses. *Nutrients*. 2019; 11(6): 1362.
- [6] Kumar MR, Shankar R, Singh S. Hypertension among the adults in rural Varanasi: a cross-sectional study on prevalence and health seeking behavior. *Indian Journal of Preventive and Social Medicine*. 2016;47(1-2): 78–83.
- [7] Forouzanfar MH, Liu P, Roth GA, Ng M, Biryukov S, Marczak L, Alexander L, Estep K, Abate KH. Global burden of hypertension and systolic blood pressure of at least 110 to 115mmHg. *Journal of the American Medical Association*. 2015; 317:165–182.
- [8] Tabrizi JS, Sadeghi-Bazargani H, Farahbakhsh M, Nikniaz L, Nikniaz Z. Prevalence and associated factors of prehypertension and hypertension in Iranian population: the lifestyle promotion project (LPP). *PloS one*. 2016;11(10): e0165264. <https://doi.org/10.1371/journal.pone.0165264>.
- [9] Warren HR, Evangelou E, Cabrera CP, Gao H, Ren M, Mifsud B, Ntalla I, Surendran P, Liu C, Cook JP, Kraja AT, Drenos F, Loh M, Verweij N, Marten J. International Genomics of Blood Pressure (iGEN-BP) Consortium. *Nature Genetics*. 2017; 49(3):403-415.
- [10] Prabakaran J, Vijayalakshmi N, VenkataRao E. Prevalence of hypertension among urban adult population (25–64 years) of Nellore. *International Journal of Research & Development of Health*. 2013; 1(2): 42–49.
- [11] Rani R, Mengi V, Gupta RK, Sharma HK. Hypertension and its risk factors—a cross sectional study in an urban population of a North Indian District. *Public Health Research*. 2015; 5(3): 67–72.
- [12] Yadav SC. Evaluation of serum creatinine in hypertensive patient. *Journal of Nobel Medical College*. 2016; 5(2): 56-59.
- [13] Uehara A, Kita Y, Sumi H, Shibagaki Y. Proton-pump inhibitor-induced severe hypomagnesemia and hypocalcemia are clinically masked by thiazide diuretic. *Internal Medicine (Tokyo, Japan)*. 2019;58(15): 2201-2205. <https://doi.org/10.2169/internalmedicine.2608-18>.
- [14] Duque EJ, Elias RM, Moysés RMA. Parathyroid Hormone: A Uremic Toxin. *Toxins(Basel)*. 2020; 12(3):189.
- [15] Egbi OG, Ahmed SD, Madubuko R. Prevalence and biosocial determinants of hypertension in a rural population in Edo State, Southern Nigeria. *African Journal of Primary Health Care & Family Medicine*. 2021; 13(1): e1-e7.

- [16] Haile B, Wolde M, Gebregziabihier T. Assessment of fasting blood glucose, serum electrolyte, albumin, creatinine, urea and lipid profile among hypertensive participants at wolaitasodo university teaching, SNNPR, Ethiopia. *BioRxiv*. 2020; 10(27): 356873.
- [17] Alhassan Y, Kwakye AO, Dwomoh AK, Baah-Nyarkoh E, Ganu VJ, Appiah B, Kretchy IA. Determinants of blood pressure and blood glucose control in patients with co-morbid hypertension and type-2 diabetes mellitus in Ghana: a hospital based cross-sectional study. *PLOS Global Public Health*. 2022; 2(12): e0001342. <https://doi.org/10.1371/journal.pgph.0001342>.
- [18] Choi J, Lee M, Lee JK. Correlates associated with participation in physical activity among adults: a systematic review of reviews and update. *BMC Public Health*. 2017; 17: 356.
- [19] Villa-Etchegoyen C, Lombarte M, Matamoros N, B elizan JM, Cormick G. Mechanisms involved in the relationship between low calcium intake and high blood pressure. *Nutrients*. 2019; 11(5): 1112. <https://doi.org/10.3390/nu11051112>.
- [20] Wu D, Chen Y, Guan H, Sun Y. Association of abnormal serum electrolyte levels with hypertension in a population with high salt intake. *Public Health Nutrition*. 2019; 22(9), 1635-1645. <https://doi.org/10.1017/S136898001900>.
- [21] Hua Y, Liu HL, Sun JY, Kong XQ, Sun W, Xiong YQ. Association between serum calcium and the prevalence of hypertension among US adults. *Frontiers in Cardiovascular Medicine*. 2021; 8: 719165. <https://doi.org/10.3389/fcvm.2021.719165>.
- [22] Akpotaire PA, Seriki SA. Assessment and correlation of serum urea and creatinine levels in normal, hypertensive, and diabetic persons in Auchi, Nigeria. *Arch PatholClin Res*. 2023; 7: 007-016. <https://doi.org/10.29328/journal.apcr.1001.035>.
- [23] Shekhar CY. Elevation of serum creatinine in hypertensive patient. *Journal of Noble Medical College*. 2016; 5(2): 56-59.
- [24] Petrie JR, Guzik TJ, Touyz RM. Diabetes, hypertension and cardiovascular disease: clinical insights and vascular mechanisms. *The Canadian Journal of Cardiology*. 2018; 34(5): 575-584. <https://doi.org/10.1016/j.cjca.2017.12.005>.
- [25] Yan Q, Sun D, Li X, Chen G, Zheng Q, Li L, Gu C, Feng B. Association of blood glucose level and hypertension in Elderly Chinese Subjects: a community based study. *BMC Endocrine Disorders*. 2016; 16(1): 40.
- [26] Ahn SK, Lee JM, Ji SM, Kim KH, Park JH, Hyun MK. Incidence of Hypertension and fasting blood glucose from real-world data: retrospective cohort for 7-years follow-up. *International Journal of Environmental Research and Public Health*. 2021; 18(4): 2085. <https://doi.org/10.3390/ijerph18042085>.

UNDER PEER REVIEW