

Original Research Article

Seed yield, Seed attributes and economics of alfalfa seed production as influenced by Integrated nutrient management during *kharif* season under southern dry zone of Karnataka

ABSTRACT

The field experiment was carried out at the farm field of KrishiVigyan Kendra, Konehalli, Tiptur taluk, Tumkuru, Karnataka during *Kharif* season of 2017-18 to examine the effect of integrated nutrient management on seed yield, seed attributes and net benefit of alfalfa. The experiment was laid out Randomized complete block design (RCBD) comprising nine treatments viz T₁ : RDF (25:50:25 kg NPK/ha + 10 t/ha FYM); T₂ : 75% RDF + 25% N through FYM; T₃ : 75% RDF + 25% N through Vermicompost; T₄ : 75% RDF + 25% N through Poultry manure; T₅ : 50% RDF + 25% N through FYM + Rhizobium + PSB+VAM ; T₆ : 50% RDF + 25% N through Vermicompost+ Rhizobium + PSB+VAM; T₇ : 50% RDF + 25% N through Poultry manure+ Rhizobium + PSB+VAM ; T₈ : RDF + *Rhizobium* + PSB+VAM ; T₉ : 10 t/ha FYM + 100% N through FYM and replicated four times. The results of experiments revealed that the plants supplied with of 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM shown highest fresh herb yield at first harvest (160.65 q/ha), herb yield after harvest of seeds (187.50q/ha), 50% flowering (35.00 days), pod initiation (48.50days), pod maturation (73.05days), while, lower fresh herb yield, pod initiation, pod maturation were recorded when 10 t/ha FYM + 100 % N through FYM was applied. The maximum pod (56.78 %), number of pods (228.75/plant), pod yield (19.90/plant), pod length (6.25 mm), pod weight (87.61 mg), seed set (56.50 %), number of seeds (5.50/pod), test weight (3.33g), seed yield per hectare (223.38kg/ha) and highest net return per hectare were recorded with 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM. while, poor performance with regard to these parameter were recorded with application of 10 t/ha FYM + 100 % N through FYM. Therefore, application of 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM may be recommended for commercial seed production in Alfalfa during *kharif* season under southern dry zone of Karnataka.

Keywords: INM, Kharif, nutrient, PSB, *Rhizobium*, seed and yield

1. INTRODUCTION

Alfalfa (*Medicago sativa* L.) 'Queen of the fodder crops' is an important fodder crop of India, belonging to family Fabaceae and growing extensively as an annual or perennial crop with supply of green fodder continuously for 2-3 years from the same field. It was introduced into India during 1900 from North-West [1] and cultivated in Uttar Pradesh, Punjab, Gujarat, Haryana, Tamil Nadu, Maharashtra and Karnataka in the country. It is growing approximately one million hectare of area and produced 60-130 tonnes of green fodder per hectare per year with 186 - 280 kg/ha of seed yield in India. It is locally known as 'Kuduremasale' in Karnataka.

The nutritive content of palatable fodder about 13.3 - 26.6% crude protein, calcium (0.92 - 2.9 %), phosphorus (0.14 - 0.66 %), carotene (9.27 mg/ 100g), fibre (20 -30 %) and vitamin A and C [4]. It also contains stachydrine as medicinal alkaloid and used for digestive, laxative, diuretic and treating against for blood pressure, arthritis, hair loss, acidity and dropsy. The soil health and environment are deteriorating due to continuous use of chemical fertilizers, which can be prevented by using the judicious combination of different sources of nutrients such as use of organic manures and bio-fertilizers along with balanced

inorganic fertilizers, one of the eco-friendly approaches, which can be enhancing the productivity and sustainability [13]. By considering the importance of crop and role of integrated nutrient management, the experiment was conducted to study the seed yield, quality and economics of alfalfa seed production as influenced by INM during *kharif*.

2. MATERIALS AND METHOD

The field experiment was carried out at the farm field of Krishi Vigyan Kendra, Konehalli, Tiptur taluk, Tumkuru district under southern dry zone of Karnataka during *Kharif* season 2017-18 on seed yield, quality and economics of alfalfa seed production as influenced by integrated nutrient management. The nine treatments and four replication were laid out with Randomized complete block design (RCBD). Plots were prepared with size of 4.32sq.m (2.4 m x 1.8 m) area of red sandy loam soil. Alfalfa seeds of T-9 variety were treated with *Rhizobium meliloti* bio-fertilizers, applied recommended dose of fertilizers and organic manures with bio-fertilizers viz., *Phosphoroussolubilizingbacteris (PSB)* and *vascular arabascularmycorrhiza (VAM)* as per treatments. The seeds were sown during *kharif* season with 30 cm x 15 cm spacing.

Treatment details

- T₁ : RDF (25:50:25 kg NPK/ha)+ 10 t/ha FYM
- T₂ : 75% RDF + 25% N through FYM
- T₃ :75% RDF + 25% N through Vermicompost
- T₄ :75% RDF + 25% N through Poultry manure
- T₅ : 50% RDF + 25% N through FYM + Rhizobium + PSB+VAM
- T₆ :50% RDF + 25% N through Vermicompost+ Rhizobium + PSB+VAM
- T₇ :50% RDF + 25% N through Poultry manure+ Rhizobium + PSB+VAM
- T₈ : RDF + *Rhizobium* + PSB+VAM
- T₉ : 10 t/ha FYM + 100% N through FYM

Table 1: NPK content of Poultry manure, Vermi-compost and Farm yard manure for NPK content before conducting the experiment

Organic manures	N content (%)	Pcontent (%)	Kcontent (%)
Poultry manure	2.10	1.35	1.76
Vermicompost	1.60	0.86	0.98
Farm yard manure	0.95	0.62	0.75

The first crop was harvested at 60 days after sowing and allowed it for further growth, flowering and to reach physiological maturity stage for obtaining the seeds. The observations on herb and seed yield were recorded using five plants per plot. These five plants were randomly selected by avoiding the border plants and labelled for recording the observations at harvest stage. The recorded data were analysed by suitable statistical analysis as per the procedure outlined by [14]. The results were compared at 5 per cent probability using Fischer's test.

Benefit cost ratio (B:C ratio) :The benefit cost ratio was calculated by using the below mentioned formula

$$\text{B:C ratio} = \frac{\text{Gross income (Rs. ha}^{-1}\text{)}}{\text{Total cost of cultivation (Rs. ha}^{-1}\text{)}}$$

3. RESULTS AND DISCUSSION

3.1 Fresh weight of plants and herb yield of Alfalfa affected by INM during *kharif* season

The fresh weight of plant, herb yield/plot and herb yield/hawere recorded during *kharif* season at first harvest and herb yield after harvest of seeds are presented in Table 2. The plants supplied with 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM during *kharif* season was recorded maximum fresh weight (133.11g/plant & 121.49g/plant), fresh herb yield (3.47kg/plot & 4.05kg/plot) and fresh herb yield (160.65q/ha & 187.50q/ha) at first harvest of herb and herb yield after harvest of seeds respectively, which was *at par* with the plants supplied with 50 % RDF + 25 % N through poultry manure + *Rhizobium* + PSB + VAM. Whereas application of 10 t/ha FYM + 100 % N through FYM during *kharif* season were recorded least fresh weight/plant, fresh herb yield/plot and fresh herb yield/ha at first harvest of herb and herb yield after harvest of seeds.

The highest fresh weight of plant and fresh herb yield might be due to that, vermin-compost is known to produce favorable effect on chemical, biological and physical factors in soil, that determines productivity and fertility status of soil and supply the nutrients in their available form, enhance the microbial population and provides sufficient energy for them to remain active [15]. It also provides both the vital macro-nutrient and micro-nutrients such as N, P, K, Ca, Mg, besides, *Rhizobium* has been increased the availability of N and helped in the synthesis of tryptophan as resulted in obtained maximum herb yield. These results are in agreement with [20] in alfalfa, [9, 18, 8] in ashwagandha and [9] in davana.

Table 2: Fresh weight of plants and herb yield of Alfalfa affected by INM during *kharif* season

Treatment	Fresh plant weight (g)		Fresh herb yield / plot (kg)		Fresh herb yield / hectare (q)		Cumulative herb yield
	1 st Harvest of herb	Herb weight after harvest of seeds	1 st Harvest of herb	Herb yield after harvest of seeds	1 st Harvest of herb	Herb yield after harvest of seeds	
T ₁	115.96	106.08	3.34	3.93	154.63	181.94	336.57
T ₂	103.37	96.58	3.23	3.81	149.54	176.39	325.93
T ₃	112.15	102.85	3.30	3.89	152.78	180.09	332.87
T ₄	110.15	100.06	3.27	3.87	151.39	179.17	330.56
T ₅	124.56	113.50	3.41	3.99	157.87	184.72	342.59
T ₆	133.11	121.49	3.47	4.05	160.65	187.50	348.15
T ₇	129.40	116.97	3.45	4.03	159.72	186.57	346.29
T ₈	119.87	109.68	3.38	3.60	156.48	166.67	323.15
T ₉	99.02	93.79	3.15	3.76	145.83	174.07	319.90
F- test	*	*	*	*	*	*	*
S.Em±	3.10	2.61	0.017	0.012	1.01	0.81	2.00
CD at 5 %	9.05	7.64	0.05	0.035	2.95	2.36	5.83

3.2 Effect of flowering characters by INM in Alfalfa during *kharif* season

Flowering characters such as days taken for flowering, pod initiation, pod maturation and number of flowers/plant were recorded during *kharif* season are presented in Table 3. The plants supplied with 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM during *kharif* season was recorded in early flowering (27 days), 50% flowering (35 days), pod initiation (48.5 days), pod maturation (73.05days) and highest number of flowers per plant (405) were recorded, which was *on par* with the application of 50 % RDF + 25 % N through poultry manure + *Rhizobium* + PSB + VAM in pod length. Whereas, application of 10 t/ha FYM + 100 % N through FYM during *kharif* season were recorded in delayed flowering, 50% flowering, pod initiation, pod maturation and least number of flowers.

This might be due to the integrated application of inorganic fertilizers and organic manure along with biofertilizers showing similar effect of growth regulators and provide in better uptake of nutrients by the plants as a result in better reproductive growth of the plants. The similar reports were found with [6] in Alfalfa, [12] in peppermint, [16] in Kalmegh, [2, 5] in chamomile.

Table 3: Effect of flowering characters by INM in Alfalfa during *kharif* season.

Treatment	No. of days for flowering	No. of days for 50% flowering	No. of flowers/plant	No. of days for pod initiation	No. of days for pod maturation
T ₁	28.55	36.10	386.50	50.13	75.30
T ₂	29.13	37.85	379.25	52.25	77.63
T ₃	28.60	36.10	385.75	50.13	75.35
T ₄	29.00	37.20	382.50	51.75	76.03
T ₅	27.55	35.60	396.63	49.10	74.28
T ₆	27.00	35.00	405.00	48.50	73.05
T ₇	27.10	35.50	399.25	48.75	73.75
T ₈	27.75	35.80	389.63	49.38	74.38
T ₉	29.38	38.00	376.00	53.25	78.25
F- test	*	*	*	*	*
S.Em±	0.15	0.18	2.78	0.12	0.38
CD at 5 %	0.44	0.54	8.14	0.34	1.12

3.3 Effect of pod quality by INM in Alfalfa during *kharif* season

The pod quality such as pod set per cent, number of pods/plant, number of filled pods, number of seeds/pod, pod yield, pod weight and pod length were recorded during *kharif* season are presented in Table 4. The application of 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM during *kharif* season were recorded maximum pod set per cent (56.78%), number of pods (228.75/plant), number of filled pods (129.25/plant), pod yield (19.90/plant), pod weight (87.61mg) and pod length (6.25mm). Whereas, application of 10 t/ha FYM + 100 % N through FYM during *kharif* season were recorded with minimum pod set per cent, number of pods/plant, number of filled pods/plant, pod yield/plant, pod weight and pod length. The good pod quality could be due to application of *Rhizobium*, PSB and VAM enhances vegetative growth, better of photosynthetic and metabolites as resulted in early flowering, pod initiation and which improved the pod characters. The results are similar with the works of [7] in Ambrette.

Table 4: Effect of pod quality by INM in Alfalfa during *kharif* season

Treatment	Pod set (%)	No. of pods per plant	No. of filled pods per plant	Pod yield per plant (g)	Pod length (mm)	Pod weight (mg)
T ₁	54.98	212.50	111.50	18.02	5.90	85.38
T ₂	53.92	204.50	101.50	16.95	5.63	84.35
T ₃	55.02	211.95	111.50	18.00	5.85	85.40
T ₄	54.58	208.75	106.55	17.56	5.80	85.25
T ₅	56.16	222.25	121.75	19.10	6.00	86.15
T ₆	56.78	228.75	129.25	19.90	6.25	87.61
T ₇	56.43	225.50	127.50	19.50	6.15	86.03
T ₈	56.14	220.00	119.25	18.54	5.98	85.95
T ₉	53.66	201.75	98.00	16.58	5.58	83.75
F- test	*	*	*	*	*	*
S.Em±	0.22	2.12	2.10	0.25	0.07	0.41
CD at 5 %	0.64	6.21	6.13	0.73	0.22	1.25

3.4 Effect of seed characters and seed yield by INM in Alfalfa during *kharif* season

The Effect of seed characters and seed yield by INM during *kharif* season are presented in Table 5. The plants supplied with 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM during *kharif* season was recorded maximum seed set per cent (56.50%), number of seeds (5.50/pod), test weight (3.33g) and seed yield (2.07g/plant, 48.25g/plot & 223.38kg/ha). Whereas, application of 10 t/ha FYM + 100 % N through FYM during *kharif* season was recorded minimum seed set per cent, number of seeds/pod, test weight, seed yield. This might be due to adequate availability of nutrients sources at different growth stages through the integrated application of inorganic, organic and bio-fertilizers and sequential metamorphosis from source to sink. The similar results are findings with [17] in isabgol and also increased uptake of nitrogen by *rhizobium* fixation of atmospheric nitrogen. Besides, the optimum supply of nutrient could have helped in increased plant metabolic activity and source to sink relationship, thus, higher the yield by more number of seeds. Similar results were found with [3] in Ashwagandha.

Table 5: Effect of seed characters and seed yield by INM in Alfalfa during *kharif* season

Treatment	Seed set per cent	No. of seeds per pod	Test weight (g)	Seed yield per plant (g/pl)	Seed yield per plot (g/plot)	Seed yield per hectore (kg/ha)
T ₁	52.47	4.70	3.16	1.95	43.63	201.99
T ₂	49.63	4.45	3.10	1.92	42.10	194.06
T ₃	52.61	4.71	3.14	1.95	43.53	201.34
T ₄	51.04	4.65	3.11	1.94	42.38	196.20
T ₅	54.78	5.20	3.20	2.00	46.25	214.12

T ₆	56.50	5.50	3.33	2.07	48.25	223.38
T ₇	56.04	5.38	3.24	2.03	47.00	217.59
T ₈	54.20	4.98	3.18	1.97	45.75	211.80
T ₉	48.57	4.35	3.09	1.91	41.38	191.57
F- test	*	*	*	*	*	*
S.Em±	0.24	0.10	0.42	0.02	0.64	2.75
CD at 5 %	0.70	0.29	NS	0.06	1.87	8.03

3.5 Effect of integrated nutrient management on economics of seed production in Alfalfa during *kharif* season

The economics of seed production in Alfalfa during *kharif* season are presented in Table 6. The plants supplied with 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM during *kharif* season was obtained highest net return (Rs. 65,763/ha) and obtained the maximum cost benefit ratio (2.12) during *kharif* season with the application of 50 % RDF + 25 % N through poultry manure + *Rhizobium* + PSB + VAM. Whereas, the lowest net return (Rs.51,824/ha) and B:C ratio (1.92) was obtained with the application of 10 t/ha FYM + 100 % N through FYM during *kharif* season. The highest net return and B:C ratio in seed production could be due to enhancement of growth and development by supplying of integrated nutrients to meet the required demand of the crop till physiological maturity resulted in increased seed yield. Similar reports were found by [10, 11] in alfalfa.

Table 6: Effect of INM on economics of seed production in Alfalfa during *kharif* season

Treatment	Economics of seed production during <i>kharif</i> season			
	Gross Return (Rs./ha)	Cost of Cultivation (Rs./ha)	Net return (Rs./ha)	B:C ratio
T ₁	114090	57199	56891	1.990
T ₂	109759	57063	52696	1.930
T ₃	113520	58360	55160	1.940
T ₄	110998	56987	54011	1.950
T ₅	120034	57558	62476	2.080
T ₆	124618	58855	65763	2.117
T ₇	121873	57482	64391	2.120
T ₈	118782	58487	60295	2.030
T ₉	108081	56257	51824	1.920

4. CONCLUSION

From the experiment findings of present investigation, it can be concluded that, the application of 50 % RDF + 25 % N through vermicompost + *Rhizobium* + PSB + VAM has recorded the maximum seed yield, good quality and highest profit in seed production. which may be recommended for commercial seed production in of Alfalfa during *kharif* season under southern dry zone of Karnataka

Disclaimer (Artificial intelligence)

Option 1:

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image generators have been used during the writing or editing of this manuscript.

Option 2:

Author(s) hereby declare that generative AI technologies such as Large Language Models, etc. have been used during the writing or editing of manuscripts. This explanation will include the name, version, model, and source of the generative AI technology and as well as all input prompts provided to the generative AI technology

Details of the AI usage are given below:

- 1.
- 2.
- 3.

REFERENCES

1. Ahlawa IPS. Agronomy of Rabi Crops - Alfalfa, Division of Agronomy. Indian Agricultural Research Institute, New Delhi. 2007; pp-1-9.
2. Dastborhan DS, Zehtab-Salmansi S, Nasrollahzadeh S, Tavassoli AR., Effect of plant growth promoting rhizobacteria and nitrogen fertilizer on yield and essential oil of German chamomile (*Matricariachamomilla* L.). Acta. Hort. 2012;9(4): 121-128.
3. Dubey PK, Pandey CS, Shakoorkhanday AB, Gaurav Mishra., Effect of integrated nutrient management on nutrient uptake, protein content and yield of fenugreek, International Journal of Food, Agricultural and Veterinary Sciences. 2012;2(1): 1-12.

4. Khalak A. Seed yield of Lucerne (*Medicago sativa* L.) varieties at different times of sowing and levels of fertilizer. M. Sc. (Agri.) Thesis, University Agricultural Sciences, Bengaluru. 1989,
5. Mohammadreza N, Mohammad M, Houseyn Z. Bahari B. Effect of different levels of nitrogen, phosphorous and potassium fertilizers on some agro morphological and biochemical traits of German chamomile (*Matricariachamomilla* L.). Journal of Medicinal Plant Reseach. 2012; 6(2): 277-283.
6. Pankaniya RM, Tandel BB. Pharrmar JD. Effect of integrated nutrient management on growth, yield, nutrient uptake and economic of Lucerne (*Medicagosative* L.). The Parma Innovation. 2023; 12(6): 1695-1697
7. Rajeswari R, Shakila A. Effect of integrated nutrient management practices on yield characters of ambrette (*AbelmoschusmoschatusMedic.*). Plant Archives. 2015;15(1): 537-540.
8. Raju B, Mohana Chandra CN, Sreeramu BS, Farooqui AA, Chandre Gowda M. Influence of bio-inoculants under graded levels of fertilizer on growth, yield and quality of makoi (*Solanumnigrum*L.). International Journal Science and Nature. 2014;5(2): 246-248.
9. Ravirajshetty G, Kusuma MV, Venkatesha J, Sadashiv Nadukeri. Impact of integrated nutrient management on growth and yield of Davana (*Artemisia pallens*Wall.) under hill zone of Karnataka. International Journal of Development and Research. 2014;4(3): 757-759.
10. Rominger RS, Smith D. Peterson LA. Yield and chemical composition of alfalfa as influenced by nutrient levels. American Journal of Expert Agriculture. 2015;68(4): 573-577.
11. Ruixuan X, Haiming Z, Yongliang Y, Ruixin W, Guibo L, Zhiqiang S, Bademuqiqige and Yingjun Z, Effects of intercropping, nitrogen fertilization and corn plant density on yield, crude protein accumulation and ensiling characteristics of silage corn inter seeded into alfalfa stand. Agriculture. 2022;12:1 -15
12. Sheykholeslami Z, Almdari M, Qanbari S, Akbarzadeh M. Effect of organic and chemical fertilizer on yield and yield components of peppermint (*Mentha piperita* L.). American Journal of Experi. Agriculture. 2015; 6(4): 251-257.
13. Singh R, Munnu Singh, Akshata Srinivas, Prakasa Rao EVS, Puttanna K. Assessment of organic and inorganic fertilizers for growth, yield and essential oil quality of industrially important plant Patchouli (*Pogostemoncablin*), Journal of Essential oil Bearing Plants, 2015;18(1): 1-10
14. Sundararaj N, Nagaraju SS, Venkataramu MN, Jaganath MK. Design and analysis of field experiment. University of Agricultural Sciences, Bengaluru. 1972
15. Sunil JN. Effect of dates of planting and nitrogen levels on growth and yield of kalmegh (*Andrographispaniculata*). M. Sc. (Hort.) Thesis, University of Agricultural Sciences, Dharwad. 2010.
16. Sushma M, Aruna J. Effect of INM on vegetative growth, flowering and fruiting of *Andrographis paniculata*. Universal Journal of Agricultural Resarch. 2014;2(3): 93-96
17. Venkatesha NT, Kattimani KN, Madalageri MB, Hanamashetti SI, Gangadharappa PM, Devappa V. Integrated nutrient management in isabgol (*Plantagoovato*Forsk.). The Asian Jornal of Horticulture. 2007; 25: 123- 127.
18. Vijay K, Vivek KS, Roshni A. Effect of different levels of nitrogen and phosphorous on growth, yield and quality of kalmegh (*Andrographispaniculata*). New Agriculturist, 2015; 26(1): 17-23.

19. Vishal G. Duhan BS. Kalmegh (*Andrographis paniculata*) crop as influenced by the application of farmyard manure (FYM) and inorganic phosphorous in typictorripsamments of Hisar. Journal of Medicinal Plant Research. 2013;7(46): 3337-3342.

20. Zhuanlin Ben, Yuan Li and Huimin Yang. Fertilizer application increases alfalfa yield and crude protein content: roles of fertilizer type, application rate, and environmental conditions. Research Square. 2023: pp 1-4.



Alfalfa crop after I harvest



General view of Alfalfa experiment during kharif season



Harvesting of Alfalfa



Plot treatment



Plot treatment



Flowering stage



Leaf and stem weight of all treatment

UNM



Fig 1. Harvesting stage of seeds

UNDER PEER