

# *Moringa oleifera* Leaf Meal: A Sustainable Approach for Poultry

## Production: A Review

### ABSTRACT

Poultry production is among the fastest ways to address the shortage of animal protein supply and intake in many nations. With the prohibition of antibiotics as growth promoters in the poultry industry due to rising bacterial resistance and growing public concern over health and food safety, the sector has been prompted to seek safe alternatives. This shift has led to a focus on finding natural antibiotic substitutes and optimizing feed management to enhance chicken health and growth. Consequently, phytochemicals have emerged as promising natural alternatives to antibiotics in the poultry sector. *Moringa oleifera* has attracted significant research interest recently as a natural supplement with numerous health advantages for poultry. Known for its antimicrobial, antioxidant, anti-inflammatory, and cholesterol-lowering properties, *Moringa* also aids in activating digestive enzymes in the stomach due to its wealth of bioactive compounds. This review emphasizes the potential impact of *Moringa oleifera* as a natural feed additive on growth performance, nutrient digestibility, blood biochemical profile, antioxidant effects, immune response, and egg production in poultry.

**Keywords:** Body weight; egg production; growth performance; immunity; *Moringa oleifera* leaf meal, nutrient utilization

### **1. INTRODUCTION**

Poultry production is one of the quickest methods to address the deficiency of animal protein supply and consumption in various countries. Energy and protein sources make up 95% of the total feed costs in poultry farming. Plant-based protein feedstuffs play a crucial role in meeting dietary protein needs in many practical poultry rations formulated globally. Rations that include soybean meal and groundnut meal are generally preferred over other protein sources. However, with the rising population and increasing demand, there is a looming shortage of key feed ingredients like maize and soybean (Thirumalaisamy *et al.*, 2016). *Moringa* leaves, which contain a high crude protein content (24-29%), can be used as a feed supplement for poultry (Su and Chen, 2020). They offer several benefits, including promoting growth, and possessing antibacterial, antifungal, antioxidant, and anti-inflammatory properties (Mahanta *et al.*, 2017).

*Moringa oleifera*, often referred to as the "Miracle tree," is rich in essential nutritional bioactive compounds that are vital for both livestock and human health (Anjorin *et al.*, 2010). The *Moringa* genus includes 13 species globally, with two found in India: *Moringa*

*concanensis* Nimmo ex Dalz. and *Moringa oleifera* Lam. *Moringa oleifera* Lam. is recognized in over 80 countries, including India, and is known by more than 200 local names due to its diverse applications (Bhargave *et al.*, 2015).

Synonyms for *Moringa oleifera* include various names in different languages. In Hindi, it is known as Sajana, Lingru, Shajna, Saina, Mungaera, Shaijmal, Segra, and Sahjan. In Punjabi, it is referred to as Sanjina, Soanjana, and Sejana. In Rajasthan, it is called Lal Sahinjano and Lingru. In Sindhi, it is known as Swanjera. In English, common names include Horseradish Tree, Ben Tree, Drumstick Tree, Miracle Tree, Mother's Best Friend, and Gold of the Poor. In Bengali, it is called Sajna or Sajina, in Gujarati as Saragavo, and in Pakistan as Sohangna (Fahey, 2005).

## **2. *Moringa oleifera* Tree**

*Moringa oleifera* is a rapidly growing, drought-resistant tree that belongs to the Moringaceae family and is widely cultivated in tropical and subtropical regions across the globe. It thrives in areas such as India, Africa, Arabia, the Caribbean Islands, South and Central America, Mexico, Hawaii, and throughout Southeast Asia, typically reaching heights of 1.5 to 2.0 meters (Foidl *et al.*, 2001). This species produces a significant amount of biomass, ranging from 43 to 115 tons per hectare annually (Kholif *et al.*, 2016). In terms of leaf production, each tree yields about 1 to 5 kilograms of fresh leaves each year, which translates to an impressive 10,000 to 50,000 kilograms per hectare at a spacing of 1 meter by 1 meter (Sánchez *et al.*, 2006).

*Moringa oleifera* offers numerous benefits, as various parts of the tree—including leaves, fruits, immature pods, and flowers—are edible and commonly included in traditional diets in many tropical and subtropical countries (Anhwange *et al.*, 2004). Three notable non-governmental organizations—"Trees for Life" (2005), "Church World Service," and "Educational Concerns for Hunger Organization" (2004)—have promoted *Moringa oleifera* as a source of "Natural nutrition for the tropics." The leaves of *Moringa oleifera* are particularly nutrient-rich, making the plant a valuable potential food supplement and protein source (Mendieta Araica *et al.*, 2011). Additionally, the leaves are high in zeatin, a cytokinin, along with other growth-promoting compounds such as ascorbates, phenolics, and essential minerals like calcium, potassium, and iron, which enhance crop growth (Anjorin *et al.*, 2010).

### **3. *Moringa oleifera* as a Traditional Medicine**

Ayurveda, a traditional system of medicine, suggests that *Moringa* can help prevent over 300 diseases in humans (Ganguly *et al.*, 2013). Additionally, *Moringa* oil holds significant cosmetic value and is utilized in body and hair care products as a moisturizer and skin conditioner. Its use in skin treatments and ointments dates back to ancient Egyptian times (Fugile, 2001).

### **4. *Moringa oleifera* Leaf as a Therapeutic Use**

*Moringa oleifera* leaves contain various phytochemicals, including sterols, tannins, terpenoids, saponins, alkaloids, phenolics, and flavonoids such as isoquercetin, isothiocyanates, quercetin, and kaempferitrin, as well as glycoside compounds (Jung, 2014). Extracts from both mature and young leaves exhibit strong antioxidant properties, effectively combating free radicals and preventing oxidative damage to key biomolecules, thus offering significant protection against oxidative stress (Patel, 2011). Additionally, *Moringa* leaves have demonstrated sympatholytic and antiviral effects against herpes simplex virus type-1, as well as the ability to lower cholesterol levels, antibacterial properties (Talreja, 2010), antifungal effects (Nickon *et al.*, 2003), hepatoprotective benefits (Pari and Kumar, 2002), and growth promotion alongside antimicrobial effects (Mbikay, 2012).

### **5. Nutritional Role of *Moringa oleifera* Leaf**

One of the most remarkable aspects of *Moringa oleifera* leaves is their rich nutrient profile, which includes vitamins (such as  $\beta$ -carotene, vitamins A, D, E, B-complex, and C), minerals (including calcium, chromium, copper, magnesium, manganese, phosphorus, selenium, and zinc), and amino acids (Nkukwana *et al.*, 2015). The nutritional composition of *Moringa* leaves can vary depending on the region (Anjorin *et al.*, 2010).

Dry *Moringa* leaves are particularly nutrient-dense, containing seven times more vitamin C than oranges, ten times more vitamin A than carrots, seventeen times more calcium than milk, fifteen times more potassium than bananas, twenty-five times more iron than spinach, and nine times more protein than yogurt (Fuglie, 1999; Dhakar *et al.*, 2011). The protein content in *Moringa oleifera* leaves ranges from 20% to 33% on a dry matter basis and includes all essential amino acids. Amaglo *et al.* (2010) reported that dry *Moringa* leaves contain 34.80% ether extract, 31.65% protein, 7.54% fiber, 8.9% moisture, and 6.53% ash

based on dry matter. Additionally, Kakengi *et al.* (2005) found that Moringa leaf meal has a crude protein content of 29.71%, 22.5% crude fiber, 4.38% ether extract, 27.9% calcium, and a minimal tannin content of 1.23 g/kg.

## **6. INCORPORATION OF *MORINGA OLEIFERA* LEAF MEAL IN POULTRY FOR SUSTAINABLE PRODUCTION**

### **A) EFFECT ON GROWTH AND PRODUCTION PERFORMANCE**

Ashong and Brown (2011) found that chicks fed a diet containing 10% Moringa leaf meal exhibited reduced feed intake and weight gain. In contrast, Olugbemi *et al.* (2010) reported that adding 5% *Moringa oleifera* leaf meal to cassava-based diets (20% and 30%) for broilers had no significant effect ( $P > 0.05$ ) on feed conversion ratio (FCR), body weight gain, feed consumption, or the economics of production compared to diets without cassava or Moringa. This lack of effect was attributed to the increased fiber content, which may have hindered nutrient digestibility and absorption. Similarly, Ayssiwede *et al.* (2011) observed that including *Moringa oleifera* leaf meal (MOLM) at levels of 0%, 8.0%, 16.0%, and 24.0% in the diets of growing indigenous Senegal chickens did not adversely affect live body weight, average daily weight gain, FCR, or mortality rates, even at 24% MOLM. However, a significant decrease in feed intake was noted at 16.0% and 24.0% MOLM, while birds fed 8.0% and 16.0% MOLM showed better growth performance, lower feed costs, and improved economic margins. Hassan *et al.* (2016) found that broiler chicks supplemented with 0.3% MOLM had the highest body weight gain and FCR, while Elkloub *et al.* (2015) reported that Japanese quail chicks receiving 0.2% MOLM had the lowest feed consumption ( $P \leq 0.01$ ), the best feed conversion ratio, and a higher European Production Efficiency Index (EPEI) compared to the control group. Khalifa *et al.* (2018) showed that birds on a diet with 5% *Moringa oleifera* meal gained significantly more weight than those on the control diet, while the group with 20% Moringa had the lowest feed consumption ( $P \leq 0.01$ ) and the best feed conversion ratio. Fouad and Raye (2019) demonstrated that birds fed 7g MOL/kg of diet experienced significantly higher ( $P < 0.05$ ) body weight gain and feed intake, along with improved FCR compared to other dietary treatments. Talukdar *et al.* (2020) found that adding *Moringa oleifera* leaf meal at levels of 0.25% to 0.50% to Japanese quail diets improved body weight, weight gain, FCR, and increased gross profit. Baloch *et al.* (2021) reported that 2.5% *Moringa oleifera* leaf meal supplementation led to maximum body weight and feed intake in chickens compared to other groups. The control group had the lowest feed intake,

while the 2.5% Moringa group showed significantly better FCR ( $P < 0.05$ ). Furthermore, Varalakshmi *et al.* (2021) revealed significant increases ( $P < 0.01$ ) in body weight gain, overall feed intake, and improved FCR with 3.0% *Moringa oleifera* leaf meal incorporated into the diet compared to other treatment groups of Japanese quail.

## **B) EFFECT ON NUTRIENT UTILIZATION**

Tijani *et al.* (2016) found that feeding a diet with 15% Moringa leaf meal resulted in significantly higher ( $P < 0.05$ ) digestibility values for crude protein (CP) and ether extract. In contrast, the lowest CP digestibility was observed in broiler chickens consuming a diet containing 20% Moringa leaf meal. This decline in digestibility is likely due to the high proportion of Moringa in the diet, which may lead to an increased concentration of anti-nutritional factors. Furthermore, Fouad and Raye (2019) reported that Japanese quail receiving 7g of *Moringa oleifera* leaf meal per kilogram of diet exhibited the best nutrient digestibility (dry matter, crude protein, crude fiber, ether extract, and nitrogen-free extract) compared to other dietary treatments. Similarly, Siti *et al.* (2019) found that incorporating Moringa leaf powder at levels of 4-6% in the diets of laying hens enhanced dry matter and organic matter digestibility. Harshini *et al.* (2022) noted that adding *Moringa oleifera* leaf powder (at 0%, 5%, 10%, and 15%) to the basal diet of Kadaknath chickens did not result in significant ( $P > 0.05$ ) differences in apparent nutrient digestibility, and the reduced feed intake in the supplemented groups did not negatively impact their nutrient retention.

## **C) EFFECT ON HAEMATO-BIOCHEMICAL CONSTITUENT**

The effect of *Moringa oleifera* leaf meal on various hematological and biochemical parameters in poultry has been extensively studied. Ebenebe *et al.* (2013) reported a significant ( $P > 0.05$ ) increase in packed cell volume and hemoglobin levels in broiler chickens fed with a 10% Moringa leaf diet. Similarly, Divya *et al.* (2014) found that a 1.5% Moringa leaf powder supplement had a hypocholesterolemic effect in broiler chicks. Makanjuola *et al.* (2014) observed no significant changes in serum protein, albumin, globulin, or aspartate aminotransferase levels with the inclusion of 0.2%, 0.4%, and 0.6% MOLM in laying bird diets. Elkloub *et al.* (2015) reported decreases in plasma cholesterol, aspartate aminotransferase, and alanine transaminase in Japanese quail fed 0.2%, 0.4%, and 0.6% MOLM, with increases in total protein and globulin levels at 0.6% and 0.4% inclusion rates, respectively, compared to controls. Further, Hassan *et al.* (2016) found that 0.2% and

0.3% MOLM supplementation in broilers significantly ( $P < 0.05$ ) increased hemoglobin concentration. Abbas *et al.* (2018) demonstrated that 0.25% to 1.0% MOLM inclusion significantly boosted red blood cell counts in broilers, though other metrics like packed cell volume, white blood cell count, and lymphocyte levels showed no significant differences among groups. Hemoglobin was significantly enhanced at 0.75% and 1.0% MOLM inclusion, while albumin-to-globulin ratios were highest at 0.25% and 0.5% MOLM. MOLM supplementation also led to significant reductions in serum cholesterol, low-density lipoprotein (LDL), and very-low-density lipoprotein (VLDL) levels compared to controls. Fouad and Rayes (2019) observed increases in red blood cells, hemoglobin, packed cell volume, white blood cells, total protein, albumin, calcium, and high-density lipoprotein (HDL) in Japanese quail fed with 3g, 5g, and 7g of MOLM per kg of feed, while plasma cholesterol, total lipids, aspartate aminotransferase, alanine transaminase, and glucose levels were decreased. Similarly, Ajantha *et al.* (2020) reported that MOLM-treated broilers exhibited lower serum cholesterol and triglycerides but had significantly higher ( $P < 0.01$ ) high density lipoprotein levels than the control. Sharmin *et al.* (2021) observed that a 1.5% MOLM dietary inclusion significantly reduced serum cholesterol and triglyceride levels in laying birds. Similarly, Garcia *et al.* (2021) found increased serum alkaline phosphatase at 4% MOLM in Japanese quail diets, with lower cholesterol and triglycerides at 6% MOLM. Meel *et al.* (2022) noted significantly higher hemoglobin, packed cell volume, total erythrocyte count, and lymphocyte levels in broilers fed a 1.5% MOLM diet, alongside reduced heterophil counts, heterophil-to-lymphocyte ratios, albumin, globulin, total protein, blood glucose, triglycerides, and creatinine compared to controls.

#### **D) EFFECT ON CARCASS CHARACTERISTICS AND QUALITY**

Zanu *et al.* (2012) observed that the inclusion of *Moringa oleifera* leaf meal (0, 5, 10, and 15%) in broiler diets had no significant impact ( $P > 0.05$ ) on carcass traits. In contrast, Tesfaye *et al.* (2013) reported that adding *Moringa oleifera* leaf meal at 0, 5, 10, 15, and 20% significantly reduced ( $P < 0.05$ ) parameters such as dressed weight, eviscerated weight, breast, thighs, and drumstick weights in broilers. Safa and Tazi (2014) found that supplementing broiler diets with 5% *Moringa oleifera* leaf meal improved carcass yield percentage compared to controls. Similarly, Elkloub *et al.* (2015) indicated that *Moringa oleifera* leaf meal at 0.2, 0.4, or 0.6% significantly reduced abdominal fat in Japanese quail and improved spleen proportion without notable differences from the control group. Karthivashan *et al.*

(2015) found that adding *Moringa* leaf meal extracts at 0, 0.5, 1.0, and 1.5% significantly enhanced ( $P<0.05$ ) dressing percentage in broilers, with 1.0% showing the highest dressing rate. Tijani *et al.* (2016) showed that up to 15% *Moringa oleifera* leaf meal in broiler diets significantly improved ( $P<0.05$ ) dressing percentage, though 10% and 20% levels reduced breast, wing, and abdominal fat. Khalifa *et al.* (2018) showed that 20% MOLM increased heart and liver weights ( $P<0.05$ ), though there were no significant differences in weights of the gizzard, carcass, and edible parts among other treated Japanese quails. Additionally, Fouad and Rayes (2019) found Japanese quails fed 7g MOL/kg diet had the highest carcass weight among treatments. Ajantha *et al.* (2020) reported that MOLM-supplemented groups showed a significant ( $P<0.01$ ) decrease in cholesterol in thigh and breast muscle compared to controls. Baloch *et al.* (2021) observed that diets with 1.25%, 2.5%, and 3.75% MOLM had a significant ( $P<0.05$ ) effect on broiler dressing percentage compared to controls. Furthermore, Varalakshmi *et al.* (2021) indicated that a 3.0% *Moringa oleifera* leaf meal level increased carcass and ready-to-cook yield ( $P<0.05$ ). Harshini *et al.* (2022) reported that adding 0, 5, 10, and 15% *Moringa oleifera* leaf powder to the basal diet increased crude protein and decreased crude fat in meat, though the difference was not significant ( $P>0.05$ ) from the control group. Meat oxidative stability, however, significantly improved ( $P<0.05$ ) with higher inclusion levels in Kadaknath chicken diets.

#### **E) EFFECT ON OXIDATIVE STRESS MARKERS IN SERUM AND MEAT**

Cowan (1999) reported that caffeic and cinnamic acids in *Moringa oleifera* contribute to its antioxidant properties. Saini *et al.* (2014) also highlighted that *Moringa oleifera* is rich in antioxidants, such as ascorbic acid and tocopherol, which support birds in managing stress. According to Rao *et al.* (2019), *Moringa oleifera* reduces lipid peroxidation, thereby lowering stress levels in birds. Similarly, Elkloub *et al.* (2015) found that total antioxidant capacity in plasma was highest in Japanese quail fed 0.6% MOLM, followed by 0.4% and 0.2%, compared to the control. Mousa *et al.* (2016) also found that plasma antioxidant capacity was significantly higher at 0.6% and 0.4% MOLM in Japanese quail during the laying period. Cui *et al.* (2018) observed a gradual increase in total antioxidant capacity, superoxide dismutase, and glutathione peroxidase activities in plasma, with a quadratic decrease in MDA levels in broiler breast muscle with MOL supplementation. Additionally, Khan *et al.* (2021) reported that broilers showed significantly lower stress levels with

*Moringa oleifera* leaf extract supplementation compared to an antibiotic group, indicated by reduced blood MDA concentrations during the finisher phase.

#### **F) EFFECT ON IMMUNE STATUS IN SERUM**

Chollom *et al.* (2012) demonstrated that *Moringa oleifera* seed extract showed strong antiviral effects against Newcastle disease (ND) in ovo. Eze *et al.* (2013) also found that a methanolic extract of *Moringa oleifera* at 200 mg/kg body weight increased ND HI titers in both vaccinated and non-vaccinated chicken groups, recommending it as an immune booster for non-vaccinated birds. Likewise, Liaqat *et al.* (2016) suggested that replacing canola meal with *Moringa oleifera* leaf meal as a plant protein source could enhanced immune response to Newcastle disease and Infectious bursal disease vaccinations, with no effect on growth, organ weights, or blood hematology in broilers. Ramadan (2017) further observed that adding 3%, 5%, and 8% *Moringa oleifera* leaf meal to bird feed significantly increased ( $P < 0.05$ ) antibody titers against ND compared to unsupplemented birds. Similarly, Khan *et al.* (2021) reported a significant ( $P < 0.05$ ) rise in antibody titers against ND and Infectious bronchitis (IB) when *Moringa oleifera* was added to drinking water at 60, 90, and 120 ml/litre. Kumar *et al.* (2021) noted that *Moringa oleifera* leaf extract combined with various levels of ascorbic acid improved antibody titers in response to ND vaccinations, strengthening immune resistance in broilers. Additionally, Kumari *et al.* (2022) found that supplementing *Moringa* leaf extract in drinking water significantly ( $P < 0.05$ ) raised antibody titers in the treated group compared to the control group.

#### **G) EFFECT ON EGG PRODUCTION AND EGG QUALITY**

Kwari *et al.* (2011) observed that including *Moringa oleifera* leaf meal at 1–2% in Vanaraja laying hen diets had no significant effects on feed conversion or egg weight. Similarly, Ebenebe *et al.* (2013) reported that adding MOL had no effect on egg shape index, which relates to eggshell strength and egg grading. Etalem *et al.* (2014) found that the hatchability rate was significantly higher in groups supplemented with 5% *Moringa oleifera* leaf meal than in the control group. Additionally, Makanjuola *et al.* (2014) found that *Moringa oleifera* seed meal at 5% and 7.5% levels was effective in enhancing eggshell thickness in laying hens. Alebachew (2016) noted that *Moringa oleifera* leaves, rich in zinc and vitamin E, could enhance hatchability and improve egg quality through antioxidants, essential oils, minerals (Ca, Mg, K, Se, P, Zn), and vitamins (A, C, D, K, and E). Riry *et al.* (2016) also observed a reduction in feed intake when Japanese quails were fed a 5% *Moringa oleifera*

seed meal diet compared to controls. Mousa *et al.* (2017) found that supplementing Japanese quails' diets with 0.4% MOLM improved hatchability of all eggs, while 0.6% MOLM improved hatchability of fertile eggs. Further, Abdelnour *et al.* (2018) suggested that up to 10% Moringa leaves had no negative impact on egg production in laying birds; however, levels over 10% could be detrimental due to anti-nutritional factors, leaf dustiness, and reduced energy and protein digestibility. Tesfaye *et al.* (2018) highlighted that 5% *Moringa oleifera* leaf meal could serve as a feed additive, enhancing product quality in the poultry sector. Akinola and Ovotu (2018) found that high-density lipoprotein (HDL) in eggs increased significantly ( $P<0.05$ ) with 1% and 1.5% MOLM, positively affecting egg cholesterol levels by week one. Siti *et al.* (2019) observed that diets containing 2-6% Moringa leaf powder enhanced yolk color in laying birds ( $P<0.05$ ). Similarly, Ashour *et al.* (2020) reported that *Moringa oleifera* leaf diets did not affect feed conversion ratio, egg weight, fertility, hatchability, egg and yolk index, or Haugh units, although egg production, mass, shell thickness, and hatchability improved with *Moringa oleifera* seed meal supplementation. Moreover, Atuahene *et al.* (2020) found no adverse effects on egg quality, fertility, or hatchability of quail eggs when soybean meal was partially replaced by *Moringa oleifera* leaf meal up to 15%, though yolk color was enhanced in Japanese quails. Talukdar *et al.* (2020) observed that Japanese quail egg traits, including weight, shape index, albumen index, and yolk index, showed no significant changes ( $P>0.05$ ), except for yolk color, which increased significantly ( $P<0.05$ ) in the 1% MOLM group. Organoleptic qualities, such as color, flavor, texture, juiciness, and acceptability, remained consistent ( $P>0.05$ ) across groups. Farhana Sharmin *et al.* (2021) also found that egg quality, weight, length, and width were unaffected by adding 1.5% MOL to the diet, though cholesterol levels significantly dropped ( $P<0.05$ ) at 1.5% MOLM. In addition, 1.0-1.5% MOLM supplementation increased omega-3 fatty acids in the yolk. Finally, Garcia *et al.* (2021) noted that adding Moringa reduced feed intake up to 1.2% and increased egg weight up to 3.80% with a linear increase in yolk weight and intensified yolk color with higher Moringa levels.

## **7. CONCLUSION**

*Moringa oleifera*, often called the “Miracle Tree,” offers numerous benefits for sustainable poultry production. Its leaf meal, rich in nutrients, can be seen as an invaluable, cost-effective natural resource.

## **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Authors hereby declare that no generative AI technologies such as Large Language Models (Chat GPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

## **COMPETING INTERESTS**

Author has declared that no competing interests exist.

## **REFERENCES**

Abbas, R. J., Ali, N. A. L., Alkassar, A. M., & Jameel, Y. J. (2018). Haematological and biochemical indices of broiler chicks fed at different levels of *Moringa oleifera* leaf meal. *Biochemical & Cellular Archives*, 18(2), 1931-1936.

Abdelnour, S. A., Abd El-Hack, M. E., & Ragni, M. (2018). The efficacy of high-protein tropical forages as alternative protein sources for chickens: A review. *Agriculture*, 8(6), 86.

Ajantha, A., Kathirvelan, C., Purushothaman, M. R., & Visha, P. (2020). Effect of *Moringa oleifera* leaf meal supplementation in broiler chicken on serum and muscle lipid profile. *Journal of Pharmacognosy and Phytochemistry*, 9(2S), 464-466.

Akinola, L. A. F., & Ekine, O. A. (2018). Evaluation of commercial layer feeds and their impact on performance and egg quality. *Nigerian Journal of Animal Science*, 20(2), 222-231.

Alebachew, W., Tesfaye, E., & Tamir, B. (2016). Effects of feeding different dietary levels of *Moringa oleifera* leaf meal on egg production, fertility and hatchability of dual purpose Koekoek hens. *Middle-East Journal of Scientific Research*, 24(9), 2909-2920.

Amaglo, N. K., Bennett, R. N., Curto, R. B. L., Rosa, E. A., Turco, V. L., Giuffrida, A., ... & Timpo, G. M. (2010). Profiling selected phytochemicals and nutrients in different tissues of the multipurpose tree *Moringa oleifera* L., grown in Ghana. *Food Chemistry*, 122(4), 1047-1054.

Anhwange, B. A., Ajibola, V. O., & Oniye, S. J. (2004). Chemical studies of the seeds of *Moringa oleifera* (Lam) and *Detarium microcarpum* (Guill and Sperr). *Journal of Biological Sciences*, 4, 711-715.

Anjorin, T. S., Ikokoh, P., & Okolo, S. (2010). Mineral composition of *Moringa oleifera* leaves, pods and seeds from two regions in Abuja, Nigeria. *International Journal of Agriculture and Biology*, 12: 431-434.

Ashong, J. O., & Brown, D. L. (2011). Safety and efficacy of powder for growing poultry. *Journal of Animal science*, 89, 84.

Ashour, E. A., El-Kholy, M. S., Alagawany, M., Abd El-Hack, M. E., Mohamed, L. A., Taha, A. E., & Tufarelli, V. (2020). Effect of dietary supplementation with *Moringa oleifera* leaves and/or seeds powder on production, egg characteristics, hatchability and blood chemistry of laying Japanese quails. *Sustainability*, 12(6), 2463-2469.

Atuahene, C. C., Opoku, O., Benante, V., Quaye, B., Adu, M. A., Tamathey, B. B., & Kumah, G. (2020). Effects of *Moringa oleifera* Leaf Meal on Fertility, Egg Quality and Hatchability of Japanese Quails. *Ghanaian Journal of Animal Science*, 11(1), 75-82.

Ayssiwede, S. B., Dieng, A., Bello, H., Chrysostome, C. A. A. M., Hane, M. B., Mankor, A., & Missohou, A. (2011). Effects of *Moringa oleifera* (Lam.) leaves meal incorporation in diets on growth performances, carcass characteristics and economics results of growing indigenous Senegal chickens. *Pakistan Journal of Nutrition*, 10(12), 1132-1145.

Baloch, F. N., Baloch, H. N., & Mughal, G. A. (2021). Replacement of sunflower meal with *Moringa oleifera* leaves on growth performance of broiler. *Approaches in Poultry Dairy & Veterinary Sciences*, 8(2), 756-762.

Bhargave, A., Pandey, I., Nama, K. S., & Pandey, M. (2015). *Moringa oleifera* Lam.—Sanjana (Horseradish Tree)—A miracle food plant with multipurpose uses in Rajasthan-India—an overview. *Int J Pure App Biosci*, 3(6), 237-248.

Chollom, S. C., Agada, G. O. A., Gotep, J. G., Mwankon, S. E., Dus, P. C., Bot, Y. S., ... & Okwori, A. E. J. (2012). Investigation of aqueous extract of *Moringa oleifera* lam seed for antiviral activity against newcastle disease virus in ovo. *J. Med. Plants Res*, 6(22), 3870-3875.

Cowan, M. M. (1999). Plant products as antimicrobial agents. *Clinical microbiology reviews*, 12(4), 564-582.

- Cui, Y. M., Wang, J., Lu, W., Zhang, H. J., Wu, S. G., & Qi, G. H. (2018). Effect of dietary supplementation with *Moringa oleifera* leaf on performance, meat quality, and oxidative stability of meat in broilers. *Poultry Science*, 97(8), 2836-2844.
- Dhakar, R. C., Maurya, S. D., Pooniya, B. K., Bairwa, N., & Gupta, M. (2011). Moringa: The herbal gold to combat malnutrition. *Moringa: The Herbal Gold To Combat Malnutrition*.
- Divya, D., Mandal, A. B., Biswas, A., Yadav, A. S., & Biswas, A. K. (2014). Effect of dietary *Moringa oleifera* leaves powder on growth performance, blood chemistry, meat quality and gut microflora of broiler chicks. *Animal Nutrition and Feed Technology*, 14(2), 349-357.
- Ebenebe, C. I., Anigbogu, C. C., Anizoba, M. A., & Ufele, A. N. (2013). Effect of various levels of *Moringa leaf* meal on the egg quality of ISA Brown breed of layers. *Advances in Life Science and Technology*, 14(1), 35-49.
- Elkloub, K., Moustafa, M. E. L., Riry, F. H., Mousa, M. A. M., Hanan, A. H., & Youssef, S. F. (2015). Effect of using *Moringa oleifera* leaf meal on performance of Japanese quail. *Egypt. Poult. Sci. J*, 35, 1095-1108.
- El-Tazi, S. M. (2014). Effect of feeding different levels of *Moringa oleifera* leaf meal on the performance and carcass quality of broiler chicks. *International Journal of Scientific Research*, 3(5), 147-151.
- Tesfaye, E. B., Anmut, G. M., Urge, M. L., & Dessie, T. A. (2014). Cassava root chips and *Moringa oleifera* leaf meal as alternative feed ingredients in the layer ration. *Journal of Applied Poultry Research*, 23(4), 614-624.
- Eze, D. C., Okwor, E. C., Okoye, J. A., & Onah, D. N. (2013). Immunologic effects of *Moringa oleifera* methanolic leaf extract in chickens infected with Newcastle diseasevirus (kudu 113) strain.
- Fahey, J. W. (2005). *Moringa oleifera*: a review of the medical evidence for its nutritional, therapeutic, and prophylactic properties. Part 1. *Trees for life Journal*, 1(5), 1-15.

Sharmin, F., Sarker, M. S. K., Sarker, N. R., & Faruque, S. (2021). Dietary effect of *Moringa oleifera* on native laying hens' egg quality, cholesterol and fatty-acid profile. *Italian Journal of Animal Science*, 20(1), 1544-1553.

Su, B., & Chen, X. (2020). Current status and potential of *Moringa oleifera* leaf as an alternative protein source for animal feeds. *Frontiers in veterinary science*, 7, 53.

Ahmed, W. F., & El-Rayes, T. K. (2019). Effect of using *Moringa oleifera* leaves on productive performance and some physiological parameters of japanese quail. *Egyptian Poultry Science Journal*, 39(1), 193-205.

Fuglie, L. J. (1999). The miracle tree: *Moringa oleifera*, natural nutrition for the tropics.

Subha Ganguly, S. G. (2013). Indian Ayurvedic and traditional medicinal implications of indigenously available plants, herbs and fruits: a review.

Garcia, R. G., Gandra, E. R. S., Burbarelli, M. F. C., Valentim, J. K., Felix, G. A., Lopes, B. A., & Caldara, F. R. (2021). Moringa Oleifera: an alternative ingredient to improve the egg quality of Japanese quail. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia*, 73(03), 721-732.

Harshini, A., Tomar, S., Yadav, A. S., & Awasthi, M. G. (2022). Effect of *Moringa oleifera* leaf powder feeding on the growth performance and meat quality of Kadaknath chicken. *Indian Journal of Animal Nutrition*, 39(4), 486-501.

Hassan, H. M. A., El-Moniary, M. M., Hamouda, Y., El-Daly, E. F., Youssef, A. W., & Abd El-Azeem, N. A. (2016). Effect of different levels of *Moringa oleifera* leaves meal on productive performance, carcass characteristics and some blood parameters of broiler chicks reared under heat stress conditions. *Asian Journal of Animal and Veterinary Advances*, 11(1), 60-66.

Jung, I. L. (2014). Soluble extract from *Moringa oleifera* leaves with a new anticancer activity. *PloS one*, 9(4), e95492.

Kakengi, A. M. V., Shem, M. N., Sarwatt, S. V., & Fujihara, T. (2005). Can *Moringa oleifera* be used as a protein supplement for ruminants?. *Asian-Australasian journal of animal sciences*, 18(1), 42-47.

- Karthivashan, G., Arulselvan, P., Alimon, A. R., Safinar Ismail, I., & Fakurazi, S. (2015). Competing role of bioactive constituents in *Moringa oleifera* extract and conventional nutrition feed on the performance of Cobb 500 broilers. *BioMed research international*, 2015(1), 970398.
- Khalifa, W. H., Abusinaa, G. E., Elmallah, G. M., Yassein, S. A., Elghamry, A. A., & Maghraby, N. A. (2018). Effect of using different levels of *Moringa oleifera* meal in the diets of Japanese quails on their productive performance. *Egyptian Journal of Nutrition and Feeds*, 21(2), 529-535.
- Khan, R. U. (2011). Antioxidants and poultry semen quality. *World's poultry science journal*, 67(2), 297-308.
- Kholif, A. E., Morsy, T. A., Gouda, G. A., Anele, U. Y., & Galyean, M. L. (2016). Effect of feeding diets with processed *Moringa oleifera* meal as protein source in lactating Anglo-Nubian goats. *Animal Feed Science and Technology*, 217, 45-55.
- Kumar, R., Kumar, K., Kumar, A., Kumar, S., Singh, P. K., Sinha, R. R. K., & Moni, C. (2021). Nutritional and physiological responses of broiler chicken to the dietary supplementation of *Moringa oleifera* aqueous leaf extract and ascorbic acid in tropics. *Tropical Animal Health and Production*, 53, 1-7.
- Kumari, S., Kumar, P., Kumar, M., Kumar, S., Anjay, & Kaushik, P. (2022). Effect of Moringa leaf aqueous extract feeding on humoral response in chicks. *Journal of Agricultural Research*, 9(3), 275-277.
- Kwari, I. D., Diarra, S. S., Saleh, B., Bova, P. R., Ramat, O. A., & Tochukwu, D. (2011). Growth, hematology and serology of broiler chickens fed different cultivars of sorghum as replacement for maize in the semi-arid zone of Nigeria. *International Journal of Poultry Science*, 10(8), 608-612.
- Liaqat, S., Mahmood, S., Ahmad, S., Kamran, Z., & Koutoulis, K. C. (2016). Replacement of canola meal with *Moringa oleifera* leaf powder affects performance and immune response in broilers. *Journal of Applied Poultry Research*, 25(3), 352-358.

- Mahanta, J. D., Borgohain, B., Sarma, M., Sapkota, D., & Hussain, J. (2017). Effect of dietary supplementation of herbal growth promoter on performance of commercial broiler chicken. *Indian Journal of Animal Research*, 51(6), 1097-1100.
- Makanjuola, B. A., Obi, O. O., Olorunghohunmi, T. O., Morakinyo, O. A., Oladele-Bukola, M. O., & Boladuro, B. A. (2014). Effect of *Moringa oleifera* leaf meal as a substitute for antibiotics on the performance and blood parameters of broiler chickens. *Livestock Research for Rural Development*, 26(8), 144.
- Mbikay, M. (2012). Therapeutic potential of *Moringa oleifera* leaves in chronic hyperglycemia and dyslipidemia: a review. *Frontiers in pharmacology*, 3, 24.
- Meel, M. S., Sharma, T., Dhuria, R. K., Joshi, M., Shende, K. A., Kumari, M., & Mishra, G. (2022). Influence of dietary inclusion of *Moringa oleifera* leaf meal on blood chemistry of broiler chicks. *Scientist*, 1(3), 4300-4305.
- Mendieta-Araica, B., Spörndly, R., Reyes-Sánchez, N., & Spörndly, E. (2011). *Moringa (Moringa oleifera)* leaf meal as a source of protein in locally produced concentrates for dairy cows fed low protein diets in tropical areas. *Livestock Science*, 137(1-3), 10-17.
- Mousa, M. A. M., Moustafa, K. E. M., Shata, R. F., Alghonimy, H. A., & Youssef, S. F. (2017). Effect of using *Moringa oleifera* leaf meal as feed additives on Japanese quail during lying period. *Egyptian Journal of Nutrition and Feeds*, 20(2 Special), 203-212.
- Moustafa, K. E. M. (2016). Effect of using germinated *Moringa oleifera* seeds on Japanese quail growth performance. *Egyptian Poultry Science Journal*, 36(2), 561-571.
- Nikkon, F. (2003). In vitro Antimicrobial Activity of the Compound Isolated from Chloroform Extract of *Moringa oleifera* Lam. Farjana Nikkon, Zahangir Alam Saud, M. Habibur Rahman and" Md. Ekramul Haque Department of Biochemistry and Molecular Biology," Department of Pharmacy. *Pakistan Journal of Biological Sciences*, 6(22), 1888-1890.
- Nkukwana, T. T., Muchenje, V., Masika, P. J., & Mushonga, B. (2015). Intestinal morphology, digestive organ size and digesta pH of broiler chickens fed diets supplemented with or without *Moringa oleifera* leaf meal. *South African Journal of Animal Science*, 45(4), 362-370.

Olugbemi, T. S., Mutayoba, S. K., & Lekule, F. P. (2010). Effect of Moringa (*Moringa oleifera*) inclusion in cassava based diets fed to broiler chickens. *International Journal of Poultry Science*, 9(4), 363-367.

Pari, L., & Kumar, N. A. (2002). Hepatoprotective activity of *Moringa oleifera* on antitubercular drug-induced liver damage in rats. *Journal of Medicinal Food*, 5(3), 171-177.

Patel, J. P., Gami, B., Patel, K., & Solanki, R. (2011). Antibacterial activity of methanolic and acetone extract of some medicinal plants used in Indian folklore. *International Journal of Phytomedicine*, 3(2), 261.

Ramadan, S. G. (2017). Impact of supplementation of *Moringa oleifera* in diet of broiler chicks on their behavior, welfare, performance and immune responses.

Rao, S. R., Raju, M. V. L. N., Prakash, B., Rajkumar, U., & Reddy, E. P. K. (2018). Effect of supplementing moringa (*Moringa oleifera*) leaf meal and pomegranate (*Punica granatum*) peel meal on performance, carcass attributes, immune and antioxidant responses in broiler chickens. *Animal Production Science*, 59(2), 288-294.

Riry, F. H., Elkloub, K., Moustafa, M. E. L., Mousa, M. A. M., Youssef, S. F., & Hanan, A. H. (2016, November). Effect of using *Moringa oleifera* seed meal on Japanese quail performance during growing period. In *Proceedings of the 9th International Poultry Conference, Hurghada, Egypt* (pp. 7-10).

Saini, R. K., Shetty, N. P., Prakash, M., & Giridhar, P. (2014). Effect of dehydration methods on retention of carotenoids, tocopherols, ascorbic acid and antioxidant activity in *Moringa oleifera* leaves and preparation of a RTE product. *Journal of food science and technology*, 51, 2176-2182.

Sánchez, N. R., Ledin, S., & Ledin, I. (2006). Biomass production and chemical composition of *Moringa oleifera* under different management regimes in Nicaragua. *Agroforestry Systems*, 66, 231-242.

Siti, N. W., Bidura, I. G. N. G., Mayuni, S. N., Suasta, I. M., & Utami, I. A. P. (2019). Effect of *Moringa oleifera* leaf powder in diets on feed digestibility and external egg quality characteristics in laying hens. *International Journal of Fauna and Biological Studies*, 6(4), 113-118.

- Su, B., & Chen, X. (2020). Current status and potential of *Moringa oleifera* leaf as an alternative protein source for animal feeds. *Frontiers in veterinary science*, 7, 53.
- Talreja, T. (2010). Screening of crude extract of flavonoids of *Moringa oleifera* against bacteria and fungal pathogen. *Journal of Phytochemistry*, 2(11).
- Talukdar, A., Choudhury, D., Kalita, K. P., & Saikia, R. (2020). Effect of using *Moringa oleifera* (drumstick) leaf meal on performance of Japanese quail. *JEZS*, 8(2), 1485-1490.
- Tesfaye, E., Alebachew, W., & Tamir, B. (2018). Performance of koekoek chicken fed with different levels of *Moringa oleifera* leaf meal. *Journal of Animal Science Research*, 2(3).
- Tesfaye, E., Animut, G., Urge, M., & Dessie, T. (2013). *Moringa olifera* leaf meal as an alternative protein feed ingredient in broiler ration. *International Journal of Poultry Science*, 12(5), 289-297.
- Thirumalaisamy, G., Muralidharan, J., Senthilkumar, S., Hema Sayee, R., & Priyadharsini, M. (2016). Cost-effective feeding of poultry. *International Journal of Science, Environment and Technology*, 5(6), 3997-4005.
- Tijani, L. A., Akanji, A. M., Agbalaya, K., & Onigemo, M. (2016). Effects of *Moringa oleifera* leaf meal on performance, nutrient digestibility and carcass quality of broiler chickens. *Applied Tropical Agriculture*, 21(1), 46-53.
- Varalakshmi, G., Kishore, K. R., Kumar, D. S., & Sessaiah, C. V. (2021). Growth and carcass characteristics of Japanese quails fed diets containing varying levels of *Moringa* leaf meal. *Indian Journal of Animal Nutrition*, 38(2), 207-212.
- Zanu, H. K., Asiedu, P., Tampuori, M., Abada, M., & Asante, I. (2012). Possibilities of using *Moringa (Moringa oleifera)* leaf meal as a partial substitute for fishmeal in broiler chickens diets. *Online Journal of Animal and Feed Research*, 2(1), 70-75.

UNDER PEER REVIEW